



INSTITUTE OF MATHEMATICS

THE CZECH ACADEMY OF SCIENCES

**Recursive functions and existentially
closed structures**

Emil Jeřábek

Preprint No. 65-2017

PRAHA 2017

Recursive functions and existentially closed structures

Emil Jeřábek*

Institute of Mathematics of the Czech Academy of Sciences

Žitná 25, 115 67 Praha 1, Czech Republic, email: jerabek@math.cas.cz

October 26, 2017

Abstract

The purpose of this paper is to clarify the relationship between various conditions implying essential undecidability: our main result is that there exists a theory T in which all partially recursive functions are representable, yet T does not interpret Robinson’s theory R . To this end, we borrow tools from model theory—specifically, we investigate model-theoretic properties of the model completion of the empty theory in a language with function symbols. We obtain a certain characterization of $\exists\forall$ theories interpretable in existential theories in the process.

1 Introduction

First-order theories studied by logicians may be broadly divided in two classes. One class comprises theories of “arithmetical strength”, such as various fragments and extensions of Peano arithmetic, or set theories. They are distinguished by their great expressive power that, on the one hand, allows them to work with all kinds of objects from mathematical practice in suitable encoding (indeed, some of these theories are designed to serve as foundation for all of mathematics, e.g., ZFC), and on the other hand, makes them subject to Gödel’s incompleteness theorems and related phenomena. The other class are theories of “tame” structures, for example algebraically closed or real closed fields, vector spaces, generic structures such as the random graph, etc. These theories have low expressive power (often manifested in classification of definable sets stemming from partial quantifier elimination), and consequently their models have a manageable structure of geometric nature. Tame theories tend to be decidable.

The borderline between arithmetical and tame theories is not sharply demarcated, but one typical feature of arithmetical theories is their *essential undecidability*, meaning that all consistent extensions of the theory are undecidable. This notion was isolated by Tarski, Mostowski, and Robinson [9]. This classic monograph also includes convenient methods for proving essential undecidability of a theory T , which can be viewed as stand-alone properties implying essential undecidability. In order of increasing strength, these are:

- T can represent all partially recursive functions (prf; see below for a precise definition).
- T can interpret Robinson’s theory R .
- T can interpret Robinson’s arithmetic Q , or equivalently, the adjunctive set theory.

*Supported by grant IAA100190902 of GA AV ČR, Center of Excellence CE-ITI under the grant P202/12/G061 of GA ČR, and RVO: 67985840.

An even stronger condition is that of being an *sequential theory* [7, 11].

Recall that Robinson’s R is, essentially, a theory axiomatizing the true Σ_1 sentences of the standard model of arithmetic \mathbb{N} ; while it is in some ways less convenient to work with than the better-known arithmetic Q (e.g., R is not finitely axiomatizable), it is distinguished by its interpretability properties—see Visser [12].

The above-mentioned conditions on theories form an increasing chain. For most of the inclusions in this chain, it is clear (or at least, reasonably well known) that the inclusions are strict: in particular, there are theories interpreting Q that are not sequential (in fact, Q itself is such a theory [10]), R does not interpret Q (as Q is a finitely axiomatized theory with no finite model, whereas R is locally finitely satisfiable), and there are essentially undecidable recursively axiomatized theories that do not represent prf . However, one of these inclusions is not as easy to resolve, leading to the question that motivated this paper:

Question 1.1 *If a theory represents all partial recursive functions, does it interpret Robinson’s theory R ?*

This may look plausible at first sight: R is a very weak theory that only fixes the values of elementary arithmetic operations on standard natural numbers, and requires virtually nothing else from the rest of the model. Now, the definition of representability of prf does provide for natural number constants and definable functions on them that behave like elementary arithmetic operations as these operations are prf , so everything seems to be in order.

Despite this, the answer turns out to be negative. The devil is in the “virtually nothing else”: R does, after all, involve universally quantified conditions that may look innocuous (in our favourite formulation of R , these universal quantifiers are bounded by a constant, hence ostensibly “finite”), but actually turn out to be crucially important. Using Visser’s [12] characterization, R interprets nontrivial universal theories such as the theory of infinite discrete linear order. In contrast, prf can be represented in a theory axiomatized purely by quantifier-free sentences, with no universal quantifiers lurking behind.

We are going to prove that consistent theories with quantifier-free—or even existential—axioms cannot interpret infinite linear orders and a couple of similar universal theories, and a fortiori, cannot interpret R . This is not easy to work out directly: the weakness of existential theories—which should intuitively be the reason for nonexistence of such interpretations—backfires in that we have absolutely no control over the complexity of formulas that make up potential interpretations, and over the sets they define in models.

Our strategy to solve this problem is to consistently extend the interpreting theory to a theory with quantifier elimination, using the fact that the empty theory in an arbitrary language L has a model completion (which we denote EC_L , being the theory of existentially closed L -structures). This fact is well known for relational languages, in which case EC_L is the theory of the “random L -structure”. However, we need it for languages with function symbols, which case is mostly neglected in common literature, though the existence of EC_L was proved in full generality already by Winkler [14].

It follows that if a theory is interpretable in a consistent quantifier-free or existential theory, it is weakly interpretable in EC_L for some L , and the interpretation can be taken quantifier-free. In order to see that this is heading in the right direction, we establish a converse result: if an $\exists\forall$ theory is weakly interpretable in EC_L , it is interpretable in a quantifier-free theory.

We proceed to prove that EC_L does not, actually, weakly interpret various theories of interest. At this point, we are heading further and further into model theory, having left the realm of arithmetical theories. It turns out that our non-interpretability results can be naturally expressed in the language of *classification theory*. Arising through the work of Shelah [8], classification theory

studies the landscape of “dividing lines” between tame and wild theories, and their structural consequences. Many dividing lines have the following form: a theory is wild if it has a model that contains a certain complex combinatorial arrangement. Usually, conditions of this form can be reformulated as (weak) interpretability of a specific $\exists\forall$ theory. For a concrete example, a theory T has the *strict order property* (SOP) if there exists a model $M \models T$, a formula $\varphi(\bar{x}, \bar{y})$, and tuples $\bar{a}_n \in M$ for $n \in \mathbb{N}$ such that φ defines in M a strict partial order, and $\varphi(\bar{a}_n, \bar{a}_m)$ whenever $n < m$. Otherwise, T is said to have the *no-strict-order property* (NSOP).

We observe that theories that can represent recursive functions, as well as consistent extensions of EC_L for sufficiently rich languages L , are moderately wild in that they always have the tree property TP_2 . However, we will prove that EC_L (for arbitrary L) has certain tameness properties: specifically, it has the no-strong-order property $NSOP_3$ (which implies NSOP), and it has elimination of infinity. Using a characterization of $NSOP_1$ theories by Chernikov and Ramsey [2], we show that it even has the $NSOP_1$ property. On the other hand, theories interpreting R are firmly on the wild side of all generally considered dividing lines.

For completeness, the paper also includes discussion of basic model-theoretic properties of EC_L in the appendix.

2 Preliminaries

Let us first agree on a few bits of general notation. We will use \mathbb{N} and ω more or less interchangeably to denote the set of nonnegative integers; \mathbb{N} may also denote the standard model of arithmetic $\langle \mathbb{N}, 0, \text{succ}, +, \cdot, < \rangle$. We denote sequences by angle brackets, and consider them indexed starting from 0; tuples of finite-but-unspecified length will be denoted by placing a bar over a variable name, so that \bar{x} may stand for the n -tuple $\langle x_0, \dots, x_{n-1} \rangle$.

We will write $F: X \dashrightarrow Y$ to denote that F is a partial function from X to Y . (We use this notation in the context of partial recursive functions, so virtually always we will have $X = \mathbb{N}^k$, $Y = \mathbb{N}$.)

The notation $t \doteq s$ means that t and s are syntactically identical terms; we may also apply it to formulas and other syntactic objects.

2.1 Theories and interpretations

In this paper, a *language* consists of an arbitrary number of relation and function symbols of arbitrary finite arity (including 0: nullary functions are constants, nullary relations are propositional variables). A *theory* is a deductively closed set of sentences in a particular language. A theory in language L is also called an L -theory. We will often consider theories specified by a set of axioms, in which case the theory is taken to be their deductive closure; we will frequently omit outer universal quantifiers from axioms. We will generally employ a form of first-order logic that allows *empty models*.

Many considerations in this paper revolve around the notion of *interpretation* of one theory in another, so we need to be somewhat specific about its meaning. However, since a precise technical definition of interpretations would get quite lengthy, we advise the reader to consult e.g. Visser [12, §2] for the details if necessary; we will only indicate the main distinctive features below.

Let T be a theory in a language L_T , and S a theory in a language L_S . In its most simple form, a *translation* I of language L_T into language L_S is specified by:

- An L_S -formula $\delta_I(x)$ denoting the domain of I .
- For each relation symbol R of L_T , as well as the equality relation $=$, an L_S -formula R_I of the same arity.

- For each function symbol F of L_T of arity k , an L_S -formula F_I of arity $k + 1$.

If φ is an L_T -formula, its I -translation φ^I is an L_S -formula constructed as follows: we rewrite the formula in an equivalent way so that function symbols only occur in atomic subformulas of the form $F(\bar{x}) = y$, where x_i, y are variables; then we replace each such atomic formula with $F_I(\bar{x}, y)$, we replace each atomic formula of the form $R(\bar{x})$ with $R_I(\bar{x})$, and we restrict all quantifiers and free variables to objects satisfying δ_I . We take care to rename bound variables to avoid variable capture during the process.

A translation I of L_T into L_S is an *interpretation* of T in S if S proves:

- For each function symbol F of L_T , the formula expressing that F_I is total on δ_I :

$$\forall x_0, \dots, x_{k-1} (\delta_I(x_0) \wedge \dots \wedge \delta_I(x_{k-1}) \rightarrow \exists y (\delta_I(y) \wedge F_I(\bar{x}, y))) \quad (1)$$

- The I -translations of all axioms of T , and axioms of equality.

It follows that S proves the I -translations of all sentences provable in T .

The simplified picture of translations and interpretations above actually describes only *one-dimensional*, *parameter-free*, and *one-piece* translations. In the full generality, we allow the following:

- Translations may be *multi-dimensional*. That is, we use n -tuples of L_S -objects to represent L_T -objects (where n is a fixed natural number, called the dimension of the translation): thus, δ_I has n free variables, R_I has kn free variables for a k -ary relation $R \in L_T$, and similarly for functions; and when constructing φ^I , each quantifier is replaced with a block of n quantifiers.
- Translations may use *parameters*. This means that the formulas δ_I , R_I , and F_I may include parameter variables \bar{w} that are assumed distinct from any proper variables used in the target formulas, and the specification of I includes an L_S -formula $\pi_I(\bar{w})$ that describes which parameters are admissible. Parameters carry through the translation unchanged, so they appear as free variables in φ^I . The definition of interpretation is modified so that S proves $\forall \bar{w} (\pi_I(\bar{w}) \rightarrow \varphi^I(\bar{w}))$ for each axiom φ , and likewise for (1).
- Translations may be *piece-wise*: the interpreted domain of L_T -objects may be stitched together from finitely many pieces (possibly of different dimensions, and possibly overlapping). Each piece has its own δ_I formula, there is a separate R_I formula for each choice of a sequence of pieces for the arguments of R , etc.

A translation I is called *unrelativized* if, on each piece, $\delta_I(\bar{x})$ is a tautologically true formula, and it has *absolute equality* if, on each piece, $\bar{x} =_I \bar{y}$ is the formula $\bigwedge_i x_i = y_i$.

Under suitable conditions, we do not need the full generality of interpretations:

- Assume that S proves the existence of at least two distinct objects. Then whenever T has an interpretation in S , it also has a one-piece interpretation. (The new interpretation may have larger dimension, but needs no extra parameters.) This can be achieved by using the pattern of equalities on an extra tuple of variables to distinguish pieces. For this reason, we will mostly think of interpretations as one-piece, to avoid unnecessary technical baggage.
- If T has a definable object, then an interpretation of T in S may be converted to an unrelativized interpretation by “equating” tuples outside the original domain with the definable object. If we do not mind using extra parameters, the same can be achieved even if T just proves the existence of at least one object. This construction may not be always desirable, hence relativized interpretations will remain the norm for us.

- A theory S has (non-functional) *pairing* if there is a formula $\pi(x, y, z)$ such that S proves

$$\begin{aligned} \pi(x, y, z) \wedge \pi(x', y', z) \rightarrow x = x' \wedge y = y' \\ \forall x \forall y \exists z \pi(x, y, z). \end{aligned}$$

If T has an interpretation in a theory S with pairing, it also has a one-dimensional interpretation, as we can use single elements to code tuples.

If I_1 is a translation of language L_1 into L_0 , and I_2 a translation of language L_2 into L_1 , the *composition* $I_1 \circ I_2$ is a translation of L_2 into L_0 , and it is defined in an expected way. Note that if I_1 is an interpretation of a theory T_1 in T_0 , and I_2 an interpretation of T_2 in T_1 , then $I_1 \circ I_2$ is an interpretation of T_2 in T_0 .

Let T and S be theories. Some variants on the notion of interpretation of T in S are:

- A *weak interpretation* of T in S is an interpretation of T in a consistent extension of S (in the same language as S), or equivalently, in a completion of S .
- A *cointerpretation* of T in S is a translation I of language L_S into L_T (sic!) such that $T \vdash \varphi^I$ implies $S \vdash \varphi$ for every L_S -sentence φ .
- A *faithful interpretation* of T in S is an interpretation of T in S that is at the same time a cointerpretation of S in T .

A theory T is *interpretable* (*weakly interpretable*, *cointerpretable*) in a theory S if there exists an interpretation (weak interpretation, cointerpretation, resp.) of T in S .

If T and S are complete theories, a translation I of L_T in L_S is an interpretation of T in S iff it is a weak interpretation iff (assuming I is parameter-free) it is a cointerpretation of S in T .

Lemma 2.1 *If I is a weak interpretation of U in T , and J a cointerpretation of S in T , then $J \circ I$ is a weak interpretation of U in S . \square*

An interpretation I of T in S , as defined, is a syntactic transformation of formulas provable in T into formulas provable in S . However, it can be also viewed semantically: it provides a uniform way of building “internally definable” models of T out of models of S .

Assume first I is a parameter-free one-piece interpretation with absolute equality, and let $M \models S$. We construct a model $M^I \models T$ as follows: if I is n -dimensional, the domain of M^I is $\delta_I(M) = \{\bar{a} \in M^n : M \models \delta_I(\bar{a})\}$; a k -ary relation symbol $R \in L_T$ is realized in M^I by $\{\langle \bar{a}_0, \dots, \bar{a}_{k-1} \rangle \in \delta_I(M)^k : M \models R_I(\bar{a}_0, \dots, \bar{a}_{k-1})\}$, and similarly, a k -ary function symbol $F \in L_T$ is realized by the function whose graph is the subset of $(\delta_I(M))^{k+1}$ defined in M by the formula F_I .

Next, if I does not have absolute equality, we build the structure as before, and let M^I be its quotient by the binary relation defined on it by the formula $=_I$; this relation is in fact a congruence, as S proves the translations of equality axioms.

If I is a piece-wise interpretation, we construct the domain of M^I as the disjoint union of the finitely many pieces, each defined as above; we define relations and functions in the appropriate way.

Finally, if I is an interpretation with parameters, we will not obtain a single model M^I , but one model for each choice of parameters: that is, if \bar{a} is a tuple such that $M \models \pi_I(\bar{a})$, then $M^{I, \bar{a}}$ is a model of T built from the expanded structure $\langle M, \bar{a} \rangle$ by the procedure above.

2.2 Representation of recursive functions

The notion of representable¹ predicates and functions in first-order theories was introduced in [9]. We summarize it below, with a few inessential modifications. (Warning: we are going to relax the definition a bit later in this section.)

Definition 2.2 Let T be a theory in a language L , and $\sigma = \{\underline{n} : n \in \mathbb{N}\}$ a fixed *sequence of numerals*: i.e., a sequence of closed terms \underline{n} such that

$$T \vdash \underline{n} \neq \underline{m}$$

for $n, m \in \mathbb{N}$, $n \neq m$.

A recursive predicate (rp) $P \subseteq \mathbb{N}^k$ is *represented* in T w.r.t. σ by a formula $\varphi(x_0, \dots, x_{k-1})$ if

$$\begin{aligned} \langle n_0, \dots, n_{k-1} \rangle \in P &\implies T \vdash \varphi(\underline{n}_0, \dots, \underline{n}_{k-1}), \\ \langle n_0, \dots, n_{k-1} \rangle \notin P &\implies T \vdash \neg\varphi(\underline{n}_0, \dots, \underline{n}_{k-1}) \end{aligned}$$

for all $n_0, \dots, n_{k-1} \in \mathbb{N}$.

A partial recursive function (prf) $F: \mathbb{N}^k \rightarrow \mathbb{N}$ is *represented* w.r.t. σ by a formula $\varphi(\bar{x}, y)$ if

$$T \vdash \varphi(\underline{n}_0, \dots, \underline{n}_{k-1}, y) \leftrightarrow y = \underline{m}$$

whenever $n_0, \dots, n_{k-1}, m \in \mathbb{N}$ are such that $F(\bar{n}) = m$.

A set \mathcal{R} of prf and rp is representable in T if there exists a sequence of numerals σ such that each member of \mathcal{R} is representable in T w.r.t. σ .

In fact, [9] only consider representation of total recursive functions (trf), but it can be obviously generalized to partial functions in the indicated fashion. Likewise, we can generalize representation of rp to representation of *disjoint pairs of r.e. predicates* (dprp): such a disjoint pair $\langle P^+, P^- \rangle$, where $P^+, P^- \subseteq \mathbb{N}^k$, is represented by a formula $\varphi(\bar{x})$ if

$$\begin{aligned} \langle n_0, \dots, n_{k-1} \rangle \in P^+ &\implies T \vdash \varphi(\underline{n}_0, \dots, \underline{n}_{k-1}), \\ \langle n_0, \dots, n_{k-1} \rangle \in P^- &\implies T \vdash \neg\varphi(\underline{n}_0, \dots, \underline{n}_{k-1}) \end{aligned}$$

for all $n_0, \dots, n_{k-1} \in \mathbb{N}$. We identify any relation $P \subseteq \mathbb{N}^k$ with the disjoint pair $\langle P, \mathbb{N}^k \setminus P \rangle$.

Notice that a representation of a rp $P \subseteq \mathbb{N}^k$ is essentially the same as a representation of its characteristic function $\chi_P: \mathbb{N}^k \rightarrow \mathbb{N}$; likewise for disjoint pairs (their characteristic functions are partial). Consequently, representability of all prf in T implies representability of all trf and representability of all dprp; in turn, either of the latter two properties implies representability of all rp.

The definition of representation of functions does not demand anything from $\varphi(\bar{x}, y)$ when \bar{x} is not one of the tuples \bar{n} in the domain of the original function. However, if $\varphi(\bar{x}, y)$ represents a partial function $F: \mathbb{N}^k \rightarrow \mathbb{N}$ in T , we may define

$$\begin{aligned} \varphi'(\bar{x}, y) &\simeq \forall z (\varphi(\bar{x}, z) \leftrightarrow z = y), \\ \varphi''(\bar{x}, y) &\simeq \varphi'(\bar{x}, y) \vee (y = \underline{0} \wedge \neg\exists z \varphi'(\bar{x}, z)). \end{aligned}$$

Then φ' and φ'' also represent F in T ; moreover, φ' is T -provably a partial function, and φ'' is T -provably a total function. Thus, we could have included either condition in the definition with no ill effects.

¹In the terminology of [9], *definable*. We reserve the latter word for something else, in accordance with current standard usage.

A desirable condition that we did not include in the definition is that the sequence of numerals σ be *recursive*: that is, we can compute the term \underline{n} on input n . For most purposes, this is actually redundant if T can represent recursive functions with respect to σ : using a formula representing the (recursive) successor function $\text{succ}(n) = n + 1$, we can build a recursive sequence of formulas $\varphi_n(x)$ that define \underline{n} .

Definition 2.2 formally makes sense for representation of *arbitrary* predicates or partial functions in T . However, there is little point in that: if T is recursively axiomatizable, and the given numeral sequence is recursive (or if we can represent succ), then all predicates and total functions represented in T are actually recursive, and each partial function represented in T extends to a partial recursive function represented in T . (This is not necessarily true for non-recursive numeral sequences, see Proposition C.2.)

The primary reason for discussing representability of recursive functions in [9] is that it implies essential undecidability. We include the argument below for completeness.

Proposition 2.3 *If the set of all unary rp is representable in a theory T w.r.t. a recursive sequence of numerals, then T is essentially undecidable.*

Proof: Let $S \supseteq T$ be decidable. This makes the predicate

$$P(n) \iff n \text{ is the Gödel number of a formula } \varphi(x) \text{ s.t. } S \vdash \neg\varphi(\underline{n})$$

recursive, hence P is represented in S by a formula $\varphi(x)$. Let $n = \ulcorner \varphi \urcorner$ be its Gödel number. If $\neg P(n)$, then $S \vdash \neg\varphi(\underline{n})$ by representability, hence $P(n)$ by the definition of P , which is a contradiction. Thus, $P(n)$. Then $S \vdash \neg\varphi(\underline{n})$ by the definition of P , and $S \vdash \varphi(\underline{n})$ by representability, hence S is inconsistent. \square

Again, the assumption of recursivity of the numeral sequence in Proposition 2.3 may be replaced with representability of succ . However, it cannot be dropped entirely, as shown in the appendix (Propositions C.2 and C.3).

Likewise, it is essential in Proposition 2.3 that all unary rp are representable at once: we show in Proposition C.1 that any finite (or uniformly recursive) set of rp and trf is representable in a decidable theory. In contrast, there is one fixed unary dprp (or: prf) whose representability in a theory w.r.t. a recursive numeral sequence implies essential undecidability: in fact, any recursively inseparable pair has this property.

The reader may have realized that representation of recursive functions and predicates in T amounts to an interpretation of a particular theory in T . We now make this connection explicit.

Definition 2.4 Let \mathcal{R} be a set of prf and dprp. The language $L_{\mathcal{R}}$ consists of constants $\{\underline{n} : n \in \mathbb{N}\}$, function symbols \underline{F} of appropriate arity for every prf $F \in \mathcal{R}$, and likewise relation symbols \underline{P} for every dprp $P \in \mathcal{R}$. The theory $REP_{\mathcal{R}}$ in language $L_{\mathcal{R}}$ is axiomatized by

$$\underline{n} \neq \underline{m}$$

for $n \neq m \in \mathbb{N}$;

$$\underline{F}(\underline{n}_0, \dots, \underline{n}_{k-1}) = \underline{m}$$

for each k -ary function $F \in \mathcal{R}$, and $n_0, \dots, n_{k-1}, m \in \mathbb{N}$ such that $F(\bar{n}) = m$; and for each k -ary disjoint pair $P = \langle P^+, P^- \rangle \in \mathcal{R}$, the axioms

$$\underline{P}(\underline{n}_0, \dots, \underline{n}_{k-1})$$

for $\langle n_0, \dots, n_{k-1} \rangle \in P^+$, and

$$\neg \underline{P}(\underline{n}_0, \dots, \underline{n}_{k-1})$$

for $\langle n_0, \dots, n_{k-1} \rangle \in P^-$. This definition also applies to $\text{rp } P \subseteq \mathbb{N}^k$ using their identification with $\text{dprp } \langle P, \mathbb{N}^k \setminus P \rangle$.

Note that that the theory $REP_{\mathcal{R}}$ is axiomatized by open (= quantifier-free) sentences.

Let PRF, TRF, DPRP, and RP denote the sets of all prf, trf, dprp, and rp, respectively (where we consider $\text{TRF} \subseteq \text{PRF}$ and $\text{RP} \subseteq \text{DPRP}$). Since REP_{DPRP} is included in an extension of REP_{PRF} by quantifier-free definitions, we will use REP_{PRF} as a proxy for $REP_{\text{PRF} \cup \text{DPRP}}$.

For convenience, we also consider a finite-language formulation of REP_{PRF} . Let $U(x, y)$ be the prf defined by

$$\begin{aligned} U(0, m) &= m + 1, \\ U(n + 1, m + 1) &= [n, m], \\ U([n, m] + 1, 0) &\simeq \phi_n(m), \end{aligned}$$

where $[n, m]$ denotes a recursive bijective pairing function $\mathbb{N}^2 \rightarrow \mathbb{N}$ (e.g., the Cantor pairing function $(n + m)(n + m + 1)/2 + n$), and $\phi_n(m)$ an efficient numeration of unary prf. Let REP_U be the fragment of REP_{PRF} in the language $\langle \underline{0}, \underline{U} \rangle$; it can be axiomatized by

$$\begin{aligned} \underline{0} &\neq S(\underline{0}), \\ \underline{U}(S^n(\underline{0}), S^m(\underline{0})) &= S^k(\underline{0}) \end{aligned}$$

for all $n, m, k \in \mathbb{N}$ such that $U(n, m) = k$, where $S(x)$ denotes $\underline{U}(\underline{0}, x)$.

Lemma 2.5 *REP_{PRF} is included in an extension of REP_U by definitions of function symbols by terms, thus a theory interprets REP_{PRF} iff it interprets REP_U .*

Proof: We can read the definition of U backwards to obtain definitions of S , $[x, y]$, and $\phi_n(x)$ in terms of U and $\underline{0}$: $S(x) = U(0, x)$, $[x, y] = U(S(x), S(y))$, $\phi_n(x) = U(S([S^n(\underline{0}), x]), \underline{0})$. Then any prf $F: \mathbb{N}^k \rightarrow \mathbb{N}$ can be written in the form $F(x_0, \dots, x_{k-1}) = \phi_n([x_0, [x_1, \dots [x_{k-2}, x_{k-1}] \dots]])$ for a suitable n . \square

Using the above-mentioned fact that representations of (partial) functions may be assumed to be actual definable functions, we see:

Observation 2.6 *A set \mathcal{R} of prf and dprp is representable in a theory T according to Definition 2.2 iff $REP_{\mathcal{R}}$ is interpretable in T by a one-piece one-dimensional parameter-free interpretation I with absolute equality such that each \underline{n}^I is definable in T by a closed term.* \square

Now, the restrictions on the interpretation in Observation 2.6 are mostly irrelevant and arbitrary; as we are looking at the concept of representations from the viewpoint of interpretability, it seems we obtain a cleaner concept if we just drop them:

Definition 2.7 *A loose representation of $\mathcal{R} \subseteq \text{PRF} \cup \text{DPRP}$ in a theory T is an interpretation of $REP_{\mathcal{R}}$ in T .*

In particular, a theory T loosely represents all prf iff it interprets the theory REP_U .

2.3 The theory R

Robinson's theory R was originally defined in [9]. Some inessential variants (mutually interpretable) of the theory appear in the literature; we prefer the following form in this paper.

Definition 2.8 Let R denote the theory in the language $L_R = \langle 0, \text{succ}, +, \cdot, < \rangle$ axiomatized by

$$\underline{n} + \underline{m} = \underline{n + m}, \quad (2)$$

$$\underline{n} \cdot \underline{m} = \underline{nm}, \quad (3)$$

$$x < \underline{n} \leftrightarrow x = \underline{0} \vee \dots \vee x = \underline{n-1} \quad (4)$$

for all $n, m \in \mathbb{N}$, where $\underline{n} \simeq \text{succ}^n(0)$.

(In particular, note that axiom (4) for $n = 0$ states $\neg(x < 0)$.) It is easy to show that R implies $\underline{n} \neq \underline{m}$ for distinct $n, m \in \mathbb{N}$.

Observe that an L_R -structure is a model of R iff it contains the standard model \mathbb{N} as an initial (i.e., closed downward under $<$) substructure.

As usual, *bounded quantifiers* are introduced in L_R as the short-hands

$$\begin{aligned} \exists y < t(\bar{x}) \varphi(\bar{x}, y) &\simeq \exists y (y < t(\bar{x}) \wedge \varphi(\bar{x}, y)), \\ \forall y < t(\bar{x}) \varphi(\bar{x}, y) &\simeq \forall y (y < t(\bar{x}) \rightarrow \varphi(\bar{x}, y)), \end{aligned}$$

where t is a term not containing the variable y . An L_R -formula $\varphi(\bar{x})$ is Δ_0 (or *bounded*) if all quantifiers in φ are bounded. A formula is Σ_1 if it consists of a block of existential quantifiers followed by a Δ_0 formula.

Proposition 2.9 R proves all Σ_1 sentences true in the standard model \mathbb{N} . Conversely, it can be axiomatized by a set of true (universal) Δ_0 sentences. \square

As already proved in [9] (for the original, slightly stronger definition of the theory), R can represent recursive functions. We briefly sketch the argument below for completeness.

Proposition 2.10 Every prf $F: \mathbb{N}^k \rightarrow \mathbb{N}$ is representable in R by a Σ_1 formula w.r.t. the usual sequence of numerals as in Definition 2.8.

Proof: The graph $\{\langle \bar{x}, y \rangle : F(\bar{x}) = y\}$ is definable in \mathbb{N} by a Σ_1 formula of the form $\exists z \vartheta(\bar{x}, y, z)$, where $\vartheta \in \Delta_0$. Put

$$\begin{aligned} \alpha(w) &\simeq 0 < w \wedge \forall z < w (\text{succ}(z) = w \vee \text{succ}(z) < w), \\ \varphi(\bar{x}, y) &\simeq \exists w, z (\alpha(w) \wedge y < w \wedge z < w \wedge \vartheta(\bar{x}, y, z) \wedge \forall y', z' < w (\vartheta(\bar{x}, y', z') \rightarrow y' = y)). \end{aligned}$$

One can check

$$R \vdash \alpha(w) \rightarrow w = \underline{1} \vee \dots \vee w = \underline{r} \vee \underline{r} < w \quad (5)$$

for any $r \in \mathbb{N}$.

We claim that φ represents F in R . Assume $F(\underline{\bar{n}}) = \underline{m}$. On the one hand, $\varphi(\underline{\bar{n}}, \underline{m})$ is a true Σ_1 sentence, and as such it is provable in R . On the other hand, fix $r \in \mathbb{N}$ that witnesses the z quantifier in $\mathbb{N} \models \varphi(\underline{\bar{n}}, \underline{m})$. Working in R , assume $\varphi(\underline{\bar{n}}, y')$, we need to show $y' = \underline{m}$. Let w', z' witness the existential quantifiers in $\varphi(\underline{\bar{n}}, y')$. Using (5), either w' equals a standard numeral, or $w' > \underline{m}, \underline{r}$. In the latter case, $\theta(\underline{\bar{n}}, \underline{m}, \underline{r})$ implies $y' = \underline{m}$ as needed. In the former case, $y', z' < w'$ are also standard. It again follows that $y' = \underline{m}$, as otherwise $\neg\theta(\underline{\bar{n}}, y', z')$ would be a true Δ_0 sentence, thus provable in R . \square

Consequently, R is essentially undecidable.

It is easy to see that R (therefore any theory interpretable in R) is *locally finitely satisfiable*, i.e., every finite subset has a finite model: indeed, if we identify all elements of \mathbb{N} above $b + 1$, we obtain a model satisfying (2), (3), and (4) for $n \leq b$. Visser [12] proved a striking converse to this observation:

Theorem 2.11 *Every locally finitely satisfiable, recursively axiomatizable theory in a finite language is interpretable in R , using a one-piece one-dimensional parameter-free interpretation.* \square

Since relational $\exists\forall$ sentences have the finite model property, this in particular implies that R interprets any consistent theory axiomatized by a recursive set of $\exists\forall$ sentences in a finite relational language.

2.4 Model theory

Since this paper is intended to be accessible to a non-model-theoretic audience (and the author is not a model theorist either), it will only assume modest prerequisites in model theory—mostly common knowledge among logicians. The material needed should be covered by a textbook such as [3], except that we will also need a few concepts from classification theory. We will review a few selected topics in more detail below.

First, let us start with a few basic conventions. Recall that we allow models to be empty, and that we denote finite tuples as \bar{x} . For any structure M , we denote by $\text{Diag}(M)$ its diagram: the set of quantifier-free sentences true in M in the language of M augmented with constants for each element of M . By a slight abuse of language, we will also use this notation to denote the set of quantifier-free sentences true in M in its *original* language, if every element of M is the value of a closed term (i.e., if M is 0-generated).

Even though we normally work with one-sorted logic, the following construction is best thought of as yielding a multi-sorted structure. For any structure M , let M^{eq} be the structure that has M itself as one of its sorts, and for each equivalence relation $E(\bar{x}, \bar{y})$ on M^n definable without parameters in M , it has a sort whose elements are the equivalence classes of E ; the structure includes the projection function to this sort from M^n . It is easy to see that each such equivalence relation is definable in M by a formula that *provably* defines an equivalence relation in predicate logic; thus, the following makes sense: for any theory T , let T^{eq} be the multi-sorted theory whose models are exactly the structures M^{eq} for $M \models T$. (Officially, M^{eq} and T^{eq} can be coded in a suitable one-sorted language.) Note that T^{eq} is interpretable in T , and any interpretation of another theory S in T can be made into an interpretation with absolute equality of S in T^{eq} .

Since we will work a lot with model completions, let us recall the related background. Let \mathcal{K} be a class of structures in the same language. A model $M \in \mathcal{K}$ is *existentially closed (e.c.)* in \mathcal{K} if for every model $N \supseteq M$ such that $N \in \mathcal{K}$, we have $M \preceq_1 N$: i.e., every existential formula with parameters from M which is satisfied in N is already satisfied in M . We will often speak of (absolutely) e.c. models without reference to \mathcal{K} , in which case it is understood that \mathcal{K} is the class of all models in the given language. An e.c. model of a theory T is an e.c. structure in the class of models of T . If T is a $\forall\exists$ -axiomatized theory, then every model $M \models T$ embeds in an e.c. model of T . (More generally, this holds for any class \mathcal{K} closed under limits of chains.)

A theory T is *model-complete* if all models $M \models T$ are e.c. models of T ; this implies the stronger condition that for all $M, N \models T$, $M \subseteq N$ implies $M \preceq N$. Equivalently, T is model-complete iff every formula φ is in T equivalent to an existential formula; it is enough to test this for universal formulas φ . A stronger condition is that T has *quantifier elimination*, meaning that every formula φ is in T equivalent to a quantifier-free formula; it is enough to test this for existential formulas φ with only one quantifier. Any model-complete theory T is axiomatizable by $\forall\exists$ sentences.

Theories T and S in the same language are *companions* if every model of T embeds in a model of S , and vice versa; equivalently, $T_{\forall} = S_{\forall}$, where T_{\forall} denotes the universal fragment of T . A *model companion* of a theory T is a model-complete theory T^* that is a companion of T . There are theories with no model companion (e.g., the theory of groups), but if a theory T has a model

companion T^* , it is unique: the models of T^* are exactly the e.c. models of T_\forall . A theory has a model companion iff the class of e.c. models of T_\forall is elementary. Notice that a model companion of T is the same thing as a model companion of T_\forall , hence we can as well restrict attention to universal theories T .

If T is a theory, and $\varphi(\bar{x})$ an existential formula, the *resultant* Res_φ^T is the set of all universal formulas $\psi(\bar{x})$ such that $T \vdash \varphi(\bar{x}) \rightarrow \psi(\bar{x})$. For any structure M and $\bar{a} \in M$, we then have: $M \models \text{Res}_\varphi^T(\bar{a})$ iff M embeds in a model of T satisfying $\varphi(\bar{a})$. It follows that the class of e.c. models of T is axiomatized by the following set of *infinitary* formulas:

$$T + \left\{ \forall \bar{x} \left(\bigwedge \text{Res}_\varphi^T(\bar{x}) \rightarrow \varphi(\bar{x}) \right) : \varphi \in \exists_1 \right\}. \quad (6)$$

A universal theory T has a model companion T^* iff the resultants Res_φ^T are finitely axiomatizable (over T) for all $\varphi \in \exists_1$, in which case (6) provides an explicit axiomatization of T^* .

A *model completion* of a theory T is a model companion T^* of T such that for every $M \models T$, the theory $T^* + \text{Diag}(M)$ is complete. Equivalently, a model companion T^* of T is a model completion of T iff T has the amalgamation property (cf. Definition B.4). If T is a universal theory (which is the case we are primarily interested in), a companion T^* of T is a model completion of T iff T^* has quantifier elimination. A universal theory T has a model completion iff the resultant Res_φ^T is equivalent to a quantifier-free formula over T for every $\varphi \in \exists_1$.

For completeness, we also mention that every theory T has a unique *Kaiser hull* T^{KH} , which is a largest $\forall\exists$ -axiomatized companion of T . Any e.c. model of T_\forall is also a model of T^{KH} . If T has a model companion T^* , then $T^* = T^{\text{KH}}$.

A convenient trick when studying models of a complete theory T is to use *monster models*. A monster model of T is a model $\mathbb{M} \models T$ sufficiently rich so that all models we need to discuss can be assumed to be submodels of \mathbb{M} ; in order for this to work, we make \mathbb{M} highly saturated: to be specific, let us posit that \mathbb{M} is κ -saturated (i.e., every type over $< \kappa$ parameters from \mathbb{M} is realized in \mathbb{M}) and strongly κ -homogeneous (i.e., every partial elementary self-map of \mathbb{M} of size $< \kappa$ extends to an automorphism of \mathbb{M}), where κ is a “large” cardinal number (in particular, larger than the size of the language, as well as any models of T that we are going to encounter during the argument). This also implies that \mathbb{M} is κ^+ -universal (every model of T of size $\leq \kappa$ elementarily embeds in \mathbb{M}). (If it were not for foundational issues that we prefer not to be dragged into, we could even take \mathbb{M} as an “Ord-saturated” model: a proper class model of T saturated w.r.t. types over any *set* of parameters.) Having fixed the monster model \mathbb{M} , a *small set* is a subset of \mathbb{M} of size $< \kappa$ (likewise for sequences and other similar objects); a *small model* is an elementary submodel of \mathbb{M} of size $< \kappa$.

Finally, let us introduce a few notions from *classification theory*. Classification theory was developed by Shelah [8] (and subsequently many others) as an elaboration of Morley’s theory of stability; one of its main themes is identifying useful “dividing lines” between tame and wild theories. The dividing lines we are going to mention here are essentially of two kinds: first, variants of stability based on counting of types, and second, combinatorial properties based on the appearance of certain arrangements of points and definable sets in models.

While model theorists prefer to work with complete theories, the properties below are all stated in such a way that a theory T has a “tameness” property P iff every completion of T has property P . Also, it will be generally the case that T has a (tameness) property P iff every countable-language fragment of T has property P .

For an overview of inclusions among the properties below, see Figure 1.

Theory T is κ -*stable* if for every $M \models T$ and $A \subseteq M$ of size $|A| \leq \kappa$, there are at most κ complete types over A . We say that T is *stable* if it is κ -stable for some infinite cardinal κ , and *superstable* if it is κ -stable for all sufficiently large cardinals κ . An even stronger condition




Figure 1: Main dividing lines

is that T be *totally transcendental*; officially, this means that every formula φ has Morley rank $\text{MR}(\varphi) < \infty$, but as we do not want to get into a definition of the rank, we can use the following characterization: T is totally transcendental iff all countable-language fragments of T are ω -stable. For countable theories, ω -stability is equivalent to total transcendence, and implies κ -stability for all infinite cardinals κ ; the class of ω -stable theories includes uncountably categorical theories, which in turn include *strongly minimal* theories (meaning that for all $M \models T$, the only subsets of M definable with parameters are finite or cofinite).

Concerning the second kind of tameness properties, we will start with an alternative definition of stability as the first example. A theory T has the *order property* (OP) if there exists a formula $\varphi(\bar{x}, \bar{y})$ (where \bar{x} and \bar{y} are tuples of the same length), a model $M \models T$, and a sequence of tuples $\{\bar{a}_i : i \in \omega\}$ in M such that

$$M \models \varphi(\bar{a}_i, \bar{a}_j) \iff i < j$$

for all $i, j \in \omega$; otherwise, T has the *no-order property* (NOP). It turns out that T has NOP if and only if it is stable.

Beware of the terminological peculiarity that the base form of this condition on theories (OP) is “negative” (wild), whereas the corresponding “positive” (tame) condition is denoted as its negation (NOP). All properties below follow the same naming pattern.

A theory T has the *independence property* (IP) if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_i : i \in \omega\}$ and $\{\bar{b}_I : I \subseteq \omega\}$ in M such that

$$M \models \varphi(\bar{a}_i, \bar{b}_I) \iff i \in I$$

for all $i \in \omega$ and $I \subseteq \omega$. Otherwise, T is NIP (also called *dependent*).

A theory T has the *strict order property* (SOP) if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_i : i \in \omega\}$ in M such that

$$M \models \exists \bar{x} (\varphi(\bar{x}, \bar{a}_i) \wedge \neg \varphi(\bar{x}, \bar{a}_j)) \iff i < j$$

for all $i, j \in \omega$; equivalently, T is SOP iff there is a formula $\varphi(\bar{x}, \bar{y})$ that T -provably defines a strict partial order, and there is a model $M \models T$ in which the partial order defined by φ has an infinite chain. Otherwise, T is NSOP.

A theory is stable (NOP) if and only if it is both NIP and NSOP. The class of theories that are NIP but SOP includes *o-minimal* theories (i.e., theories featuring a linear order $<$ such that every definable subset of the universe is a union of finitely many intervals, possibly degenerate).

Recall that $X^{<\omega}$ denotes the set of finite sequences with entries from X , ordered by the initial subsequence relation (which we write as $t \subseteq s$) to form an X -branching tree; X^ω is the corresponding set of infinite sequences (which are branches of the tree). If $s \in X^{<\omega} \cup X^\omega$, and $n < \text{lh}(s)$, then $s \upharpoonright n$ is the initial subsequence of s of length n ; if $s \in X^{<\omega}$ and $x \in X$, then $s \hat{\ } x$ is s extended with a new entry x at the end. For clarity, we will write von Neumann numerals as $\mathbf{n} = \{0, \dots, n-1\}$.

A set of formulas is *k-inconsistent* if each k -element subset is inconsistent.

A theory T has the *tree property* (TP) if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, tuples $\{\bar{a}_s : s \in \omega^{<\omega}\}$ in M , and $k \geq 2$ such that

- for each $\sigma \in \omega^\omega$, the type $\{\varphi(\bar{x}, \bar{a}_{\sigma \upharpoonright n}) : n < \omega\}$ is consistent, and
- for each $s \in \omega^{<\omega}$, $\{\varphi(\bar{x}, \bar{a}_{s \hat{\ } i}) : i < \omega\}$ is k -inconsistent.

Otherwise, T is called NTP or *simple*. Simplicity is officially defined in terms of properties of forking, but it is equivalent to NTP; there is a related stronger condition called *supersimplicity*, see e.g. [13]. Stable theories are simple, and simple theories are NSOP. Superstable theories are exactly the stable supersimple theories.

The tree property has two important variants. A theory T has the *tree property* TP_1 if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_s : s \in \omega^{<\omega}\}$ in M such that

- for each $\sigma \in \omega^\omega$, $\{\varphi(\bar{x}, \bar{a}_{\sigma \upharpoonright n}) : n < \omega\}$ is consistent, and
- for each incomparable $s, t \in \omega^{<\omega}$, $\{\varphi(\bar{x}, \bar{a}_s), \varphi(\bar{x}, \bar{a}_t)\}$ is inconsistent.

T has the *tree property* TP_2 if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_{n,i} : n, i \in \omega\}$ in M such that

- for each $\sigma \in \omega^\omega$, $\{\varphi(\bar{x}, \bar{a}_{n, \sigma(n)}) : n < \omega\}$ is consistent, and
- for each $n, i, j \in \omega$ such that $i < j$, $\{\varphi(\bar{x}, \bar{a}_{n,i}), \varphi(\bar{x}, \bar{a}_{n,j})\}$ is inconsistent.

As usual, if T is not TP_i , it is NTP_i . A theory is NTP if and only if it is both NTP_1 and NTP_2 . All NIP theories are NTP_2 , and all NTP_1 theories are NSOP.

The region between simple and NSOP theories is further stratified by levels of the *strong order property*. For $k \geq 3$, a theory T has the *strong order property* SOP_k if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_i : i < \omega\}$ in M such that $M \models \varphi(\bar{a}_i, \bar{a}_j)$ for all $i < j < \omega$, but

$$\{\varphi(\bar{x}_0, \bar{x}_1), \varphi(\bar{x}_1, \bar{x}_2), \dots, \varphi(\bar{x}_{k-2}, \bar{x}_{k-1}), \varphi(\bar{x}_{k-1}, \bar{x}_0)\} \quad (7)$$

is inconsistent; otherwise, T has NSOP_k . A theory T has the *strong order property* SOP_∞ if there are data as above such that (7) is inconsistent for all $k \geq 3$; otherwise, T has NSOP_∞ . For any theory T , we have

$$\text{NTP}_1 \implies \text{NSOP}_3 \implies \text{NSOP}_4 \implies \dots \implies \text{NSOP}_\infty \implies \text{NSOP}.$$

We warn the reader that usage of the abbreviations (N)SOP $_\infty$ and (N)SOP varies in the literature.

Notice that the definition of NSOP_k above is only interesting for $k \geq 3$, as stated: taking it blindly for $k = 2$ would give a condition equivalent to stability, and for $k = 1$ a condition false for any theory with infinite models. Instead, the names NSOP_1 and NSOP_2 were given ad hoc to variants of the tree property that fit nicely in the picture. Since NSOP_2 is equivalent to NTP_1 , we will not bother to define it separately. A theory T has property SOP_1 if there is a formula $\varphi(\bar{x}, \bar{y})$, a model $M \models T$, and tuples $\{\bar{a}_s : s \in \mathbf{2}^{<\omega}\}$ in M such that

- for each $\sigma \in \mathbf{2}^\omega$, $\{\varphi(\bar{x}, \bar{a}_{\sigma \upharpoonright n}) : n < \omega\}$ is consistent, and

- for each $s, t \in \mathbf{2}^{<\omega}$, if $s \frown 0 \subseteq t$, then $\{\varphi(\bar{x}, \bar{a}_{s \frown 1}), \varphi(\bar{x}, \bar{a}_t)\}$ is inconsistent;

otherwise, T is NSOP₁. We have

$$\text{NTP} \implies \text{NSOP}_1 \implies \text{NSOP}_2 \iff \text{NTP}_1 \implies \text{NSOP}_3 \implies \dots$$

for any theory T .

We observe that each of the combinatorial properties above (NOP, NSOP, NIP, NTP_{1,2}, NSOP_k, NSOP_∞) can be expressed as weak non-interpretability of a particular recursively $\exists\forall$ -axiomatized theory in a finite relational language. (Incidentally, notice that any such theory is interpretable in R by Theorem 2.11, hence R is “wild” according to all of these dividing lines.)

For example, by compactness, a theory T has SOP iff it has a model with a definable strict order (on k -tuples, for some k) with arbitrarily long finite chains. It makes no difference if the defining formula is allowed extra parameters, or if we allow to relativize the domain of the order. Thus, T has SOP iff it has a completion that interprets the theory T_{SOP} of strict orders with arbitrarily long chains, axiomatized by

$$\begin{aligned} \forall x, y, z (x < y \wedge y < z \rightarrow x < z), \\ \forall x \neg(x < x), \\ \exists x_0, \dots, x_n \bigwedge_{i < n} x_i < x_{i+1} \end{aligned}$$

for $n \in \omega$.

For a more complicated example, T has TP₁ iff it weakly interprets the theory in a language with a single binary relation $R(x, y)$, and axioms

$$\exists\{y_s : s \in \mathbf{n}^{\leq n}\} \left(\bigwedge_{s \in \mathbf{n}^n} \exists x \bigwedge_{i \leq n} R(x, y_{s \upharpoonright i}) \wedge \bigwedge_{\substack{s, t \in \mathbf{n}^{\leq n} \\ s \not\subseteq t \not\subseteq s}} \forall x \neg(R(x, y_s) \wedge R(x, y_t)) \right)$$

for $n \in \omega$.

3 Model completion of the empty theory

Recall that our original motivation was to find a theory T that represents prf, but does not interpret R . Now, the weaker T is, the lower its chances of interpreting R , so the obvious choice is to take $T = \text{REP}_{\text{PRF}}$. This theory, axiomatized by quantifier-free sentences, essentially just states that the universe includes a copy of a certain model based on the integers; it does not take a big leap of faith to surmise it is too weak to interpret much of anything. It is, however, another matter to actually prove this. A possible strategy is to consider an arbitrary translation I of the language of R into REP_{PRF} , and try to argue that in some models of REP_{PRF} , R^I is not valid. But here the weakness of REP_{PRF} that we were hoping to exploit becomes our worst enemy: I may involve formulas of arbitrary high quantifier complexity that may potentially denote very complicated combinatorial properties, and we just have no handle how to understand them. What we need is that definable sets have manageable structure.

Ideally, we would like to extend REP_{PRF} to a (consistent) theory T with full quantifier elimination. Now, a moment of reflection tells us that any possible configuration of finitely many functions on a finite set may be realized by suitable recursive functions, and as such should embed into a model of T . By compactness, *any* L_{PRF} -structure should embed in a model of T , thus if such a T exists, it is unique: T must be the model completion of the empty theory in L_{PRF} . (By the empty theory, we mean the theory with no extra-logical axioms.)

The model completion of the empty L -theory is well known and well understood for finite relational languages L . The theory can be axiomatized by a transparent set of “extension axioms”, and it coincides with the set of all formulas that hold in random finite L -structures with asymptotic probability 1. The theory is ω -categorical, and its unique countable model is the countable random L -structure generalizing the Erdős–Rényi–Rado random graph; alternatively, it can be described as the Fraïssé limit of the class of all finite L -structures.

It is much less known that the model completion of the empty L -theory does, indeed, exist for arbitrary languages L , as we need here. This was proved by Winkler [14] as a corollary of more general results on model companions of Skolem expansions of model-complete theories. Note that for languages with functions, most of the above-mentioned properties of the theory of the random relational structure break down: first-order logic with functions has no 0–1 law (or even limit law) on finite structures, and there does not seem to be a sensible way of defining a probability distribution on functions on infinite sets; we will see that the model completion is not ω -categorical, and not locally finitely satisfiable.

We will now give a self-contained argument that the model completion exists, including an explicit axiomatization by extension formulas; this will also help us later to determine (syntactically) what open formulas are consistent with the theory. We will denote the model completion as EC_L , as its models are exactly the (absolutely) existentially closed L -structures.

Definition 3.1 Let L be a finite language, and Θ be a finite set of L -terms closed under subterms such that the variables in Θ are among $x_0, \dots, x_{n-1}, y_0, \dots, y_{m-1}$. Let $\varepsilon_t^R \in \{0, 1\}$ for every k -ary relation $R \in L$, and every $t_0, \dots, t_{k-1} \in \Theta$. Let \sim be an equivalence relation on Θ such that:

- (i) If $R \in L$ is k -ary, and $t_i \sim s_i$ for each $i < k$, then $\varepsilon_t^R = \varepsilon_s^R$.
- (ii) If $F \in L$ is k -ary, and $t \simeq F(\bar{t}) \in \Theta$ and $s \simeq F(\bar{s}) \in \Theta$ satisfy $t_i \sim s_i$ for each $i < k$, then $t \sim s$.

Then the elementary existential formula $\exists \bar{y} \theta_{\Theta, \sim, \varepsilon}(\bar{x}, \bar{y})$ is defined by

$$\theta_{\Theta, \sim, \varepsilon}(\bar{x}, \bar{y}) \simeq \bigwedge_{\substack{t, s \in \Theta \\ t \sim s}} t = s \wedge \bigwedge_{\substack{t, s \in \Theta \\ t \not\sim s}} t \neq s \wedge \bigwedge_{\substack{R \in L \\ \bar{t} \in \Theta}} R^{\varepsilon_{\bar{t}}^R}(\bar{t}), \quad (8)$$

where $\psi^1 \simeq \psi$, $\psi^0 \simeq \neg\psi$.

Lemma 3.2 Every existential formula $\exists \bar{y} \theta(\bar{x}, \bar{y})$ in a finite language L is equivalent to a disjunction of elementary existential formulas with the same free and bound variables.

Proof: Let Θ be the set of all subterms of $\theta(\bar{x}, \bar{y})$, and Φ the set of all (finitely many) atomic formulas using terms from Θ . We can write θ in full disjunctive normal form in atoms Φ , and switch disjunctions with existential quantifiers. Each disjunct has the form (8), except for the conditions on \sim and ε . However, it is easy to see that if \sim is not an equivalence relation, or if (i) or (ii) is violated, then $\theta_{\Theta, \sim, \varepsilon}$ is contradictory. \square

Definition 3.3 Let $L, \Theta, \bar{x}, \bar{y}, \sim, \varepsilon$ be as in Definition 3.1. We define a subset $\Xi \subseteq \Theta$, and for each $t \in \Xi$ a term $t^*(\bar{x})$, as follows:

- (i) Every variable x_i is in Ξ , and $x_i^* \simeq x_i$.
- (ii) If $t \sim s \in \Xi$, then $t \in \Xi$, and $t^* \simeq s^*$.
- (iii) If $t \simeq F(t_0, \dots, t_{k-1}) \in \Theta$, and $t_0, \dots, t_{k-1} \in \Xi$, then $t \in \Xi$, and $t^* \simeq F(t_0^*, \dots, t_{k-1}^*)$.

If more than one clause applies to put $t \in \Xi$, we define t^* using any of them; the choice does not matter. The only relevant property is the following observation:

$$\vdash \theta_{\Theta, \sim, \varepsilon}(\bar{x}, \bar{y}) \rightarrow t(\bar{x}, \bar{y}) = t^*(\bar{x}) \quad (9)$$

for every $t \in \Xi$. Finally, we define an open formula $\theta_{\Theta, \sim, \varepsilon}^*(\bar{x})$ as

$$\bigwedge_{\substack{t, s \in \Xi \\ t \sim s}} t^* = s^* \wedge \bigwedge_{\substack{t, s \in \Xi \\ t \not\sim s}} t^* \neq s^* \wedge \bigwedge_{\substack{R \in L \\ \bar{t} \in \Xi}} R^{\varepsilon_{\bar{t}}^R}(\bar{t}^*) \wedge \bigwedge_{\substack{t = F(\bar{t}) \in \Xi \\ \bar{t} \in \Xi}} t^* = F(\bar{t}^*).$$

Proposition 3.4 *Let $L, \Theta, \sim, \varepsilon$ be as in Definition 3.1, M an L -structure, and $\bar{u} \in M$. The following are equivalent.*

- (i) *There exists an extension $N \supseteq M$ such that $N \models \exists \bar{y} \theta_{\Theta, \sim, \varepsilon}(\bar{u}, \bar{y})$.*
- (ii) *$M \models \theta_{\Theta, \sim, \varepsilon}^*(\bar{u})$.*

Proof: (i) \rightarrow (ii) follows immediately from the definitions and property (9).

(ii) \rightarrow (i): Put $N = M \dot{\cup} (\Theta \setminus \Xi) / \sim$, and identify t / \sim with $t^*(\bar{u})$ for $t \in \Xi$. If $t_0, \dots, t_{k-1} \in \Theta$, $R \in L$, and $t = F(\bar{t}) \in \Theta$, define

$$\begin{aligned} R^N(t_0 / \sim, \dots, t_{k-1} / \sim) &\iff \varepsilon_{\bar{t}}^R = 1, \\ F^N(t_0 / \sim, \dots, t_{k-1} / \sim) &= t / \sim. \end{aligned}$$

Using $M \models \theta_{\Theta, \sim, \varepsilon}^*(\bar{u})$ and the properties of \sim and Ξ , it is easy to check that the definition is independent of the choice of representatives, and agrees with M if $t_0, \dots, t_{k-1} \in \Xi$. We can thus extend the definition with the original structure of M , and then arbitrarily to the remaining tuples from N . The definition ensures

$$N \models \theta_{\Theta, \sim, \varepsilon}(\bar{x} / \sim, \bar{y} / \sim),$$

where $x_i / \sim = u_i$. □

In other words, Proposition 3.4 shows that $\theta_{\Theta, \sim, \varepsilon}^*$ is equivalent to the resultant of the elementary existential formula $\exists \bar{y} \theta_{\Theta, \sim, \varepsilon}(\bar{x}, \bar{y})$ in the empty theory.

Lemma 3.5 *Let M be an L -structure.*

- (i) *If M is existentially closed, then so is $M \upharpoonright L'$ for every $L' \subseteq L$.*
- (ii) *If $M \upharpoonright L'$ is e.c. for every finite $L' \subseteq L$, then M is e.c.*

Proof: (i): Every extension of $M \upharpoonright L'$ can be expanded to an extension of M . (ii): Assume that $M \subseteq N \models \varphi(\bar{u})$, where $\bar{u} \in M$, and φ is an existential formula. Let $L' \subseteq L$ be a finite sublanguage containing all symbols occurring in φ . Then $\varphi(\bar{u})$ holds in $N \upharpoonright L'$, hence in $M \upharpoonright L'$ (and M) as $M \upharpoonright L'$ is e.c. □

Definition 3.6 *If L is a finite language, let EC_L denote the theory axiomatized by the formulas*

$$\theta_{\Theta, \sim, \varepsilon}^*(\bar{x}) \rightarrow \exists y \theta_{\Theta, \sim, \varepsilon}(\bar{x}, y)$$

for all $\Theta, \sim, \varepsilon$ as in Definition 3.1 with $m = 1$.

For infinite L , we put $EC_L = \bigcup \{EC_{L'} : L' \subseteq L \text{ finite}\}$.

Theorem 3.7 *For any language L , EC_L is a model completion of the empty L -theory:*

- (i) EC_L has elimination of quantifiers.
- (ii) Models of EC_L are exactly the existentially closed L -structures; in particular, every L -structure embeds in a model of EC_L .

Moreover, if L is recursively presented, then EC_L is decidable.

Proof: By Lemma 3.5, we may assume that L is finite.

(i): Proposition 3.4 implies the converse implications

$$\vdash \exists y \theta_{\Theta, \sim, \varepsilon}(\bar{x}, y) \rightarrow \theta_{\Theta, \sim, \varepsilon}^*(\bar{x})$$

for each axiom of EC_L . In view of Lemma 3.2, this shows that every formula with one existential quantifier is equivalent to an open formula over EC_L , hence the same follows for all formulas by induction on complexity.

(ii): Proposition 3.4 implies that any e.c. model validates EC_L . The converse would also follow from Proposition 3.4, were it not for the restriction to $m = 1$ in the definition of EC_L . However, if $\varphi(\bar{x})$ is an existential formula, $N \supseteq M \models EC_L$, and $N \models \varphi(\bar{u})$ for some $\bar{u} \in M$, we can assume N is e.c. by extending it further if necessary. Thus, $N \models EC_L$, and M is an elementary substructure of N by (i), hence $M \models \varphi(\bar{u})$.

As for decidability, EC_L is clearly r.e., hence quantifier elimination is effective (in fact, the argument above gives an explicit algorithm). Now, if φ is a quantifier-free (or universal) sentence, then $EC_L \vdash \varphi$ iff $\emptyset \vdash \varphi$ iff φ holds in all finite models of cardinality bounded by the number of subterms t of φ : if $M \not\models \varphi$, let M_0 be the set of realizations of all such t in M , and make it an L -structure in such a way that realizations of all symbols agree with M whenever possible. Then $M_0 \not\models \varphi$. \square

Corollary 3.8 *If M is a recursive L -structure, $EC_L + \text{Diag}(M)$ is a decidable complete theory.* \square

4 Interpretability in existential theories

As explained in the beginning of Section 3, our intention for discussing EC_L is that we want to show noninterpretability of certain theories S in REP_{PRF} (which is an existential—in fact, quantifier-free—theory) by showing their noninterpretability in completions of EC_L . Now, it is not a priori clear if this strategy is any good: why should we expect that S is, indeed, not interpretable in an extension of EC_L ? After all, EC_L is a fairly nontrivial theory, hence it may interpret S even if no existential theory can; to begin with, it interprets EC_L itself.

To allay our fears, we will prove in the present section that a large class of theories S is immune to such shenanigans: specifically, a theory axiomatized by $\exists\forall$ sentences is interpretable in a completion of some EC_L if and only if it is interpretable in a consistent quantifier-free or existential theory. We believe this characterization to be of independent interest, which is why we prove it in detail even though it is only of indirect relevance for our original goal, in that it assures us that we set off in the right direction; of course, we would eventually find that anyway when we get to the point.

We start with a few auxiliary lemmas whose basic intention is that if a theory is (weakly) interpretable in EC_L , we can make L and the interpretation “nice”.

Lemma 4.1

- (i) If $L' \supseteq L$, $EC_{L'}$ is a conservative extension of EC_L .
- (ii) If $L' = L \cup \{c_i : i \in I\}$, where $c_i \notin L$ are constants, then $EC_{L'}$ is axiomatized by EC_L . In particular, $EC_{L'} \vdash \varphi(\bar{c})$ iff $EC_L \vdash \forall \bar{x} \varphi(\bar{x})$.
- (iii) If L contains a constant c , and $F, R \notin L$ are a k -ary function and relation symbol (resp.), then

$$R^I(\bar{x}) \simeq F(\bar{x}) = c$$

provides a faithful interpretation of $EC_{L \cup \{R\}}$ in $EC_{L \cup \{F\}}$.

- (iv) If L contains an at least binary function, or at least two unary functions, then EC_L has pairing.

Proof: (i): $EC_{L'} \supseteq EC_L$ follows from Lemma 3.5. On the other hand, any model $M \models EC_L$ has an expansion to an L' -structure M' , which has an extension $N' \models EC_{L'}$. M is an elementary substructure of $N' \upharpoonright L$ by quantifier elimination for EC_L .

(ii) follows from (i) and the fact that an expansion of an e.c. model by constants is e.c.

(iii): Let $M \models EC_{L \cup \{F\}}$, and $\varphi(\bar{u})$ be an existential $(L \cup \{R\})$ -formula with $\bar{u} \in M$, satisfied in an extension $N \supseteq M^I$. We can extend F^M to N so that $F^N(\bar{a}) = c$ iff $R^N(\bar{a})$ for all $\bar{a} \in N$. Then $\langle N \upharpoonright L, F^N \rangle \models \varphi^I(\bar{u})$, where φ^I is an existential formula, hence $M \models \varphi^I(\bar{u})$ by e.c., and $M^I \models \varphi(\bar{u})$. Thus, $M^I \models EC_{L \cup \{R\}}$, which shows that $EC_{L \cup \{F\}} \vdash EC_{L \cup \{R\}}^I$. On the other hand, assume $EC_{L \cup \{F\}} \vdash \varphi^I$, and $M \models EC_{L \cup \{R\}}$. Let $F^M : M^k \rightarrow M$ be such that $F^M(\bar{a}) = c^M$ iff $R^M(\bar{a})$, and $N \supseteq \langle M \upharpoonright L, F^M \rangle$ be a model of $EC_{L \cup \{F\}}$. We have $N^I \models EC_{L \cup \{R\}} + \varphi$, and $M \preceq N^I$ by quantifier elimination, hence $M \models \varphi$.

(iv): If L contains two distinct unary function symbols $L(x), R(x)$, we claim that

$$EC_L \vdash \forall x, y \exists z (L(z) = x \wedge R(z) = y).$$

Let $M \models EC_L$, and $a, b \in M$. Put $N = M \dot{\cup} \{c\}$, where $L^N(c) = a, R^N(c) = b$, and the realization of other functions or relations on tuples involving c is arbitrary. Then $N \models \exists z (L(z) = a \wedge R(z) = b)$, hence the same holds in M by e.c.

If L contains a k -ary function F for $k \geq 2$, we may use a similar argument with e.g.

$$\begin{aligned} L(x) &= F(F(x, \dots, x), x, \dots, x), \\ R(x) &= F(x, \dots, x, F(x, \dots, x)), \end{aligned}$$

and $N = M \dot{\cup} \{c, F(c, \dots, c)\}$. □

Definition 4.2 We will call a parameter-free translation *quantifier-free* if its domain, and the translations of all predicate symbols as well as equality are given by quantifier-free formulas, and the translations of all function symbols are given piecewise by terms, where the pieces are finitely many and quantifier-free definable.

Recall Lemma 2.1.

Lemma 4.3 For any language L , there is a quantifier-free one-piece one-dimensional parameter-free unrelativized cointerpretation with absolute equality of EC_{L_2} in EC_L , where L_2 consists of a single binary function, and at most $|L|$ constants. If L is countable, one constant suffices.

Proof: Using Lemma 4.1 (iii) (which may be applied in parallel to all relations using the same argument), we may assume L contains no relations. Let L_2 be the language consisting of a binary function (x, y) , the constants of L , and new constants c_F for every nonconstant function $F \in L$. For $n \geq 1$, write

$$(x_0, \dots, x_{n-1}) := (x_0, (x_1, \dots, (x_{n-2}, x_{n-1}) \dots)).$$

Let I be the translation of L into L_2 defined by $c^I = c$ for constants $c \in L$, and

$$F^I(x_0, \dots, x_{n-1}) = ((c_F, x_0, \dots, x_{n-1}), x_0, \dots, x_{n-1}, x_0, \dots, x_{n-1})$$

for n -ary functions $F \in L$, $n > 0$. Let

$$EC_{L_2}^* = EC_{L_2} + \{c_F \neq c_G : F \neq G \in L\}.$$

We will show that I is a faithful interpretation of EC_L in $EC_{L_2}^*$, which implies it is also a cointerpretation of EC_{L_2} in EC_L .

Claim 4.3.1

- (i) If N is an L_2 -structure such that the constants c_F^N are pairwise distinct, and M is an extension of the L -structure N^I , there is an extension $K \supseteq N$ such that $K^I \supseteq M$.
- (ii) If M is an L -structure, there is an L_2 -structure N such that $N^I \supseteq M$, and the constants c_F^N are distinct.

Proof: (i): Let K be the disjoint union $M \dot{\cup} M^{<\omega}$, with constants realized as in N , and

$$\begin{aligned} (a, b)^K &= (a, b)^N & a, b \in N, \\ (a_0, \langle a_1, \dots, a_n \rangle)^K &= \langle a_0, \dots, a_n \rangle & a_0 \in M, \\ (\langle c_F, u_0, \dots, u_{i-1} \rangle, \langle a_0, \dots, a_{n-1}, v_0, \dots, v_{j-1} \rangle)^K &= F^M(a_0, \dots, a_{n-1}) & F \text{ } n\text{-ary, } i, j \geq 0, \\ (a, b)^K &= \langle a \rangle & a, b \in M, \{a, b\} \not\subseteq N, \\ (a, b)^K &= \langle \rangle & \text{all other cases.} \end{aligned}$$

We need to check that if $F \in L$ is n -ary with $n > 0$, and $a_0, \dots, a_{n-1} \in M$, then

$$F^M(\bar{a}) = (F^I(\bar{a}))^K.$$

We may assume $\bar{a} \notin N^n$, as otherwise the statement follows from $N^I \subseteq M$. Let $i < n$ be maximal such that $a_i \notin N$, and $i' = \min\{i, n-2\}$. It follows from the definition that

$$\begin{aligned} (a_0, \dots, a_{n-1}, a_0, \dots, a_{n-1})^K &= \langle a_0, \dots, a_{n-1}, a_0, \dots, a_{i'} \rangle, \\ (c_F, a_0, \dots, a_{n-1})^K &= \langle c_F, a_0, \dots, a_{i'} \rangle, \end{aligned}$$

hence

$$((c_F, a_0, \dots, a_{n-1}), a_0, \dots, a_{n-1}, a_0, \dots, a_{n-1})^K = F^M(a_0, \dots, a_{n-1})$$

as required.

(ii): By extending M if necessary, we may assume $|M| \geq |L|$, hence we can fix pairwise distinct elements $c_F^N \in M$. Put $N = M \dot{\cup} M^{<\omega}$, and define

$$\begin{aligned} (a, b)^N &= \langle a, b \rangle & a, b \in M, \\ (a_0, \langle a_1, \dots, a_n \rangle)^N &= \langle a_0, \dots, a_n \rangle & a_0 \in M, \\ (\langle c_F, u_0, \dots, u_{i-1} \rangle, \langle a_0, \dots, a_{n-1}, v_0, \dots, v_{j-1} \rangle)^N &= F^M(a_0, \dots, a_{n-1}) & F \text{ } n\text{-ary, } i, j \geq 0, \\ (a, b)^N &= \langle \rangle & \text{otherwise.} \end{aligned}$$

We have $M \subseteq N^I$ by a similar (but easier) argument as in (i).

□ (Claim 4.3.1)

In order that I interprets EC_L in $EC_{L_2}^*$, it suffices to show that if $N \models EC_{L_2}^*$, then N^I is e.c. Now, if an existential L -formula φ with parameters from N is satisfiable in $M \supseteq N^I$, then $M \subseteq K^I$ for some $K \supseteq N$ by the claim, which thus satisfies the existential formula φ^I . It follows that $N \models \varphi^I$ as N is e.c., i.e., $N^I \models \varphi$.

To show that I is faithful, let $EC_{L_2}^* \vdash \varphi^I$, and $M \models EC_L$. By the claim, there is N with the elements c_F^N pairwise distinct such that $N^I \supseteq M$. By extending it if necessary, we may assume $N \models EC_{L_2}^*$, hence $N^I \models \varphi$. Also, $N^I \models EC_L$, hence $M \preceq N^I$ by quantifier elimination, which gives $M \models \varphi$.

Finally, let L be countable, and enumerate it as $\{F_k : k \in \omega\}$. Let L_2 be the language consisting of (x, y) and a single constant c . We modify the construction above as follows: we employ the closed terms

$$c_k = (\dots \underbrace{((c, c), c), \dots, c}_{n+2 \text{ times}}).$$

in place of c_{F_k} , and if F_k is a constant, we put $F_k^I = (c_k, c_k)$. (In particular, we redefine $EC_{L_2}^*$ to state that all the c_k are pairwise distinct.) Then it is easy to check that the argument still goes through: the only place where the exact composition of c_F matters is in the proof of part (ii) of the claim, and we can fix it e.g. by making N the set of all finite binary trees with leaves labelled by $M \cup \{c\}$, where $a \in M$ is identified with a one-node tree, and $(x, y)^N$ is the tree whose root has children x, y , except for

$$\begin{aligned} ((c_k, a_0, \dots, a_{n-1})^N, (a_0, \dots, a_{n-1}, a_0, \dots, a_{n-1})^N)^N &= F_k^M(a_0, \dots, a_{n-1}) && F_k \text{ } n\text{-ary, } \bar{a} \in M, \\ (c_k^N, c_k^N)^N &= F_k^M && F_k \text{ constant,} \end{aligned}$$

as needed to make the interpretation work. \square

Corollary 4.4 *If a theory T is weakly interpretable in EC_L for some L , it has a one-piece one-dimensional parameter-free unrelativized interpretation in a consistent extension of some EC_{L_2} , where L_2 consists of a binary function, and at most $|L|$ constants. If L is countable, one constant suffices.*

Proof: We can make the interpretation one-piece as EC_L proves there are at least two elements. We can assume L contains a constant by Lemma 4.1 (i), and that it is purely functional by (iii). Then we can make the interpretation one-dimensional by (iv), and parameter-free by expanding L with constants for the parameters, using (ii). We can also assume to have a constant c denoting an element in the domain of the interpretation, and then it is easy to make the interpretation unrelativized by equating (i.e., extending the interpreted equality) elements outside the domain with c . Finally, we can compose the interpretation with the one from Lemma 4.3 to make the language as needed. \square

Note that the argument in Corollary 4.4 does not guarantee that the interpretation is quantifier-free: while the domain and the translations of all symbols can be made quantifier-free formulas just by quantifier elimination, this does not ensure function symbols are given piecewise by terms. This will in fact pose a serious challenge in the proof of the characterization below, and we will need results on elimination of imaginaries from Appendix A to deal with it.

Theorem 4.5 *Let T be an $\exists\forall$ -axiomatized theory in a language L_T . The following are equivalent.*

- (i) T is interpretable in a consistent existential theory.
- (ii) T has a quantifier-free interpretation I in a consistent quantifier-free theory S such that I and the language of S obey the conditions in Corollary 4.4, except that I may be multi-dimensional if L_T contains a proper function symbol.

(iii) T is weakly interpretable in EC_L for some language L , w.l.o.g. obeying the same conditions as in (ii).

If L_T is finite, and T is recursively axiomatized, we can also make the interpreting theories recursively axiomatized.

Proof: (ii) \rightarrow (i) is trivial, and (i) \rightarrow (iii) follows from the fact that every consistent existential theory is consistent with EC_L in the same language by Theorem 3.7.

(iii) \rightarrow (ii): By expanding L and L_T with Henkin constants for the existential quantifiers in axioms of T using Lemma 4.1 (ii), we may assume that T is universal. By Corollary 4.4, T has a one-piece one-dimensional parameter-free unrelativized interpretation J in a consistent theory $EC_L + S$, where L consists of a binary function and constants. By quantifier elimination, we may assume S is a set of quantifier-free sentences, and the J -translations of equality and all symbols of L_T are given by quantifier-free formulas. By expanding the language L further, we may assume that constants of L_T are interpreted by constants (or constant terms) of L . In the countable case, we may apply Lemma 4.3 again to reduce the number of constants to one.

If L_T contains proper function symbols, we need more work, as we cannot add Skolem functions in the same way as constants. As we will explain in the appendix, EC_L has weak elimination of imaginaries, and as a corollary, we obtain in Proposition A.9 an explicit description of definable equivalence relations that we apply to $=^J$. Using (13), we see that the collection of equivalence classes that make up the domain of J can be definably split in finitely many pieces, where the i -th piece is in definable bijection with a collection of m_i -element sets of r -tuples (represented by an equivalence relation on $m_i r$ -tuples as in Definition A.1). The upshot is that we may replace J with an equivalent piece-wise interpretation I that almost has absolute equality, in the sense that all equivalence classes of $=^I$ have bounded finite size. Consequently, the translation F^I of any function symbol $F \in L_T$, when viewed as a relation on tuples rather than on their equivalence classes, is a total multifunction with only finitely many values. By Lemma A.2 and a compactness argument, there is a piecewise term-definable function that picks one possible value of such a multivalued function. Thus, I is a quantifier-free interpretation. Since we may assume L includes a pair of constants c, d such that $S \vdash c \neq d$, we can make I a one-piece interpretation; it is still parameter-free, and we can make it unrelativized as above, but it may be multi-dimensional. (Lemma 4.1 does not give a pairing *function*, hence it is unclear if we can make the interpretation one-dimensional without sacrificing the property that translations of functions are piecewise term-definable.)

The result of these manipulations is that T^I is a *universal* subtheory of $EC_L + S$, as we made sure all existential quantifiers needed are witnessed (piecewise) by terms. Thus, T^I is in fact included in S , i.e., I is an interpretation of T in S which satisfies all the requirements.

Finally, let T be an r.e. theory in a finite language. We have shown that if T is interpretable in a consistent existential theory, there is a quantifier-free unrelativized one-piece parameter-free interpretation of T in a consistent extension of EC_{L_2} , where L_2 consists of a constant and a binary function. (The interpretation is automatically recursive, as the language is finite.) The universal Henkin expansion T^H of T is still r.e., and we can assign the Henkin constants in a recursive way to new constants added to L_2 so that we get an interpretation I with the same properties of T^H in a consistent extension of EC_L , where L consists of a binary function and countably many constants, and I is recursive. The cointerpretation from Lemma 4.3 is also recursive, hence we can reduce the language back to L_2 . Then $EC_{L_2} + (T^H)^I$ is an r.e. theory, hence by effectiveness of quantifier elimination, it is equivalent to $EC_{L_2} + S$ for an r.e. quantifier-free L_2 -theory S . By the argument above, I is an interpretation of T in S , as $(T^H)^I$ is a universal theory. \square

Remark 4.6 Theorem 4.5 does not extend to $\forall\exists$ theories T . On the one hand, any theory interpretable in a consistent existential theory is locally finitely satisfiable (notice also that any consistent $\exists\forall$ theory in a relational language is locally finitely satisfiable). On the other hand, EC_L itself is a $\forall\exists$ theory interpretable in EC_L , and if L contains a nonconstant function symbol, then EC_L is not locally finitely satisfiable: for example, if we have a unary function $F(x)$, then EC_L proves the formula

$$\forall x, y \exists z (z \neq x \wedge F(z) = y)$$

with no finite model.

We note that if L contains only at most unary relations and constants, then EC_L and any its consistent extension is an existential theory, and easily seen to be interpretable in REP_U for L finite. If L consists of relations and constants, but is not unary, then EC_L (i.e., essentially the theory of the random structure) is genuinely $\forall\exists$, but still locally finitely satisfiable, hence interpretable in R for L finite by Visser's Theorem 2.11.

Question 4.7 *Is every consistent r.e. existential theory interpretable in REP_{PRF} ?*

Question 4.8 *Is the theory of the random graph interpretable in a consistent existential theory?*

5 Classification of EC_L

We now proceed to the main results of the paper, showing that certain theories are not interpretable in any existentially axiomatized theory by way of establishing tameness properties of EC_L . We will mostly deduce them from the following statement, showing the impossibility of certain configurations in models of EC_L .

In order to keep the proof self-contained and accessible to wider audience, we will not use any results on indiscernibles (though they are lurking in our application of Ramsey's theorem).

Recall that a relation $R \subseteq X^2$ is *asymmetric* if there are no $a, b \in X$ such that $R(a, b) \wedge R(b, a)$.

Theorem 5.1 *For any language L and formula $\varphi(\bar{z}, \bar{x}, \bar{y})$ with $\text{lh}(\bar{x}) = \text{lh}(\bar{y})$, there is a constant n with the following property. Let $M \models EC_L$ and $\bar{u} \in M$ be such that*

$$M \models \exists \bar{x}_0, \dots, \bar{x}_{n-1} \bigwedge_{i < j < n} \varphi(\bar{u}, \bar{x}_i, \bar{x}_j).$$

Then for every $m \in \omega$ and an asymmetric relation R on $\{0, \dots, m-1\}$,

$$M \models \exists \bar{x}_0, \dots, \bar{x}_{m-1} \bigwedge_{\langle \alpha, \beta \rangle \in R} \varphi(\bar{u}, \bar{x}_\alpha, \bar{x}_\beta).$$

Proof: By Theorem 3.7 and Lemma 4.1, we may assume L contains no relations, all the tuples have length one, and φ is open. Let τ be the number of subterms of φ , and $\tau^* = 2^{256\tau^2}$. Using Ramsey's theorem, let n be sufficiently large so that

$$n \rightarrow (\tau^*)_{\tau^*}^4.$$

Fix $M \models EC_L$, $u \in M$, and $\{a_i : i < n\} \subseteq M$ such that $M \models \varphi(u, a_i, a_j)$ for $i < j < n$. In order to simplify the notation, we will assume u is given by a constant of L , and write just $\varphi(x, y)$; this does not increase the number of subterms of φ . Let S be the set of all subterms $t(x, y)$ of φ , and for every $i_0 < i_1 < i_2 < i_3 < n$, define

$$\text{tp}(i_0, i_1, i_2, i_3) = \{\langle u_0, u_1, u_2, u_3, t, s \rangle \in \mathbf{4}^4 \times S^2 : t^M(a_{i_{u_0}}, a_{i_{u_1}}) = s^M(a_{i_{u_2}}, a_{i_{u_3}})\}.$$

Since $|S| \leq \tau$, tp is a colouring of quadruples of numbers below n by at most τ^* colours. Thus, we can find a 7-element homogeneous set $H \subseteq \{0, \dots, n-1\}$ for tp ; without loss of generality $H = \{0, \dots, 6\}$.

Fix a set of variables $\{y_\alpha : \alpha < m\}$, and put

$$\Theta = \{t(y_\alpha, y_\beta) : \langle \alpha, \beta \rangle \in R, t(x, y) \in S\}.$$

If $t \in \Theta$, let $V(t)$ denote the set of $\alpha < m$ such that y_α occurs in t ; note that $|V(t)| \leq 2$. A *realization* of t is an injective mapping $r: V(t) \rightarrow H$ such that

$$\alpha, \beta \in V(t) \wedge \langle \alpha, \beta \rangle \in R \implies r(\alpha) < r(\beta).$$

Notice that this condition is void if t depends on at most one variable; otherwise it concerns a unique pair $\langle \alpha, \beta \rangle$. If r is a realization of t , let $r(t) \in M$ be the value of the term resulting from t by replacing each variable y_α with $a_{r(\alpha)}$.

A *joint realization* of a set of terms $\{t_0, \dots, t_{k-1}\} \subseteq \Theta$ is an injective mapping $r: V(t_0) \cup \dots \cup V(t_{k-1}) \rightarrow H$ such that $r \upharpoonright V(t_i)$ is a realization of t_i for $i < k$. Note that any pair $\{t, s\} \subseteq \Theta$ has a joint realization, as R has no cycles of length at most 2.

If $t, s \in \Theta$, and r is a joint realization of t and s , we define

$$t \sim s \iff r(t) = r(s).$$

Claim 5.1.1 *The definition of \sim is independent of the choice of r .*

Proof: First, if two joint realizations r, r' satisfy

$$r(\alpha) < r(\beta) \iff r'(\alpha) < r'(\beta)$$

for all $\alpha, \beta \in V(t) \cup V(s)$, then

$$r(t) = r(s) \iff r'(t) = r'(s) \tag{10}$$

by homogeneity for tp . This condition holds automatically if

- $V(t) \subseteq V(s)$ or $V(s) \subseteq V(t)$, or
- $V(t) = \{\alpha, \beta\}$, $V(s) = \{\beta, \gamma\}$, where $\langle \alpha, \beta \rangle, \langle \beta, \gamma \rangle \in R$, or vice versa.

Assume $t = t(y_\alpha, y_\beta)$, $s = s(y_\alpha, y_\gamma)$, where $\beta \neq \gamma$, and $\langle \alpha, \beta \rangle, \langle \alpha, \gamma \rangle \in R$ (the case with $\langle \beta, \alpha \rangle, \langle \gamma, \alpha \rangle \in R$ is symmetric). By (10), it suffices to consider the case where $r(\alpha) = r'(\alpha) = 0$, $r(\beta) = 1$, $r(\gamma) = r'(\gamma) = 2$, $r'(\beta) = 3$. Using (10), we have

$$t^M(a_0, a_1) = s^M(a_0, a_2) \implies t^M(a_0, a_3) = s^M(a_0, a_4) = t^M(a_0, a_1) = s^M(a_0, a_2),$$

and the converse implication is symmetric.

The remaining case is when $V(t)$ and $V(s)$ are disjoint and nonempty. It suffices to show that if $r(t) = r(s)$ for some joint realization r , there is a constant $a \in M$ such that $r'(t) = a$ for every realization r' of t (whence the same holds for s by symmetry). Assume t depends on two variables y_α, y_β with $\langle \alpha, \beta \rangle \in R$ (the unary case is easier). Using (10), we may assume that the realization $r_1(\alpha) = r(\alpha) - 1$, $r_1(\beta) = r(\beta) + 1$ of t is within bounds, and disjoint from $r(V(s))$, thus $r_1 \cup (r \upharpoonright V(s))$ is a joint realization of t, s ordered in the same way as r . It follows that

$$t^M(a_{r(\alpha)-1}, a_{r(\beta)+1}) = r(s) = t^M(a_{r(\alpha)}, a_{r(\beta)}).$$

Applying homogeneity, we have

$$t^M(a_i, a_j) = t^M(a_0, a_6) = t^M(a_k, a_l)$$

for every $0 < i < j < 6$, $0 < k < l < 6$. Since every set $\{i, j, k, l\}$ is order-isomorphic to some not involving 0, 6, we obtain

$$t^M(a_i, a_j) = t^M(a_k, a_l)$$

for all $i < j$, $k < l$ using homogeneity again. □ (Claim 5.1.1)

Thus, \sim is a well-defined relation on Θ . It is clearly reflexive and symmetric. If $t \simeq F(t_0, \dots, t_{k-1})$ and $s \simeq F(s_0, \dots, s_{k-1})$ are in Θ , and r is a joint realization of t and s , it is also a joint realization of each $\{t_i, s_i\}$, hence

$$t_0 \sim s_0, \dots, t_{k-1} \sim s_{k-1} \implies t \sim s.$$

Claim 5.1.2 \sim is transitive.

Proof: Assume that $t \sim s \sim u$. If there exists a joint realization r of $\{t, s, u\}$, we immediately obtain $r(t) = r(s) = r(u)$, hence $t \sim u$. If not, we must have $t = t(y_\alpha, y_\beta)$, $s = s(y_\beta, y_\gamma)$, $u = u(y_\gamma, y_\alpha)$, where $\langle \alpha, \beta \rangle, \langle \beta, \gamma \rangle, \langle \gamma, \alpha \rangle \in R$. Applying alternately $t \sim s$ and $s \sim u$, we obtain

$$t^M(a_3, a_4) = s^M(a_4, a_5) = u^M(a_5, a_6) = s^M(a_1, a_5) = t^M(a_0, a_1) = s^M(a_1, a_2) = u^M(a_2, a_3),$$

hence $r(t) = r(u)$ under the joint realization of t, u such that $r(\gamma) = 2$, $r(\alpha) = 3$, and $r(\beta) = 4$. (This argument in fact shows that with such a cyclic dependency, the values of all three terms are independent of the realization.) □ (Claim 5.1.2)

Let $\Xi \subseteq \Theta$ and $\{t^* : t \in \Xi\}$ be as in Definition 3.1 (for empty \bar{x}). By induction on the definition of $t \in \Xi$, we see that the value of the closed term t^* in M coincides with $r(t)$ for any realization r of t . This and the definition of \sim implies that

$$M \models \theta_{\Theta, \sim}^*$$

hence by Lemma 3.5 and existential closedness of M ,

$$M \models \exists y_0, \dots, y_{m-1} \theta_{\Theta, \sim}(\bar{y}).$$

If $b_0, \dots, b_{m-1} \in M$ witness this, and $\langle \alpha, \beta \rangle \in R$ and $i < j \in H$, we have

$$M \models \psi(b_\alpha, b_\beta) \iff M \models \psi(a_i, a_j)$$

for every subformula ψ of φ . It follows that

$$M \models \bigwedge_{\langle \alpha, \beta \rangle \in R} \varphi(b_\alpha, b_\beta)$$

as required. □

We draw two principal conclusions from Theorem 5.1. For the first one, notice that the theory below is interpretable in the theory R just by taking $<$ for \in : then (11) is witnessed by $x_i = \underline{i}$, $z = \underline{n}$ due to axiom (4).

Corollary 5.2 *The theory in the language $\langle \in \rangle$ axiomatized by the sentences*

$$\exists z, x_0, \dots, x_{n-1} \left(\bigwedge_{i < j < n} x_i \neq x_j \wedge \forall y \left(y \in z \leftrightarrow \bigvee_{i < n} y = x_i \right) \right) \quad (11)$$

for all $n \in \omega$ is not weakly interpretable in EC_L , and consequently not interpretable in any consistent existential theory.

Proof: Apply Theorem 5.1 to the formula interpreting $x \in z \wedge y \in z \wedge x \neq y$, and R a chain longer than n . □

We can restate this in proper model-theoretic terminology. A theory T is said to *eliminate* \exists^∞ (or *eliminate infinity*) if for every formula $\varphi(\bar{z}, x)$, there exists n such that for every model $M \models T$ and $\bar{a} \in M$, if $|\varphi(\bar{a}, M)| \geq n$, then it is infinite. Specializing this to the theory T^{eq} , this means that for every $\varphi(\bar{z}, \bar{x})$ and $\psi(\bar{x}, \bar{y})$ (where $\text{lh}(\bar{x}) = \text{lh}(\bar{y}) = k$), there exists n such that for every $M \models T$ and $\bar{a} \in M$, if ψ defines an equivalence relation on M^k , and $\varphi(\bar{a}, M^k)$ hits at least n equivalence classes, then it hits infinitely many.

As in Corollary 5.2, an application of Theorem 5.1 to the formula $\varphi(\bar{z}, \bar{x}) \wedge \varphi(\bar{z}, \bar{y}) \wedge \neg\psi(\bar{x}, \bar{y})$ yields:

Corollary 5.3 $(EC_L)^{\text{eq}}$ has elimination of the \exists^∞ quantifier. □

Our second principal conclusion is the following tameness result on EC_L :

Corollary 5.4 For any language L , EC_L has NSOP₃. That is, the theory axiomatized by

$$\forall x, y, z \neg(x < y \wedge y < z \wedge z < x),$$

$$\exists x_0, \dots, x_{n-1} \bigwedge_{i < j < n} x_i < x_j$$

for $n \in \omega$ is not weakly interpretable in EC_L , and is not interpretable in any consistent existential theory.

Consequently, EC_L has the NSOP property, i.e., no theory consistent with EC_L interprets a partial order with arbitrarily long chains.

Proof: Apply Theorem 5.1 with R being a directed 3-cycle. □

Let us state for the record that Corollaries 5.2 or 5.4 solve our original problem:

Corollary 5.5 The theory REP_{PRF} represents all partially recursive functions, but it does not interpret R . □

On the other hand, it should be stressed that EC_L is not *that* tame, if the language L is sufficiently complicated (note that the observation below also stands in contrast to properties of random relational structures, i.e., EC_L with L purely relational, which is a simple theory, thus NTP₂).

Proposition 5.6 If L contains an at least binary function symbol, then EC_L has TP₂, hence it is not simple. More generally, any theory weakly interpreting REP_{TRF} (i.e., with a consistent extension that loosely represents *trf*) has TP₂.

Proof: It suffices to show the latter claim. Let $\underline{F}(x, y)$ be the REP_{TRF} -function representing the recursive function that interprets x as a Gödel number of a finite sequence, and outputs its y th element. Let $\bar{a}_{i,j} = (\underline{i}, \underline{j})$, and $\varphi(x, y_1, y_2)$ be the formula

$$\underline{F}(x, y_1) = y_2.$$

Clearly, $\varphi(x, \bar{a}_{i,j}) \wedge \varphi(x, \bar{a}_{i,k})$ is inconsistent for $j \neq k$. On the other hand, if $\sigma \in \omega^\omega$, and $n \in \omega$, let s be the Gödel number of $\sigma \upharpoonright n$. Then $\varphi(\underline{s}, \bar{a}_{i, \sigma(i)})$ for all $i < n$. Thus, the type

$$\{\varphi(x, \bar{a}_{i, \sigma(i)}) : i < \omega\}$$

is consistent. □

The assumption on L in Proposition 5.6 is essential; see Theorem B.1 for more detailed model-theoretic classification of the theories EC_L as L varies.

Now, in the most general case when L contains an at least binary function symbol, there is still a gap left between Proposition 5.6 and Corollary 5.4. We can close it by improving Corollary 5.4 from NSOP₃ to NSOP₁, but the proof will no longer be self-contained: we will rely on a characterization of NSOP₁ theories due to Chernikov and Ramsey [2] using an independence relation in the spirit of the Kim–Pillay theorem.

We will work inside a *monster model* \mathbb{M} of a completion $T \supseteq EC_L$, as in Section 2.4: a κ -saturated, strongly κ -homogeneous model of T , where κ is a cardinal larger than $\|L\|$ and all structures we intend to handle; recall that subsets of \mathbb{M} of size $< \kappa$ are called small.

Definition 5.7 If A , B , and C are small tuples (sequences), we say that A is *independent from B over C* , written as $A \downarrow_C B$, if $\langle AC \rangle \cap \langle BC \rangle = \langle C \rangle$, where $\langle X \rangle$ denotes the substructure generated by X , and the juxtaposition of two sequences denotes their concatenation. We will often treat these tuples as sets where the context permits, seeing as the definition of \downarrow does not depend on their ordering.

The definition of $A \downarrow_C B$ is stated here in more general circumstances than what is required for [2] (in particular, their characterization only needs the case when C is a small *model*, i.e., an elementary submodel of the monster). We do it partly because we can—at no additional cost—and partly because we also want the definition to conform to the shape of independence relations from the original Kim–Pillay theorem, which we will use elsewhere in the paper. For the same reason, the next Lemma includes properties of independence relations that are not directly relevant to the characterization from [2].

Lemma 5.8 *Let \mathbb{M} be a monster model of a completion T of EC_L . The independence relation \downarrow has the following properties for all small tuples A, A', B, B', C, D :*

- (i) (*Invariance*) If f is an automorphism of \mathbb{M} , then $A \downarrow_C B$ implies $f(A) \downarrow_{f(C)} f(B)$.
- (ii) (*Symmetry*) $A \downarrow_C B$ implies $B \downarrow_C A$.
- (iii) (*Monotonicity*) If $A' \subseteq A$, and $B' \subseteq B$, then $A \downarrow_C B$ implies $A' \downarrow_C B'$.
- (iv) (*Weak transitivity*) $A \downarrow_B C$ and $A \downarrow_{BC} D$ implies $A \downarrow_B CD$.
- (v) (*Existence*) $A \downarrow_B B$.
- (vi) (*Strong finite character*) If $A \not\downarrow_C B$, there is a formula $\varphi(\bar{x}, \bar{b}, \bar{c}) \in \text{tp}(A/BC)$ such that $\bar{a} \not\downarrow_C B$ whenever $\mathbb{M} \models \varphi(\bar{a}, \bar{b}, \bar{c})$.
- (vii) (*Extension*) For any A, B, C , there is $A' \equiv_C A$ such that $A' \downarrow_C B$.
- (viii) (*Local character*) For any B , and finite A , there is $B' \subseteq B$ such that $|B'| \leq \|L\|$, and $A \downarrow_{B'} B$.
- (ix) (*Independence theorem*) If $A \downarrow_C B$, $B \downarrow_C B'$, $B' \downarrow_C A'$, and $A' \equiv_C A$, there exists A'' such that $A'' \equiv_{CB} A$, $A'' \equiv_{CB'} A'$, and $A'' \downarrow_C BB'$.

Proof: Properties (i)–(v) are clear.

(vi): By definition, $A \not\downarrow_C B$ implies that $t(\bar{a}, \bar{c}) = s(\bar{b}, \bar{c}) \notin \langle C \rangle$ for some terms t, s , and $\bar{a} \subseteq A$, $\bar{b} \subseteq B$, $\bar{c} \subseteq C$. Then we can take $t(\bar{x}, \bar{c}) = s(\bar{b}, \bar{c})$ for the formula φ .

(vii): We can extend the structure $\langle BC \rangle$ with a disjoint copy of $\langle AC \rangle \setminus \langle C \rangle$: that is, let us define a structure D with domain

$$\langle BC \rangle \dot{\cup} \{\bar{x} : x \in \langle AC \rangle \setminus \langle C \rangle\},$$

with relations and functions defined so that they agree with the original structure on $\langle BC \rangle$, and so that $f = \text{id}_{\langle C \rangle} \cup \bar{x}$ is an isomorphism of $\langle AC \rangle$ to $\langle C \rangle \cup \overline{\langle AC \rangle \setminus \langle C \rangle} \subseteq D$. We can ensure D is a substructure of \mathbb{M} (extending $\langle BC \rangle$) using EC_L and κ -saturation. Then $A' = f(A)$ has the required properties.

(viii): Let us construct a chain $B_0 \subseteq B_1 \subseteq B_2 \subseteq \dots$ of subsets of B of size $|B_n| \leq \lambda := \|L\|$ as follows. We put $B_0 = \emptyset$. Given B_n , let $M_n = \langle AB_n \rangle \cap \langle B \rangle$. Since $|M_n| \leq \lambda$, there exists a set $B_n \subseteq B_{n+1} \subseteq B$ of size $|B_{n+1}| \leq \lambda$ such that $M_n \subseteq \langle B_{n+1} \rangle$.

Let $B' = \bigcup_{n \in \omega} B_n$. Then $\langle AB' \rangle \cap \langle B \rangle = \bigcup_n M_n \subseteq \langle B' \rangle$ by construction, hence $A \perp_{B'} B$, and $|B'| \leq \lambda$.

(ix): In order to simplify the notation, we may assume without loss of generality that C, A, A', B , and B' are structures, with $C \subseteq A, A', B, B'$. By the assumption, we have $A \cap B = A' \cap B' = B \cap B' = C$, and we can fix an isomorphism $f: A \simeq A'$ identical on C . Put $D = \langle AA'BB' \rangle$. We will extend D into a model D' with domain

$$D' = D \dot{\cup} \{\bar{x} : x \in A \setminus C\} \dot{\cup} \{\underline{x} : x \in \langle AB \rangle \setminus (A \cup B)\} \dot{\cup} \{\underline{x} : x \in \langle A'B' \rangle \setminus (A' \cup B')\}$$

using copies of parts of D . We will also write \bar{y} for elements $y \in A' \setminus C$, so that $\bar{x} = \bar{y}$ if $f(x) = y$.

We define relations and functions on D' so that

- $g = \text{id}_B \cup \bar{x} \cup \underline{x}$ is an isomorphism of $\langle AB \rangle$ to $B \cup \overline{A \setminus C} \cup \langle AB \rangle \setminus (A \cup B)$, and
- $g' = \text{id}_{B'} \cup \bar{x} \cup \underline{x}$ is an isomorphism of $\langle A'B' \rangle$ to $B' \cup \overline{A' \setminus C} \cup \langle A'B' \rangle \setminus (A' \cup B')$.

It is important to note there is no conflict between the two clauses: the intersection of the two targets is $C \cup \overline{A \setminus C} = C \cup \overline{A' \setminus C}$, which is asked to be made isomorphic to A via $\text{id}_C \cup \bar{x}$, and to A' via $\text{id}_C \cup \bar{x} = \text{id}_C \cup f^{-1}(\bar{x})$; these two requirements are equivalent, as the two mappings commute with the isomorphism $f: A \rightarrow A'$.

Now, using EC_L and κ -saturation, we can embed D' as a substructure of \mathbb{M} extending D . Let $A'' = g(A) = g'(A')$. Then g is an isomorphism of A to A'' identical on $B \supseteq C$, thus $A'' \equiv_{CB} A$, and similarly $A'' \equiv_{CB'} A'$ via g' . Finally, $\langle A''C \rangle = A''$, $\langle BB'C \rangle \subseteq D$, and $A'' \cap D = C$, thus $A'' \perp_C BB'$. \square

The following is a restatement of Proposition 5.8 in Chernikov and Ramsey [2].

Theorem 5.9 *Let \mathbb{M} be a monster model of a complete theory T , and $A \perp_M B$ an independence relation on small tuples A, B , and small models $M \models T$, that satisfies the appropriate restrictions of properties (i), (ii), (iii), (v), (vi), and (ix) from Lemma 5.8. Then T is NSOP₁.* \square

Corollary 5.10 *For any language L , EC_L is NSOP₁.* \square

Remark 5.11 Using Theorem 9.1 in [4], it can be seen that our \perp coincides with the relation of *Kim independence*.

We mention that Corollary 5.10 was independently discovered by Kruckman and Ramsey [6], who learned of the problem from an earlier unpublished version of this paper where it was posed as an open problem.

6 Conclusion

We succeeded in our original goal of separating interpretability of R from representability of recursive functions. More generally, we obtained a criterion for interpretability of $\exists\forall$ theories in existential theories, showing in particular that we may assume such interpretations to be quantifier-free. We believe these results are interesting in their own right, of course, but at the same time we place as much value on the connection between formal arithmetic and model theory that it revealed: while model-theoretic methods are often used in the study of arithmetic, typically this means to work with models of the (fairly strong) theories of arithmetic themselves, which are quite unlike the kind of tame model theory we encountered in this paper. It would be interesting to see if more such connections are waiting to be discovered.

Acknowledgement

The question that led to this paper arose from a fruitful discussion with Albert Visser. I would also like to thank Nick Ramsey for clarification of his work and other comments on the topic.

References

- [1] Enrique Casanovas and Rafael Farré, *Weak forms of elimination of imaginaries*, Mathematical Logic Quarterly 50 (2004), no. 2, pp. 126–140.
- [2] Artem Chernikov and Nicholas Ramsey, *On model-theoretic tree properties*, Journal of Mathematical Logic 16 (2016), no. 2, 1650009 (41 pages).
- [3] Wilfrid Hodges, *A shorter model theory*, Cambridge University Press, Cambridge, 1997.
- [4] Itay Kaplan and Nicholas Ramsey, *On Kim-independence*, arXiv:1702.03894 [math.LO], 2017, <http://arxiv.org/abs/1702.03894>.
- [5] Byunghan Kim and Anand Pillay, *Simple theories*, Annals of Pure and Applied Logic 88 (1997), no. 2–3, pp. 149–164.
- [6] Alex Kruckman and Nicholas Ramsey, *Generic expansion and Skolemization in NSOP₁ theories*, arXiv:1706.06616 [math.LO], 2017, <http://arxiv.org/abs/1706.06616>.
- [7] Pavel Pudlák, *Cuts, consistency statements and interpretations*, Journal of Symbolic Logic 50 (1985), no. 2, pp. 423–441.
- [8] Shaharon Shelah, *Classification theory and the number of non-isomorphic models*, Studies in Logic and the Foundations of Mathematics vol. 92, North-Holland Publishing Company, Amsterdam, New York, 1978, second edition 1990.
- [9] Alfred Tarski, Andrzej Mostowski, and Rafael M. Robinson, *Undecidable theories*, North-Holland, Amsterdam, 1953.
- [10] Albert Visser, *Pairs, sets and sequences in first-order theories*, Archive for Mathematical Logic 47 (2008), no. 4, pp. 299–326.
- [11] ———, *What is the right notion of sequentiality?*, <http://www.phil.uu.nl/preprints/lgps/number/288>, 2010.

- [12] ———, *Why the theory R is special*, in: Foundational Adventures: Essays in Honor of Harvey M. Friedman (N. Tennant, ed.), Tributes vol. 22, College Publications, London, 2014, originally published online by Templeton Press, <http://foundationaladventures.com>, 2012.
- [13] Frank O. Wagner, *Simple theories*, Mathematics and Its Applications vol. 503, Kluwer, 2000.
- [14] Peter M. Winkler, *Model-completeness and Skolem expansions*, in: Model Theory and Algebra: A Memorial Tribute to Abraham Robinson (D. H. Saracino and V. B. Weispfenning, eds.), Lecture Notes in Mathematics vol. 498, Springer, 1975, pp. 408–463.

A Elimination of imaginaries

In this section, we discuss elimination of imaginaries in the theories EC_L . We put it here in the appendix as it is rather tangential to our main topic; we only need it in the proof of Theorem 4.5.

One way to describe elimination of imaginaries is that, loosely speaking, it allows to replace any interpretation with an interpretation with absolute equality. We recall the proper definition below, along with some important variants of the notion.

Definition A.1 A theory T has *elimination of imaginaries (e.i.)* if for every $M \models T$ and $e \in M^{\text{eq}}$, there is a tuple $\bar{b} \in M$ such that $\text{dcl}_{M^{\text{eq}}}(e) = \text{dcl}_{M^{\text{eq}}}(\bar{b})$.

T has *weak e.i.* if for every $M \models T$ and $e \in M^{\text{eq}}$, there is $\bar{b} \in M$ such that $\bar{b} \in \text{acl}_{M^{\text{eq}}}(e)$ and $e \in \text{dcl}_{M^{\text{eq}}}(\bar{b})$.

T has *Galois e.i.* if for every $M \models T$, $\bar{a} \in M$, and $e \in \text{acl}_{M^{\text{eq}}}(\bar{a})$, there is $\bar{b} \in M$ such that $\text{dcl}_{M^{\text{eq}}}(\bar{a}e) = \text{dcl}_{M^{\text{eq}}}(\bar{a}\bar{b})$.

As a special case, for any $k, l > 0$ let $\sim_{k,l}$ be the equivalence relation on injective l -tuples of k -tuples (represented as kl -tuples) defined by

$$\langle \bar{a}_i : i < l \rangle \sim_{k,l} \langle \bar{b}_i : i < l \rangle \iff \{\bar{a}_i : i < l\} = \{\bar{b}_i : i < l\},$$

so that $M^{kl}/\sim_{k,l}$ represents l -element subsets of M^k . If for all $M \models T$ and all $k, l > 0$, every $a \in M^{kl}/\sim_{k,l}$ is interdefinable (in M^{eq}) with a tuple $\bar{b} \in M$, then T has *coding of finite sets*. See [1] for an exposition of various forms of e.i.; in particular, T has e.i. iff it has weak e.i. and coding of finite sets.

It would be nice if EC_L had e.i., however this is too good to be true: as we will prove shortly, the theory does not have coding of finite sets.

Lemma A.2 *If $M \models EC_L$, and $A \subseteq M$, then $\text{acl}_M(A) = \langle A \rangle$.*

Proof: Assume A is a submodel, and $M \models \varphi(b, \bar{a})$, where $b \in M \setminus A$, and φ is open. Let $N = M \dot{\cup} (\omega \times (M \setminus A))$, where each $A \cup (\{i\} \times (M \setminus A))$ is an isomorphic copy of M , as in the proof of Lemma 5.8 (vii). Then $N \models \varphi(\langle i, b \rangle, \bar{a})$ for each $i < \omega$, hence $\varphi(M, \bar{a})$ is also infinite by existential closedness. \square

Lemma A.3 *Every $M \models EC_L$ has elementary extensions with arbitrarily large sets X totally indiscernible over M . Moreover, we can choose X to have the additional property that any L_M -term $t(x_1, \dots, x_k)$ containing all the indicated variables defines an injective function on X^k .*

Proof: Let $M \models EC_L$, and X be a set disjoint from M . An L_M -term in variables X is *reduced* if it has no constant subterms other than M -constants; that is, variables and M -constants are reduced,

and if t_1, \dots, t_k are reduced terms, and F is a k -ary function symbol of L , then $F(t_1, \dots, t_k)$ is reduced unless all the t_i are M -constants.

Let R be the model whose domain is the set of all reduced terms, with realizations of relations the same as in M (i.e., unsatisfied by tuples involving any non-constant terms), and functions realized in the obvious way. Let N be an e.c. extension of R . Since every permutation of X extends to an automorphism of R fixing M , any tuples of distinct elements x_1, \dots, x_k and x'_1, \dots, x'_k of X satisfy the same atomic formulas with parameters from M . Thus in N , X is a totally indiscernible set over M by quantifier elimination. \square

Proposition A.4 *Neither EC_L nor any its completion has coding of unordered pairs of elements, and a fortiori elimination of imaginaries.*

Proof: Let N be a model of EC_L with $\{a, b\}$ a 2-element totally indiscernible set satisfying the property from Lemma A.3. Without loss of generality, N is strongly ω -homogeneous. Assume for contradiction that there is a tuple \bar{u} in N interdefinable with the representation of $\{a, b\}$ (that is, with the element $\langle a, b \rangle / \sim_{1,2}$ of N^{eq}). By Lemma A.2, all elements of \bar{u} are in the submodel generated by a, b , hence they are given by a tuple of terms $\bar{t}(a, b)$. By $a, b \equiv b, a$ and homogeneity, there is an automorphism f such that $f(a) = b$ and $f(b) = a$; since f preserves $\{a, b\}$, it also preserves \bar{u} , hence $t_i(a, b) = t_i(b, a)$. Using the extra property, this can only happen if all the t_i are closed terms. Thus \bar{u} , hence $\{a, b\}$, is \emptyset -definable, and $a, b \in \text{acl}_N(\emptyset)$; using Lemma A.2 again, a and b are in fact values of closed terms, but this contradicts $a \equiv b$. \square

Short of full e.i., the next best thing we can hope for is weak e.i. This will turn out to hold for EC_L , and thankfully it is still enough for our intended application.

Let us work again in a monster model \mathbb{M} of a completion of EC_L . Recall the following characterization [1, Facts 1.2]: weak e.i. holds iff for every relation R definable with parameters in \mathbb{M} , there exists a smallest algebraically closed set defining R . (By Lemma A.2, algebraically closed set = substructure for us.) We observe easily that EC_L satisfies a somewhat weaker property:

Lemma A.5 *For any definable relation $R(\bar{x})$, the class of substructures that define R is directed. That is, if R is definable over \bar{b} , and over \bar{c} , it is also definable over $\langle \bar{b} \rangle \cap \langle \bar{c} \rangle$.*

Proof: In order to simplify the notation, we will omit bars over finite tuples.

Put $A = \langle b \rangle \cap \langle c \rangle$. By saturation, it suffices to show that

$$x \equiv_A x' \implies (R(x) \leftrightarrow R(x')). \quad (12)$$

Assume that $R(x) \leftrightarrow \varphi(b, x) \leftrightarrow \psi(c, x)$.

Claim A.5.1

- (i) *If $b \equiv_c b'$, then $R(x) \leftrightarrow \varphi(b', x)$.*
- (ii) *If $c \equiv_b c'$, and $b \equiv_{c'} b'$, then $R(x) \leftrightarrow \varphi(b', x)$.*
- (iii) *If $b \equiv_A b'$, and $x \perp_A b'$, then $R(x) \leftrightarrow \varphi(b', x)$.*
- (iv) *If $x \equiv_A x'$, and $x \perp_A b$, then $R(x) \leftrightarrow R(x')$.*

Proof: (i): Let $\alpha(v, w)$ denote $\forall x (\varphi(v, x) \leftrightarrow \psi(w, x))$. Since $\alpha(b, c)$, and $b \equiv_c b'$, we have $\alpha(b', c)$.

(ii): By the dual statement to (i), we have $R(x) \leftrightarrow \psi(c', x)$. Thus, $\alpha(b, c')$, whence $\alpha(b', c')$.

(iii): By Lemma 5.8 (vii), there exists $c' \equiv_b c$ such that $c' \downarrow_b x$. Since $c \downarrow_A b$ by the definition of A , $c' \equiv_b c$ implies $c' \downarrow_A b$. By Lemma 5.8 (iv), we obtain $c' \downarrow_A x$. Thus, by Lemma 5.8 (ix), there exists b'' such that $b'' \equiv_{c'} b$, and $b'' \equiv_{Ax} b'$. Using (ii), this implies

$$R(x) \leftrightarrow \varphi(b'', x) \leftrightarrow \varphi(b', x).$$

(iv): There exists b' such that $xb \equiv_A x'b'$. Then $x' \downarrow_A b'$, thus

$$R(x) \leftrightarrow \varphi(b, x) \leftrightarrow \varphi(b', x') \leftrightarrow R(x')$$

by (iii). □ (Claim A.5.1)

Finally, to prove (12), let $x'' \equiv_A x$ be such that $x'' \downarrow_A b$ using Lemma 5.8 (vii). Then $R(x) \leftrightarrow R(x'') \leftrightarrow R(x')$ by (iv) of the Claim. □

Incidentally, the previous lemma implies another variant of e.i.:

Proposition A.6 *For any language L , EC_L has Galois elimination of imaginaries.*

Proof: By [1, Prop. 3.9], Galois e.i. is equivalent to the conjunction of elimination of strong types (ST), and coding of Galois finite sets. The latter follows from our Lemma A.2 by [1, Facts 3.7]. It thus suffices to show ST; by [1, Prop. 3.2], this is equivalent (in view of Lemma A.2) to the claim that $\text{acl}_{M^{\text{eq}}}(A) = \text{dcl}_{M^{\text{eq}}}(A)$ for real sets A in a monster model $\mathbb{M} \models EC_L$. We can reformulate this as follows: if E is an equivalence relation (on k -tuples) definable over A with finitely many equivalence classes, then all the equivalence class are individually definable over A .

So, let us fix a k -tuple \bar{x} , we will show that the equivalence class of \bar{x} is A -definable. Assume that E has n classes. By repeated use of Lemma 5.8 (vii), we can find a sequence $\{\bar{x}^i : i \leq n\}$ of tuples such that $\bar{x} \equiv_A \bar{x}^i$, and $\bar{x}^i \downarrow_A \bar{x}^0 \dots \bar{x}^{i-1} \bar{x}$. By the pigeonhole principle, there exist $0 \leq i < j \leq n$ such that \bar{x}^i and \bar{x}^j are in the same class C of E . It follows that C is definable over $A\bar{x}^i$ and over $A\bar{x}^j$, hence by Lemma A.5, it is definable over $\langle A\bar{x}^i \rangle \cap \langle A\bar{x}^j \rangle = \langle A \rangle$, i.e., over A . But then $\bar{x} \equiv_A \bar{x}^i$ implies that $\bar{x} \in C$. □

It may come as anticlimactic that we will state the result we are most interested in, viz. weak e.i. for EC_L , without proof: while the author has figured out a long and cumbersome argument, it was independently shown in an easier way by Kruckman and Ramsey [6, §3.4], and we invite the interested reader to consult their paper.

Theorem A.7 *For any language L , EC_L has weak elimination of imaginaries.* □

We end this section by stating explicitly the consequence of Theorem A.7 for definable functions of EC_L^{eq} that we will need in the proof of Theorem 4.5. While it is somewhat hairy to formulate, it follows by a simple compactness argument.

Lemma A.8 *Let $T \supseteq EC_L$. If*

$$T \vdash \alpha(\bar{x}) \rightarrow \exists^{\equiv m} \bar{u} \beta(\bar{u}, \bar{x}),$$

there are formulas $\{\alpha_i(\bar{x}) : i < n\}$ for some $n > 0$, and for each $i < n$, tuples of terms $\bar{t}^{i,j}(\bar{x})$, $j < m$, such that T proves that $\{\alpha_i : i < n\}$ define a partition of α (i.e., $\alpha(\bar{x}) \leftrightarrow \bigvee_{i < n} \alpha_i(\bar{x})$, and $\alpha_i(\bar{x}) \rightarrow \neg \alpha_{i'}(\bar{x})$ for $i \neq i'$), and

$$\alpha_i(\bar{x}) \rightarrow \left(\beta(\bar{u}, \bar{x}) \leftrightarrow \bigvee_{j < m} \bar{u} = \bar{t}^{i,j}(\bar{x}) \right).$$

Moreover, T proves $\alpha_i(\bar{x}) \rightarrow \bar{t}^{i,j}(\bar{x}) \neq \bar{t}^{i',j'}(\bar{x})$ for $j \neq j'$.

Proof: Note that $\alpha_i(\bar{x}) \rightarrow \bar{t}^{i,j}(\bar{x}) \neq \bar{t}^{i,j'}(\bar{x})$ for $j \neq j'$ follows from the rest, specifically $\alpha(\bar{x})$ implies that there are *at least* m tuples satisfying $\beta(\bar{u}, \bar{x})$.

We will prove the statement by induction on m . If $m = 0$, there is nothing to prove. Assuming the result holds for m , we will prove it for $m + 1$.

By Lemma A.2 and a compactness argument, there are tuples of terms $\bar{t}^{i,m}(\bar{x})$, $i < n$, such that T proves

$$\alpha(\bar{x}) \rightarrow \bigvee_{i < n} \beta(\bar{t}^{i,m}(\bar{x}), \bar{x}).$$

Putting

$$\alpha_i(\bar{x}) \leftrightarrow \beta(\bar{t}^{i,m}(\bar{x}), \bar{x}) \wedge \bigwedge_{i' < i} \neg \beta(\bar{t}^{i',m}(\bar{x}), \bar{x}),$$

we have a partition of α into formulas α_i such that T proves

$$\alpha_i(\bar{x}) \rightarrow \beta(\bar{t}^i(\bar{x}), \bar{x}).$$

Let us write

$$\beta_i(\bar{u}, \bar{x}) \leftrightarrow \beta(\bar{u}, \bar{x}) \wedge \bar{u} \neq \bar{t}^i(\bar{x}).$$

Then T proves

$$\alpha_i(\bar{x}) \rightarrow \exists^{=m} \bar{u} \beta_i(\bar{u}, \bar{x}).$$

Using the induction hypothesis, we can further refine the partition so that there are tuples of terms $\bar{t}^{i,j}(\bar{x})$, $j < m$, such that

$$T \vdash \alpha_i(\bar{x}) \rightarrow \left(\beta_i(\bar{u}, \bar{x}) \leftrightarrow \bigvee_{j < m} \bar{u} = \bar{t}^{i,j}(\bar{x}) \right).$$

Then T proves

$$\alpha_i(\bar{x}) \rightarrow \left(\beta(\bar{u}, \bar{x}) \leftrightarrow \bigvee_{j < m+1} \bar{u} = \bar{t}^{i,j}(\bar{x}) \right)$$

as required. □

Proposition A.9 *Let $T \supseteq EC_L$, and E be an equivalence relation on k -tuples definable in T . Then there are*

- integers $n > 0$, $r > 0$, and $m_i \geq 0$ for $i < n$,
- formulas $\varphi_i(\bar{x})$ for $i < n$, and
- terms $t_l^{i,j}(\bar{x})$ for $i < n$, $j < m_i$, $l < r$,

such that T proves

$$E(\bar{x}, \bar{y}) \leftrightarrow \bigvee_{i < n} \left(\varphi_i(\bar{x}) \wedge \varphi_i(\bar{y}) \wedge \{ \bar{t}^{i,j}(\bar{x}) : j < m_i \} = \{ \bar{t}^{i,j}(\bar{y}) : j < m_i \} \right), \quad (13)$$

where $\bar{t}^{i,j}(\bar{x})$ denotes the tuple $\langle t_l^{i,j}(\bar{x}) : l < r \rangle$, and the following condition hold:

- The formulas φ_i form a partition, i.e., T proves $\bigvee_{i < n} \varphi_i(\bar{x})$, and $\varphi_i(\bar{x}) \rightarrow \neg \varphi_{i'}(\bar{x})$ for $i \neq i'$.
- On each part φ_i , the tuples $\bar{t}^{i,j}$ are pairwise distinct, i.e.,

$$T \vdash \varphi_i(\bar{x}) \rightarrow \bigvee_{l < r} t_l^{i,j}(\bar{x}) \neq t_l^{i,j'}(\bar{x})$$

for each $i < n$, and $j < j' < m_i$.

Proof: By Theorem A.7, for each $M \models T$ and an equivalence class e of E , there is a real tuple $\bar{a} \in M$ such that $\bar{a} \in \text{acl}_{M^{\text{eq}}}(e)$, and $e \in \text{dcl}_{M^{\text{eq}}}(\bar{a})$. Thus, for each such M , e , and \bar{a} , we can find a formula $\psi(\bar{u}, \bar{x})$ and an integer $m \geq 1$ such that:

- $\psi(\bar{a}, \bar{x})$ for all $\bar{x} \in e$.
- $\psi(\bar{u}, \bar{x}) \rightarrow (\psi(\bar{u}, \bar{y}) \leftrightarrow E(\bar{x}, \bar{y}))$; that is, for a given \bar{u} , the set $\{\bar{x} : M \models \psi(\bar{u}, \bar{x})\}$ is either empty, or an equivalence class of E .
- For a given \bar{x} , there are either none or exactly m tuples \bar{u} such that $\psi(\bar{u}, \bar{x})$.

Using a compactness argument, there is a finite partition coarser than E definable in T by formulas $\{\varphi_i(\bar{x}) : i < n\}$, and for each $i < n$, there is a formula $\psi_i(\bar{u}, \bar{x})$, and an integer $m_i \geq 1$, such that T proves:

$$\begin{aligned} \psi_i(\bar{u}, \bar{x}) &\rightarrow (\psi_i(\bar{u}, \bar{y}) \leftrightarrow E(\bar{x}, \bar{y})), \\ \psi_i(\bar{u}, \bar{x}) &\rightarrow \varphi_i(\bar{x}), \\ \varphi_i(\bar{x}) &\rightarrow \exists^{=m_i} \bar{u} \psi_i(\bar{u}, \bar{x}). \end{aligned}$$

For a given i , if the tuple \bar{u} in $\psi_i(\bar{x}, \bar{u})$ has length 0, then $\varphi_i(\bar{x})$ is equivalent to $\psi_i(\bar{x})$, and it defines a single equivalence class of E ; in this case, we formally replace ψ_i with an always false formula using a dummy variable u , and put $m_i = 0$. In this way, we may ensure that all the \bar{u} are nonempty; by repeating one of its elements, we may in fact assume that they all have the same length $l > 0$ independent of i .

Using Lemma A.8, after possibly refining the partition $\{\varphi_i : i < n\}$, we can find tuples of terms $\bar{t}^{i,j}(\bar{x})$ such that T proves

$$\psi_i(\bar{u}, \bar{x}) \leftrightarrow \varphi_i(\bar{x}) \wedge \bigvee_{j < m_i} \bar{u} = \bar{t}^{i,j}(\bar{x}).$$

This implies (13), that is,

$$\varphi_i(\bar{x}) \wedge \varphi_i(\bar{y}) \rightarrow \left(E(\bar{x}, \bar{y}) \leftrightarrow \{\bar{t}^{i,j}(\bar{x}) : j < m_i\} = \{\bar{t}^{i,j}(\bar{y}) : j < m_i\} \right).$$

Indeed, $\{\bar{t}^{i,j}(\bar{x}) : j < m_i\} = \{\bar{u} : \psi_i(\bar{u}, \bar{x})\}$, and similarly for \bar{y} . The properties of ψ_i ensure that if $E(\bar{x}, \bar{y})$, these sets are equal, whereas if not (which can only happen if $m_i > 0$), they are disjoint, hence distinct. \square

B Dependence on language

The results in Section 5 give fairly tight model-theoretic classification of EC_L (NSOP₁, but TP₂) in the case that L includes at least one at least binary function symbol. However, the theories behave in different ways for other languages L ; in this section, we summarize the main model-theoretic properties of the theories EC_L in dependence on L .

Theorem B.1 *Let L be a language. Then any complete extension of EC_L is*

- (i) *strongly minimal iff L consists of nullary symbols;*
- (ii) *totally transcendental iff L consists of nullary symbols, and either finitely many unary relations, or one unary function;*
- (iii) *superstable iff L consists of at most unary symbols, at most one of which is a unary function;*

- (iv) *stable iff it has NIP iff L consists of at most unary symbols;*
- (v) *supersimple iff L consists of relations, constants, and at most one unary function;*
- (vi) *simple iff it has NTP_2 iff L consists of relations and at most unary functions;*
- (vii) NSOP_1 .

Proof: We may assume L contains no nullary relations, as these are fixed to true or false in any complete extension.

First, we establish the right-to-left implications. (i) is obvious. (ii): We may assume L is countable, we will verify EC_L is ω -stable. Let $M \models EC_L$, and $A \subseteq M$ be a set of parameters, which we may assume to be a submodel. If L consists of constants and finitely many unary relations $P_i(x)$, a type $\text{tp}(b/A)$ of an element $b \notin A$ is determined by $\{i : P_i(b)\}$, hence there are only finitely many (plus $|A|$ trivial types for $b \in A$). If L consists of constants and a unary function $F(x)$, $\text{tp}(b/A)$ is determined by the least n such that $F^n(b) \in A$ and the value of $F^n(b)$, or in case there is no such n , by the shape of the chain $\{F^n(b) : n < \omega\}$. This makes $|A| + \aleph_0$ possibilities.

(iii)²: Assume L consists of constants, unary relations $\{P_i(x) : i \in I\}$, and one unary function $F(x)$. Let $A \subseteq M \models EC_L$ be a submodel, and $b \in M$. Similarly to the ω -stable case, $\text{tp}(b/A)$ is determined by the type of b over A in the relation-free reduct of M , for which there are $|A| + \aleph_0$ possibilities, and by $\{(i, n) : i \in I, n < \omega, P_i(F^n(b))\}$. Thus, $|S_1(A)| \leq |A| + 2^{\|L\|}$.

(iv): Let L be unary, and M, A, b as before. Atomic formulas involved in $\text{tp}(b/A)$ are of the forms $t(b) = s(b)$ or $P(t(b))$ (i.e., not referring to A , hence at most $\|L\|$ many), or $t(b) = a$ for $a \in A$. For any $t(x)$, the type either contains one formula of the form $t(b) = a$, or all of $\{t(b) \neq a : a \in A\}$. Thus, $|S_1(A)| \leq |A|^{\|L\|}$; in particular, EC_L is κ -stable whenever $\kappa = \kappa^{\|L\|}$.

(vi): The Kim–Pillay theorem [5] (see also [13, 2.6.1]) states that a theory is simple if we can define an independence relation $A \downarrow_C B$ that satisfies properties (i), (ii), (iv), [a weaker form of] (vi), (vii), (viii), and (ix) from Lemma 5.8, and the converse implication to (iv); the latter amounts to *base monotonicity*: if $C' \subseteq B$, then $A \downarrow_C B$ implies $A \downarrow_{CC'} B$.

Now, if L contains no functions of arity 2 or more, the relation $A \downarrow_C B$ from Definition 5.7 can be restated as

$$A \downarrow_C B \iff \forall a \in A \forall b \in B \langle a \rangle \cap \langle b \rangle \subseteq \langle C \rangle.$$

This shows it satisfies base monotonicity, even in the stronger form that $A \downarrow_C B$ implies $A \downarrow_{CC'} B$ for arbitrary C' .

(v): In terms of the independence relation, a simple theory is supersimple iff it satisfies a strong form of local character: for every B and finite A , there is a *finite* $B' \subseteq B$ such that $A \downarrow_{B'} B$ (cf. [13]). If L consists of relations and constants, we can take $B' = A \cap B$. If L also contains one unary function F , we construct a set $B' \subseteq B$ with $|B'| \leq |A|$ such that for each $u \in A$, if $F^n(u) \in \langle B \rangle \setminus \langle \emptyset \rangle$ for some n , then the least such n satisfies $F^n(u) = F^m(v)$ for some $v \in B'$ and $m \in \omega$.

(vii) is Corollary 5.10.

Now we turn to the left-to-right implications.

(vi) follows from Proposition 5.6.

(v): We may assume T is simple, i.e., all functions in L are at most unary. If there are two unary functions $F(x), G(x)$, a monster model $\mathbb{M} \models T$ will contain an a such that $t(a) \neq s(a)$ for any pair of distinct unary terms t, s . Let $B = \{G(F^n(a)) : n < \omega\}$. Then any finite $C \subseteq B$ satisfies $\langle C \rangle \cap B = C \subsetneq B \subseteq \langle a \rangle$, hence $a \not\downarrow_C B$. Thus T is not supersimple.

²A theory is superstable iff it is stable and supersimple, hence (iii) follows from (iv) and (v). However, we prefer to give a direct proof not relying on sophisticated tools like the Kim–Pillay theorem.

(iv): If L contains an at least binary function, it is even TP_2 . If L contains an at least binary relation, wlog $P(x, y)$, then the formula $P(x, y)$ has IP, as every finite model embeds into any model of EC_L .

(iii): We may assume L is unary, lest the theory is not even stable. So, assume L contains two unary functions $F(x), G(x)$. Let κ be an arbitrarily large cardinal such that $\kappa^\omega > \kappa$, $M \models EC_L$, and $A = \{a_\alpha : \alpha < \kappa\} \subseteq M$ of cardinality κ . For every $\sigma : \omega \rightarrow \kappa$, the type

$$p_\sigma(x) = \{G(F^n(x)) = a_{\sigma(n)} : n \in \omega\}$$

over A is consistent, and p_σ and p_τ are incompatible for $\sigma \neq \tau$. Thus, $|S_1(A)| \geq \kappa^\omega$, and T is not κ -stable.

(ii): We may assume T is superstable. If L contains a function $F(x)$, and a relation $P(x)$, the 2^ω types

$$p_I(x) = \{P(F^n(x)) : n \in I\} \cup \{\neg P(F^n(x)) : n \notin I\}$$

for $I \subseteq \omega$ witness that the $\langle F, P \rangle$ -fragment of T is not ω -stable, hence T is not totally transcendental. If L contains infinitely many predicates $P_n(x)$, $n < \omega$, we can likewise use the types $\{P_n(x) : n \in I\} \cup \{\neg P_n(x) : n \notin I\}$.

(i): If L contains a non-nullary (wlog unary) predicate $P(x)$, then both $P(M)$ and its complement are infinite for any $M \models EC_L$, thus M is not minimal. Likewise, if L contains a function $F(x)$, the formula $F(x) = x$ defines an infinite set with infinite complement. \square

Apart from tameness properties from classification theory, we also discuss some more elementary invariants of the theories, namely the number of types, and the number of complete extensions. We only consider countable languages for the rest of this section.

Recall that a countable complete theory is called *small* if it has countable many complete n -types for all $n \in \omega$; this holds if and only if it has a saturated countable model.

Proposition B.2 *Let L be countable, and T a complete extension of EC_L .*

- (i) *Let L consist of relations and constants, where the number of non-nullary relations and T -unequal constants is finite. Then T is ω -categorical.*
- (ii) *Let L consist either of nullary symbols and one unary function, or of nullary relations, finitely many unary relations, and infinitely many T -unequal constants. Then T has \aleph_0 complete n -types for each $0 < n < \omega$, hence it is not ω -categorical, but it is small.*
- (iii) *Otherwise T has 2^ω complete 1-types.*

Proof:

(i): By quantifier elimination, there are only finitely many formulas in n variables for every n (ignoring sentences).

(ii): On the one hand, if F is a unary function, there are infinitely many incompatible 1-types of the form

$$F^n(x) = F^{n+1}(x) \wedge \bigwedge_{i < j \leq n} F^i(x) \neq F^j(x).$$

If $\{c_i : i < \omega\}$ are provably pairwise distinct constants, then we have infinitely many 1-types extending $x = c_i$.

On the other hand, T is ω -stable by Theorem B.1.

(iii): If we have a unary function $F(x)$, and a nonnullary relation (wlog unary) $R(x)$, we have countably many independent atomic formulas $R(F^n(x))$, thus 2^ω 1-types. Likewise, if there is another unary function $G(x)$, the formulas $G(F^n(x)) = x$ are independent, and if we have

a binary (or more) function $H(x, y)$, we can consider the formulas $H(s_1(x), s_n(x)) = x$, where $s_1(x) := H(x, x)$, and $s_{n+1}(x) := H(x, s_n(x))$. If there are infinitely many proper predicates, or an at least binary predicate and infinitely many distinct constants, we are also done. \square

Corollary B.3 *Let L be countable.*

- (i) EC_L is complete iff L contains no nullary symbols, or consists of one constant.
- (ii) If L contains no constants, and its nullary relations are finitely many, then EC_L has finitely many complete extensions.
- (iii) If L is finite, and contains no nonconstant functions, then EC_L has finitely many complete extensions.
- (iv) If L consists of a unary function, and finitely many nullary symbols, at least one of which is a constant, then EC_L has countably infinitely many complete extensions.
- (v) Otherwise EC_L has 2^ω complete extensions.

Proof: (i): It is easy to see that in the other cases, there is at least one nontrivial atomic sentence.

(ii)–(v): If there are infinitely many nullary relations or constants, there are infinitely many independent atomic sentences, hence 2^ω complete extensions.

Assume there are only finitely many nullary symbols, and let L_0 be L minus constants. The only atomic sentences in L_0 are nullary relations, hence EC_{L_0} has finitely many complete extensions. If L has $k > 0$ constants, then completions of EC_L correspond to complete k -types over completions of EC_{L_0} . By Proposition B.2, these are finitely many if L_0 is finite and contains no functions, \aleph_0 if L_0 consists of one unary function and nullary predicates, and 2^ω otherwise. \square

Let us touch upon a somewhat different topic now. In Section 3, we proved the existence of the model completion EC_L in a laborious way by, essentially, computing an explicit finite axiomatization of resultants. Model theorists are not very keen on getting their hands dirty with actual formulas, and prefer higher-level methods; in particular, a very popular technique for construction of model completions is using *Fraïssé limits*. For example, if L is a finite relational language, one can show the existence of EC_L quite easily by taking the theory of the Fraïssé limit of the class of all finite L -structures. The reader may wonder why we did not use this method as well, thus we will have a look at what we can achieve with Fraïssé limits in our situation. Let us first recall the basic setup.

Definition B.4 The *age* of a structure M is the class of all finitely generated structures embeddable in M .

A *Fraïssé class* is a class \mathcal{K} of finitely generated countable structures satisfying the following conditions:

- \mathcal{K} contains only countably many structures up to isomorphism.
- Hereditary property (HP): if a finitely generated structure B embeds in $A \in \mathcal{K}$, then $B \in \mathcal{K}$.
- Joint embedding property (JEP): for any finite set $\{B_0, \dots, B_{n-1}\} \subseteq \mathcal{K}$, there exists $A \in \mathcal{K}$ such that each B_i embeds in A .
- Amalgamation property (AP): for any $C, B_0, B_1 \in \mathcal{K}$ and embeddings $g_i: C \rightarrow B_i$ ($i = 0, 1$), there exists $A \in \mathcal{K}$ and embeddings $f_i: B_i \rightarrow A$ ($i = 0, 1$) such that $f_0 \circ g_0 = f_1 \circ g_1$.

Note that JEP is equivalent to its special cases $n = 0$ and $n = 2$. The former amounts to $\mathcal{K} \neq \emptyset$; the latter is, under the assumption of AP and HP, equivalent to the simpler property that \mathcal{K} contains only one 0-generated structure up to isomorphism.

Note also that if the language is countable, finitely generated structures are automatically countable.

A structure M is *ultrahomogeneous* if for every finitely generated substructures $A, B \subseteq M$ and every isomorphism $f: A \simeq B$, there exists an automorphism g of M such that $f \subseteq g$.

Proposition B.5 *A class \mathcal{K} is a Fraïssé class if and only if it is the age of a countable ultrahomogeneous structure M . In that case, M is unique up to an isomorphism; it is called the Fraïssé limit of \mathcal{K} .* \square

Model completions can then be conveniently constructed using Fraïssé limits as follows.

Proposition B.6 *Let T be a universal theory in a finite language consisting of relations and constants. If the class of finite models of T has AP and JEP, then it is a Fraïssé class; its Fraïssé limit M is a unique countable e.c. model of T up to isomorphism. The theory of M is the model completion of T .* \square

This works well for EC_L when L is a finite language with relations and (by considering separately each quantifier-free diagram) constants. However, it is not applicable if L contains proper functions; this cannot be circumvented by somehow encoding the structures in a relational language, as the theories have fundamentally different properties: in particular, model completions constructed by Proposition B.6 are always ω -categorical, and this seems to be inherent in the method.

Thus, it seems Fraïssé limits are not helpful for showing the existence of EC_L in general. Nevertheless, we may still wonder if limits of suitable Fraïssé classes could provide some interesting models of EC_L .

Notice that in order to have JEP, whatever class of models we consider must satisfy the same quantifier-free sentences; in view of quantifier elimination of EC_L , this amounts to choosing a completion of EC_L . So, let L be a countable language, and T a complete extension of EC_L . We can write $T = EC_L + \text{Diag}(M_0)$, where M_0 is a (possibly empty) structure whose every element is the value of a closed term, i.e., M_0 is 0-generated; we denote $T_0 = \text{Diag}(M_0)$. There are two candidate Fraïssé classes of models of T_0 that immediately spring to mind:

- The class \mathcal{K}_{fg} of finitely generated models of T_0 .
- The class \mathcal{K}_{fin} of finite models of T_0 .

It is easy to verify that \mathcal{K}_{fg} satisfies HP, AP, and JEP. Thus, it is a Fraïssé class iff it contains countably many structures up to isomorphism, which happens iff T is *small*. (We already characterized when T is small in Proposition B.2.) We leave it to the reader to check that in this case, the Fraïssé limit of \mathcal{K}_{fg} is the countable saturated model of T .

Clearly, \mathcal{K}_{fin} is nonempty only if M_0 is finite, hence we will assume this for the moment. Again, it is easy to see \mathcal{K}_{fin} has HP, AP, and JEP, hence it is a Fraïssé class iff it is countable up to isomorphism. This holds iff L contains only finitely many nonnullary symbols; since the finiteness of M_0 implies there are only finitely many constants up to equality in T_0 , and nullary relations are in T_0 fixed to true or false, we can as well assume without loss of generality that L is finite. The Fraïssé limit of \mathcal{K}_{fin} is then described by the following result.

Proposition B.7 *Let $T = EC_L + \text{Diag}(M_0)$, where M_0 is a 0-generated L -structure, and L and M_0 are finite.*

Then T has a prime (equivalently: countable atomic) model, which can be characterized as the unique locally finite (i.e., such that finitely generated submodels are finite) countable model of T , and it can be constructed as the Fraïssé limit of the class of all finite L -structures that extend M_0 .

Proof: By the preceding discussion, \mathcal{K}_{fin} is a Fraïssé class, hence it has a Fraïssé limit M . Clearly, $M \supseteq M_0$ is countable, and locally finite. By general properties of Fraïssé limits (see e.g. [3]), M is existentially closed in the class of locally finite structures. This in fact implies that M is e.c. in the class of all structures, hence $M \models T$: if $\varphi(\bar{u})$ is an \exists_1 formula, $\bar{u} \in M$, and $M \subseteq N \models \varphi$, let

$$N' = M \cup \{t^N(\bar{u}) : t \text{ is a subterm of } \varphi\},$$

and make it an L -structure by preserving the values of all relations and functions in N where possible, and $f^{N'}(\bar{v}) = a$ for some fixed $a \in M$ if $f^N(\bar{v}) \notin N'$. Then $M \subseteq N' \models \varphi(\bar{u})$, and every finitely generated submodel of N' is included in $N' \setminus M$ plus a finitely generated submodel of M , hence N' is locally finite. Thus, $M \models \varphi(\bar{u})$.

Every locally finite $M' \models T$ is atomic: let $\bar{a} \in M'$, and A be the submodel of M' generated by \bar{a} . Write $A \setminus \{\bar{a}\} = \{b_i : i < m\}$. Then $\text{tp}(\bar{a})$ is generated by $\exists y_0, \dots, y_{m-1} \text{Diag}(A)$, where $\text{Diag}(A)$ is written using variables \bar{x}, \bar{y} in place of \bar{a}, \bar{b} .

Thus, M is a countable atomic, hence prime, model of T , and by uniqueness of prime models, every countable locally finite model of T is isomorphic to M . \square

There are other cases when T has a prime model: for example, if T is small (see Proposition B.2). In fact, we can give a full description (for countable languages). Recall that a complete countable theory has a prime model if and only if it is atomic.

Proposition B.8 *Let $T = EC_L + \text{Diag}(M_0)$, where L is countable, and M_0 is a 0-generated L -structure. Then T has a prime model iff it falls in of the following cases:*

- (i) $M_0 \models EC_L$.
- (ii) L contains only finitely many nonnullary symbols, all of which are unary.
- (iii) M_0 is finite, and L contains only finitely many nonnullary symbols.

Proof: Clearly, (i) implies that M_0 is a prime model of T . We constructed a prime model in case (iii) in Proposition B.7.

Assume (ii) holds; we will show T is atomic. Let $\varphi(\bar{x})$ be a T -consistent formula, which we may assume to be quantifier-free. By the argument in the proof of Proposition B.7, φ is satisfiable in a model $M \supseteq M_0$ such that $M \setminus M_0$ is finite. Let $M_0 \subseteq M \models \varphi(\bar{a})$ be such that $|M \setminus M_0|$ is minimal possible. Using minimality, we can choose for each $b \in M \setminus M_0$ a term $t_b(\bar{x})$ (in fact, a subterm of φ) such that $t_b(\bar{a}) = b$. Let C be the (finite) set of all elements of M_0 that are values of subterms of $\varphi(\bar{a})$. For each element $b \in M_0$, let us fix a constant term t_b whose value is b . Let $\psi(\bar{x})$ be the conjunction of the following formulas:

- $t_u(\bar{x}) \neq t_v(\bar{x})$ for each $u \in M \setminus M_0$, and $u \neq v \in (M \setminus M_0) \cup C$;
- $x_i = t_{a_i}(\bar{x})$ for each i ;
- $F(t_u(\bar{x})) = t_{F(u)}(\bar{x})$ for each $u \in M \setminus M_0$, and $F \in L$ a nonconstant function symbol;
- $R(t_u(\bar{x}))$ or $\neg R(t_u(\bar{x}))$ (whichever is satisfied by \bar{a}) for each $u \in M \setminus M_0$, and $R \in L$ a nonnullary relation symbol.

By construction, $M \models \psi(\bar{a})$, hence ψ is consistent. By induction on the length of s , we see that $T_0 \vdash \psi(\bar{x}) \rightarrow s(\bar{x}) = t_{s(\bar{a})}(\bar{x})$ for each subterm s of φ ; it follows easily that $T_0 \vdash \psi(\bar{x}) \rightarrow \varphi(\bar{x})$. We claim that ψ is an atom; by quantifier elimination, it suffices to show that it implies the quantifier-free type of \bar{a} . Thus, let us consider a model $M' \models \psi(\bar{a}')$. We may assume M' is generated by \bar{a}' . The conjuncts of ψ ensure that the mapping $b \mapsto t_b(\bar{a}')$ is a homomorphism $f: M \rightarrow M'$ such that $f(a_i) = a'_i$. It is the identity on M_0 , and injective on $M \setminus M_0$. Since ψ implies φ , we have $M' \models \varphi(\bar{a}')$. By the minimality of M , this implies $|M' \setminus M_0| \geq |M \setminus M_0|$, hence f must map $M \setminus M_0$ to $M' \setminus M_0$. Thus, f is in fact an isomorphism of M to M' .

On the other hand, assume that none of (i)–(iii) holds. Since M_0 does not validate some axiom of EC_L as in Definition 3.6, and all elements of M_0 can be denoted by constant terms, there exists a consistent quantifier-free formula in one variable $\theta(x)$ which is not satisfiable in M_0 . Assuming for contradiction that T is atomic, we may choose $\theta(x)$ to be an atom.

If L includes infinitely many nonnullary relation symbols, let $R(\bar{x})$ be one that does not appear in θ . Then $\theta(x) \wedge R(x, \dots, x)$ and $\theta(x) \wedge \neg R(x, \dots, x)$ are both consistent, contradicting θ being an atom: taking an arbitrary model $M \models \theta(a)$, we may flip the value of $R(a, \dots, a)$ without affecting $M \models \theta(a)$. By a similar argument, we obtain a contradiction if L contains infinitely many nonconstant function symbols.

The remaining case is that M_0 is infinite, and L includes an at least binary symbol, say, a relation symbol $R(\bar{x}, y)$ (the case of a function symbol is similar). Fix a model $M \models \theta(a)$. Since M_0 is infinite, we can find $c \in M_0$, denoted by a closed term t , such that c is not the value of any subterm of $\theta(a)$. Then we can flip the value of $R(a, \dots, a, c)$ without affecting $M \models \theta(a)$, hence the formulas $\theta(x) \wedge R(x, \dots, x, t)$ and $\theta(x) \wedge \neg R(x, \dots, x, t)$ are both consistent, a contradiction. \square

C More on representation

For completeness, let us present a few counterexamples to possible strengthenings of some of the basic claims in Section 2.2.

First, we mentioned that it is enough to represent a specific dprp, or prf, in a theory in order to show its essential undecidability. In contrast, we will prove that any finite set of trf and rp can be represented in a decidable theory. More generally, it holds even for infinite families of such functions and predicates as long as they are *uniformly recursive*: here, we call a sequence $\{F_n : n \in \omega\}$ of recursive functions $F_n : \mathbb{N}^{k_n} \rightarrow \mathbb{N}$ uniformly recursive if the functions $n \mapsto k_n$ and $\langle n, w \rangle \mapsto F_n((w)_0, \dots, (w)_{k_n-1})$ are recursive, and similarly for sequences of predicates.

Proposition C.1 *Let \mathcal{R} be a uniformly recursive sequence of trf and rp. Then there exists a consistent decidable theory T and a recursive numeral sequence σ such that \mathcal{R} is representable in T w.r.t. σ .*

Proof: Let L be the (recursive) language $L_{\mathcal{R}}$ from Definition 2.4, and $\langle \mathbb{N}, \mathcal{R} \rangle$ be the “standard L -model” with domain \mathbb{N} and the L -symbols realized by the corresponding elements of \mathcal{R} . Then $EC_L + \text{Diag}(\langle \mathbb{N}, \mathcal{R} \rangle) = EC_L + REP_{\mathcal{R}}$ is a decidable complete theory by Corollary 3.8, and it represents \mathcal{R} w.r.t. the sequence $n \mapsto \underline{n}$. \square

The second example serves two-fold purpose. For one, it exhibits that a recursively axiomatizable (or even decidable) theory may represent a non-recursive predicate w.r.t. a non-recursive sequence of numerals. Second, it shows that representation of rp, or even dprp, does not imply essential undecidability if the sequence of numerals is not recursive. We will first prove a simple version applying to finite languages.

Proposition C.2 *Let \mathcal{P} be a finite set of predicates and disjoint pairs. Then there exists a consistent decidable theory T that represents \mathcal{P} (w.r.t. a possibly non-recursive sequence of numerals).*

Proof: Since we do not require the elements of \mathcal{P} to be recursive to begin with, we may as well extend each disjoint pair to a predicate, thus we will assume \mathcal{P} consists of predicates without loss of generality. Let L be the finite relational language corresponding to \mathcal{P} . Since EC_L is decidable, it has a recursive model M (using the standard Henkin completion procedure; in fact, in this case, it is not difficult to construct the model explicitly). Let $T = EC_L + \text{Diag}(M)$, which is a decidable complete theory. Since EC_L is ω -categorical, M is (a recursive presentation of) its unique countable model, and the countable structure $\langle \mathbb{N}, \mathcal{P} \rangle$ embeds in M ; let us fix such an embedding $\sigma: n \mapsto \underline{n}$, where the elements $\underline{n} \in M$ are identified with the corresponding constants in the language of T . Then all \mathcal{P} -predicates are represented in T w.r.t. σ . \square

We could handle countable sets of predicates and disjoint pairs of *bounded arity* with a bit of preprocessing, but we will need more work to take care of the general case: in particular, note that EC_L is no longer ω -categorical (or even small) if L is an infinite relational language; we will use a slightly different theory instead.

Let L_0 be the language that includes an n -ary relation $R_n(x_0, \dots, x_{n-1})$ for every $n \geq 1$, and T_0 be the universal L_0 -theory axiomatized by

$$R_n(\bar{x}) \rightarrow \bigwedge_{i < j < n} x_i \neq x_j$$

for each n . The class \mathcal{K} of finite models of T_0 is easily seen to have HP, AP, and JEP. Crucially, it contains only countably many nonisomorphic structures: in fact, for any n , there are only finitely many T_0 -structures of size n , as the axioms force all but the first n L_0 -relations to be empty.

Thus, \mathcal{K} is a Fraïssé class, and it has a Fraïssé limit M . Since any isomorphism between finite submodels of M extends to an automorphism of M , and there are finitely many isomorphism types of such submodels of fixed size, it follows that the theory T^* of M is ω -categorical, and has elimination of quantifiers. Since all finite(ly generated) models of T_0 embed in M , T^* is in fact the model completion of T_0 .

It is not difficult to explicitly axiomatize T^* by suitable extension axioms, thus T^* is recursively axiomatizable, and in fact, decidable (as it is complete). Moreover, M may be presented as a recursive structure. Thus, its elementary diagram $T_M^* = T^* + \text{Diag}(M)$ is a decidable theory.

We will now show that *every* predicate and disjoint pair (with no recursivity assumption) is representable in T_M^* , using various (mostly nonrecursive) sequences of numerals. In fact, any countable set of predicates and disjoint pairs can be represented w.r.t. the *same* sequence of numerals:

Proposition C.3 *Every countable set \mathcal{P} of predicates and disjoint pairs is representable in the decidable theory T_M^* .*

Proof: As in the proof of Proposition C.2, we will assume \mathcal{P} to consist of predicates. Without loss of generality, we may also assume that \mathcal{P} is closed under identification of variables: i.e., if $P \in \mathcal{P}$ is n -ary, and $i < j < n$, the $(n-1)$ -ary predicate $P_{i,j}$ defined by

$$P_{i,j}(x_0, \dots, x_{n-2}) \iff P(x_0, \dots, x_{j-1}, x_i, x_j, \dots, x_{n-2})$$

is in \mathcal{P} . Let us also define

$$P'(x_0, \dots, x_{n-1}) \iff P(x_0, \dots, x_{n-1}) \wedge \bigwedge_{i < j < n} x_i \neq x_j.$$

Then we can reconstruct P from P' and predicates of smaller arity as

$$P(x_0, \dots, x_{n-1}) \iff P'(x_0, \dots, x_{n-1}) \vee \bigvee_{i < j < n} (x_i = x_j \wedge P_{i,j}(x_0, \dots, x_{j-1}, x_{j+1}, \dots, x_{n-1})),$$

hence by induction on the arity, we see that all \mathcal{P} -predicates are quantifier-free definable from predicates from $\mathcal{P}' = \{P' : P \in \mathcal{P}\}$. Thus, without loss of generality, we may replace \mathcal{P} with \mathcal{P}' , i.e., we may assume that all predicates $P \in \mathcal{P}$ satisfy

$$P(\bar{x}) \rightarrow \bigwedge_{i < j} x_i \neq x_j.$$

We may also assume \mathcal{P} contains predicates of every arity $n \in \mathbb{N}$.

For each $n \in \mathbb{N}$, let us fix an enumeration $\{P_{n,m} : m \in \mathbb{N}\}$ of all n -ary predicates from \mathcal{P} such that each such predicate occurs more than n times in the enumeration. Finally, for each $n \geq 1$, we define

$$P_n(x_0, \dots, x_{n-1}) \iff P_{n-1,x_0}(x_1, \dots, x_{n-1}) \wedge \bigwedge_{i > 0} x_i \neq x_0.$$

The structure $\langle \mathbb{N}, P_n : n \geq 1 \rangle$ is a countable model of T_0 , hence it embeds in M . Let us fix such an embedding $\sigma : n \mapsto \underline{n}$, where we identify $\underline{n} \in M$ with the corresponding constant in the language of T_M^* . Thus, σ serves as a sequence of numerals, and each relation P_n is represented in T_M^* w.r.t. σ by the formula $R_n(\bar{x})$.

It follows that each $P \in \mathcal{P}$ is represented in T_M^* as well: if P is n -ary, let us fix distinct $m_0, \dots, m_n \in \mathbb{N}$ such that $P = P_{n,m_0} = \dots = P_{n,m_n}$. By the pigeonhole principle, all the numbers m_0, \dots, m_n cannot simultaneously appear in any tuple satisfying P ; thus, the formula

$$\bigvee_{j \leq n} R_{n+1}(\underline{m_j}, x_0, \dots, x_{n-1}).$$

represents $P(x_0, \dots, x_{n-1})$ in T_M^* w.r.t. σ . □

The main result of this paper shows that representability of prf does not imply interpretability of R . Another problem in a similar vein is to clarify the relationship between representability of different types of recursive objects. Specifically, let us consider the following conditions on a theory T :

- (i) The set RP is representable in T .
- (ii) The set DPRP is representable in T .
- (iii) The set $\text{RP} \cup \{\text{succ}\}$ is representable in T .
- (iv) The set $\text{DPRP} \cup \{\text{succ}\}$ is representable in T .
- (v) The set TRF is representable in T .
- (vi) The set PRF is representable in T .

We discuss separately the cases with the successor function included because of pathologies exhibited by representation of predicates and disjoint pairs w.r.t. potentially non-recursive sequences of numerals, as seen in Proposition C.3.

As we already mentioned in Section 2.2, it is easy to see that (vi) \rightarrow (v), (iv) \rightarrow (iii), (ii) \rightarrow (i), (vi) \rightarrow (iv) \rightarrow (ii), and (v) \rightarrow (iii) \rightarrow (i). We wish to show now that no other implications between these six conditions hold in general. This turns out not to be quite true—once again due to pathologies exhibited by non-recursive sequences of numerals. In the spirit of battering

reality vigorously until it complies with our preformed expectations, we fix this by considering a more strict notion of implication between representability of classes \mathcal{R}_0 and \mathcal{R}_1 , namely: does representability of \mathcal{R}_0 in T w.r.t. a sequence of numerals σ imply the representability of \mathcal{R}_1 w.r.t. σ ? This leads to the desired answer.

Proposition C.4

- (i) *There exists a theory that represents $\text{DPRP} \cup \{\text{succ}\}$, but does not represent TRF.*
- (ii) *There exists a theory that represents TRF, but does not represent $\text{DPRP} \cup \{\text{succ}\}$.*
- (iii) *There exists a theory that represents DPRP, but does not represent succ.*
- (iv) *Every theory that represents RP also represents arbitrary countable sets of predicates and disjoint pairs.*
- (v) *There exists a theory that represents TRF w.r.t. a sequence of numerals σ , but does not represent DPRP w.r.t. σ .*

Proof: If $\mathcal{R} \subseteq \text{PRF} \cup \text{DPRP}$, let $\text{ECREP}_{\mathcal{R}} = \text{EC}_{L_{\mathcal{R}}} + \text{REP}_{\mathcal{R}}$, where $L_{\mathcal{R}}$ is the language of $\text{REP}_{\mathcal{R}}$.

(i): Let $\mathcal{R} = \text{DPRP} + \{\text{succ}\}$. Then the theory $\text{ECREP}_{\mathcal{R}}$ represents \mathcal{R} , and it is supersimple (hence NTP_2) by Theorem B.1. In contrast, any theory representing trf is TP_2 by Proposition 5.6.

(ii): The (complete) theory $T = \text{ECREP}_{\text{TRF}}$ represents trf. Let $P = \langle P^+, P^- \rangle$ be a recursively inseparable dprp. We claim that T cannot represent $\{P, \text{succ}\}$, i.e., it does not interpret $\text{REP}_{P, \text{succ}}$: if it did, then $\text{REP}_{P, \text{succ}}$ would be also interpretable in a finite-language fragment T_0 of T , which would make T_0 an essentially undecidable theory. However, T_0 is of the form $\text{ECREP}_{\mathcal{R}}$ for a finite $\mathcal{R} \subseteq \text{TRF}$, thus it is decidable as in the proof of Proposition C.1.

(iii): The theory $T = \text{ECREP}_{\text{DPRP}}$ represents dprp. Assume for contradiction that it represents succ, i.e., it interprets REP_{succ} . Then REP_{succ} is interpretable in a finite-language fragment T_0 of T , which is an extension of EC_L for L a finite language with relations and constants. Thus, (any completion of) T_0 is ω -categorical. It follows that REP_{succ} also has an ω -categorical extension, but this is impossible, as it has infinitely many definable constants.

(iv): The argument from the proof of Proposition C.3 shows that every countable set \mathcal{P} of predicates and disjoint pairs is representable (w.r.t. a suitable sequence of numerals) in any theory that represents all predicates definable in the model M .

(v): The theory $T = \text{ECREP}_{\text{TRF}}$ represents trf w.r.t. the sequence of numerals \underline{n} . If it also represented DPRP w.r.t. the same numeral sequence, it would in fact represent $\text{DPRP} \cup \{\text{succ}\}$, which we already know to be impossible from (ii). \square