



INSTITUTE OF MATHEMATICS

THE CZECH ACADEMY OF SCIENCES

**A finite volume scheme for the Euler  
system inspired by the two velocities  
approach**

*Eduard Feireisl*

*Mária Lukáčová-Medvidová*

*Hana Mizerová*

Preprint No. 20-2018

PRAHA 2018



# A finite volume scheme for the Euler system inspired by the two velocities approach

Eduard Feireisl <sup>\*</sup>    Mária Lukáčová-Medviďová <sup>†</sup>    Hana Mizerová <sup>\*§</sup>

May 15, 2018

<sup>\*</sup> Institute of Mathematics of the Academy of Sciences of the Czech Republic  
Žitná 25, CZ-115 67 Praha 1, Czech Republic  
feireisl@math.cas.cz  
mizerova@math.cas.cz

<sup>†</sup> Institute of Mathematics, Johannes Gutenberg-University Mainz  
Staudingerweg 9, 55128 Mainz, Germany  
lukacova@uni-mainz.de

<sup>§</sup> Department of Mathematical Analysis and Numerical Mathematics  
Faculty of Mathematics, Physics and Informatics, Comenius University in Bratislava  
Mlynská dolina, 842 48 Bratislava, Slovakia

## Abstract

We propose a new finite volume scheme for the Euler system of gas dynamics motivated by the model proposed by H. Brenner. Numerical viscosity imposed through upwinding acts on the velocity field rather than on the convected quantities. The resulting numerical method enjoys the crucial properties of the Euler system, in particular positivity of the approximate density and pressure and the minimal entropy principle. In addition, the approximate solutions generate a dissipative measure-valued solutions of the limit system. In particular, the numerical solutions converge to the smooth solution of the system as long as the latter exists.

**Keywords:** complete Euler system, finite volume method, two velocities model, dissipative measure-valued solution

---

<sup>\*</sup>The research of E.F. and H.M. leading to these results has received funding from the Czech Sciences Foundation (GAČR), Grant Agreement 18-05974S. The Institute of Mathematics of the Academy of Sciences of the Czech Republic is supported by RVO:67985840.

<sup>†</sup>The reserach of M.L. was supported by the German Science Foundation under the Collaborative Research Centers TRR 146 and TRR 165.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Complete Euler system . . . . .	3
1.2	Brenner's model . . . . .	5
1.3	Finite volume scheme . . . . .	6
<b>2</b>	<b>Numerical scheme</b>	<b>6</b>
2.1	Preliminaries . . . . .	6
2.2	Approximation scheme . . . . .	8
<b>3</b>	<b>Entropy balance</b>	<b>10</b>
3.1	Renormalization . . . . .	11
3.1.1	Continuity equation . . . . .	11
3.1.2	Transport equation . . . . .	11
3.2	Discrete entropy balance equation . . . . .	12
3.2.1	Discrete kinetic energy equation . . . . .	12
3.2.2	Discrete internal energy equation . . . . .	13
3.2.3	Discrete entropy balance . . . . .	14
3.2.4	Entropy renormalization . . . . .	16
<b>4</b>	<b>Stability</b>	<b>18</b>
4.1	Mass and energy conservation . . . . .	18
4.2	Minimum entropy principle . . . . .	19
4.3	Positivity of the pressure, existence of the temperature . . . . .	19
4.4	Positivity of the density . . . . .	20
4.5	Existence of approximate solutions . . . . .	21
4.6	Entropy estimates . . . . .	21
<b>5</b>	<b>Consistency</b>	<b>23</b>
5.1	Numerical flux . . . . .	23
5.1.1	Upwinds . . . . .	23
5.1.2	$\mu_h$ -dependent terms . . . . .	28
5.2	The artificial viscosity and the pressure terms . . . . .	29
5.3	Consistency formulation . . . . .	30
<b>6</b>	<b>Convergence</b>	<b>32</b>
6.1	Young measure generated by the approximate solutions . . . . .	32
6.2	Kinetic energy concentration defect . . . . .	32
6.3	Limit problem . . . . .	33

# 1 Introduction

In 2005, H. Brenner [4] proposed a new approach to dynamics of *viscous and heat conducting* fluids based on two velocity fields distinguishing the bulk mass transport from the purely microscopic motion. Brenner’s approach has been subjected to thorough criticism by Öttinger et al. [21], where its incompatibility with certain physical principles is shown. Nevertheless, some computational simulations have been performed by Greenshields and Reese [18], Bardow and Öttinger [2], Guo and Xu [20] showing suitability of the model in specific situations. More recently, Guermond and Popov [19] rediscovered the model pointing out its striking similarity with certain numerical methods based on the finite volume approximation of the *inviscid* fluids. In particular, unlike the conventional and well accepted Navier–Stokes–Fourier system, Brenner’s model reflects the basic properties of the complete Euler system in the asymptotic limit of vanishing transport coefficients.

Inspired by these observations, we propose a new finite volume scheme for the complete Euler system based on Brenner’s ideas. In particular, the new scheme enjoys the following properties:

- **Positivity of the discrete density and temperature**

The approximate density and temperature remain strictly positive on any finite time interval.

- **Entropy stability**

The discrete entropy inequality in the sense of Tadmor is satisfied, see [25, 24].

- **Minimum entropy principle**

The entropy attains its minimum at the initial time, cf. [26, 19].

- **Weak BV estimates**

We control suitable weak BV norms of the discrete density, temperature and velocity.

In comparison with the conventional convergence results based on unrealistic hypothesis on uniform boundedness of all physical quantities our scheme produces convergent solutions as long as the gas remains in its non-degenerate regime, cf. Section 6.

## 1.1 Complete Euler system

The complete Euler system describes the time evolution of the *standard* physical fields: the mass density  $\varrho = \varrho(t, x)$ , the macroscopic velocity  $\mathbf{u} = \mathbf{u}(t, x)$ , and the (absolute) temperature  $\vartheta = \vartheta(t, x)$  of a perfect compressible fluid,

$$\begin{aligned} \partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) &= 0, \\ \partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p &= 0, \\ \partial_t \left( \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e \right) + \operatorname{div}_x \left[ \left( \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e + p \right) \mathbf{u} \right] &= 0. \end{aligned}$$

For the sake of simplicity, we consider the standard polytropic EOS with the Boyle–Marriot pressure law,

$$p = (\gamma - 1)\varrho e = \varrho\vartheta, \quad e = c_v\vartheta, \quad c_v = \frac{1}{\gamma - 1}.$$

Accordingly, the *physical* entropy reads

$$s(\varrho, \vartheta) = \log\left(\frac{\vartheta^{c_v}}{\varrho}\right)$$

with the associated entropy inequality,

$$\partial_t(\varrho s) + \operatorname{div}_x(\varrho s \mathbf{u}) \geq 0.$$

Note that the same inequality is automatically satisfied by any “renormalized” *mathematical* entropy  $s_\chi$

$$s_\chi = \chi\left(\log\left(\frac{\vartheta^{c_v}}{\varrho}\right)\right),$$

where  $\chi$  is a non-decreasing concave function.

Numerical schemes are based on the *conservative* variables: the density  $\varrho$ , the momentum  $\mathbf{m} = \varrho \mathbf{u}$ , and the total energy

$$E = \frac{1}{2}\varrho|\mathbf{u}|^2 + \varrho e.$$

Accordingly, the Euler system takes the form

$$\partial_t \varrho + \operatorname{div}_x \mathbf{m} = 0, \tag{1.1}$$

$$\partial_t \mathbf{m} + \operatorname{div}_x \left( \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right) + \nabla_x p = 0, \tag{1.2}$$

$$\partial_t E + \operatorname{div}_x \left[ (E + p) \frac{\mathbf{m}}{\varrho} \right] = 0, \tag{1.3}$$

where

$$p = (\gamma - 1) \left( E - \frac{1}{2} \frac{|\mathbf{m}|^2}{\varrho} \right).$$

In the conservative framework, positivity of the density as well as of the pressure becomes an issue, in which the associated entropy balance

$$\partial_t(\varrho s_\chi) + \operatorname{div}_x(s_\chi \mathbf{m}) \geq 0$$

plays a crucial role.

## 1.2 Brenner's model

Brenner's approach to modelling real *viscous* and *heat conducting* fluids postulates two velocities  $\mathbf{u}$  and  $\mathbf{v}$  interrelated through

$$\mathbf{v} = \mathbf{u} - K \nabla_x \log(\varrho).$$

For the Newtonian viscous stress

$$\mathbb{S}(\nabla_x \mathbf{u}) = \eta_1 \left( \nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \eta_2 \operatorname{div}_x \mathbf{u} \mathbb{I}$$

and the Fourier heat flux

$$\mathbf{q} = -\kappa \nabla_x \vartheta$$

the Brenner model reads

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{v}) = 0, \quad (1.4)$$

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{v}) + \nabla_x p = \operatorname{div}_x \mathbb{S}(\nabla_x \mathbf{u}), \quad (1.5)$$

$$\partial_t \left( \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e \right) + \operatorname{div}_x \left[ \left( \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e + p \right) \mathbf{v} \right] + \operatorname{div}_x \mathbf{q} = \operatorname{div}_x (\mathbb{S}(\nabla_x \mathbf{u}) \cdot \mathbf{u}), \quad (1.6)$$

see Brenner [3, 4, 5]. Moreover, if  $K$  is related to the heat conductivity coefficient  $\kappa$  through

$$K = \frac{\kappa}{\varrho c_v},$$

then the associated entropy balance takes the form

$$\begin{aligned} & \partial_t(\varrho s_\chi) + \operatorname{div}_x(\varrho s_\chi \mathbf{v}) - \operatorname{div}_x \left( \frac{\kappa}{c_v} \nabla_x s_\chi \right) \\ &= \frac{\chi'(s)}{\vartheta} \mathbb{S}(\nabla_x \mathbf{u}) : \nabla_x \mathbf{u} + \kappa \chi'(s) |\nabla_x \log(\vartheta)|^2 + \chi'(s) \frac{\kappa}{c_v} |\nabla_x \log(\varrho)|^2 - \chi''(s) \frac{\kappa}{c_v} |\nabla_x s|^2, \end{aligned} \quad (1.7)$$

see Guermond and Popov [19] and [7, Section 4.1].

As observed by Guermond and Popov [19], for the ansatz

$$\mathbb{S}(\nabla_x \mathbf{u}) = h \lambda \varrho \nabla_x \mathbf{u} + h^\alpha \nabla_x \mathbf{u}, \quad \kappa = c_v \varrho K = c_v h \varrho \lambda, \quad \lambda \geq 0,$$

the system (1.4–1.6) rewrites in the conservative variables as

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = h \operatorname{div}_x(\lambda \nabla_x \varrho), \quad (1.8)$$

$$\partial_t \mathbf{m} + \operatorname{div}_x(\mathbf{m} \otimes \mathbf{u}) + \nabla_x p = h \operatorname{div}_x(\lambda \nabla_x \mathbf{m}) + h^\alpha \Delta_x \mathbf{u}, \quad (1.9)$$

$$\partial_t E + \operatorname{div}_x(E \mathbf{u} + p \mathbf{u}) = h \operatorname{div}_x(\lambda \nabla_x E) + h^\alpha \operatorname{div}_x(\nabla_x \mathbf{u} \cdot \mathbf{u}). \quad (1.10)$$

This form, without the  $h^\alpha$ -dependent terms, is strongly reminiscent of some numerical schemes for the complete (inviscid) Euler system based on the finite volume method like the Lax–Friedrichs scheme.

### 1.3 Finite volume scheme

Motivated by Guermond and Popov [19] we propose a finite volume scheme for the complete Euler system based on (1.8–1.10). Although written exclusively in the conservative variables, the scheme relies on convective terms expressed in terms of the velocity  $\mathbf{u}$  rather than the momentum  $\mathbf{m}$ . This allows to minimize the effect of the viscous perturbations - a potential source of deviation from the target Euler system for inviscid flows. Indeed the scheme preserves all the basic properties of the continuous system, in particular, it is entropy stable. Moreover, the positivity of the density and pressure as well as the minimum entropy principle hold.

We then examine the properties of the associated semi-discrete dynamical system. We show that it generates in the asymptotic limit a dissipative measure-valued (DMV) solution of the complete Euler system introduced in [6, 7], see also [14] for the convergence of the Lax–Friedrichs method. Moreover, employing the (DMV)–strong uniqueness principle, we will obtain strong (pointwise) convergence to the unique classical solution as long as the latter exists. In contrast with the standard entropy stable finite volume methods, where convergence analysis is based on rather unrealistic *a priori* hypotheses of uniform boundedness of numerical solutions, cf. Fjordholm, Mishra, Käppeli, Tadmor [15, 16, 17, 24], the convergence for the present scheme is almost unconditional, requiring only a technical hypothesis of boundedness of the numerical temperature and the absence of vacuum.

The paper is organized as follows. Section 2 contains necessary preliminaries including the geometric properties of the mesh and the basic notation used in finite volume methods. Then we introduce the numerical method and the associated semi-discrete dynamical system. In Section 3, we show that the scheme is entropy stable. In Section 4, we study stability of the semi-discrete scheme deriving all necessary *a priori* bounds. Consistency of the scheme, based on a careful analysis of the error terms, is discussed in Section 5. Finally, we perform the limit of vanishing numerical step in Section 6.

## 2 Numerical scheme

We introduce the basic notation, function spaces, and, finally, the numerical scheme.

### 2.1 Preliminaries

We suppose the physical space to be a polyhedral domain  $\Omega_h \subset R^N$ ,  $N = 1, 2, 3$ , that is decomposed into compact elements

$$\bar{\Omega}_h = \bigcup_{K \in \mathcal{T}_h} K.$$

The elements  $K$  are sharing either a common face, edge, or vortex. The mesh  $\mathcal{T}_h$  satisfies the standard regularity assumptions, cf. [9, 10]. The set of all faces is denoted by  $\Sigma$ , while  $\Sigma_{int} = \Sigma \setminus \partial\Omega_h$  stands for the set of all interior faces. Each face is associated with a normal vector  $\mathbf{n}$ . In what



follows, we shall suppose

$$|K|_N \approx h^N, \quad |\sigma|_{N-1} \approx h^{N-1} \text{ for any } K \in \mathcal{T}_h, \sigma \in \Sigma.$$

The symbol  $Q_h$  denotes the set of functions constant on each element  $K$ . For a piecewise (element-wise) continuous function  $v$  we define

$$v^{\text{out}}(x) = \lim_{\delta \rightarrow 0^+} v(x + \delta \mathbf{n}), \quad v^{\text{in}}(x) = \lim_{\delta \rightarrow 0^+} v(x - \delta \mathbf{n}), \quad \bar{v}(x) = \frac{v^{\text{in}}(x) + v^{\text{out}}(x)}{2}, \quad [[v]] = v^{\text{out}}(x) - v^{\text{in}}(x)$$

whenever  $x \in \sigma \in \Sigma_{\text{int}}$ . We recall the product rule

$$[[uv]] = \bar{u}[[v]] + [[u]]\bar{v}.$$

For  $\Phi \in L^1(\Omega_h)$  we define the projection

$$\Pi_h[\Phi] = \sum_{K \in \mathcal{T}_h} 1_K \frac{1}{|K|} \int_K \Phi \, dx \in Q_h(\Omega_h).$$

If  $\Phi \in C^1(\overline{\Omega_h})$  we have

$$\left| [[\Pi_h[\Phi]]] \right|_{\sigma} \lesssim h \|\Phi\|_{C^1}, \quad \left| \Phi - \overline{\Pi_h[\Phi]} \right|_{\sigma} \lesssim h \|\Phi\|_{C^1} \text{ for any } x \in \sigma \in \Sigma_{\text{int}}. \quad (2.1)$$

Here and hereafter the symbol  $A \lesssim B$  means  $A \leq cB$  for a generic positive constant  $c$  independent of  $h$ . If  $\Phi \in C^2(\overline{\Omega_h})$  and  $\mathcal{T}_h$  consists of uniform rectangular/cubic elements, then we moreover have

$$\frac{1}{|\sigma|} \int_{\sigma} \left| \Phi - \overline{\Pi_h[\Phi]} \right| \, dS_h \lesssim h^2 \|\Phi\|_{C^2} \quad \text{for any } \sigma \in \Sigma_{\text{int}}. \quad (2.2)$$

Indeed, any  $C^2$  function can be approximated by the piecewise linear Rannacher–Turek elements [23] (an analogue of the Crouzeix–Raviart elements on rectangles) with the error of  $\mathcal{O}(h^2)$ . Thus, it is enough to show (2.2) for the non-conforming piecewise linear Rannacher–Turek elements. Taking into account their continuity in the center of cell interfaces and the definition of projection  $\Pi_h$ , we only need to show

$$\left| \Phi(S_{\sigma}) - \frac{(\Phi(S_K) + \Phi(S_L))}{2} \right| \lesssim h^2,$$

where  $S_{\sigma}$  denotes the center of gravity of  $\sigma$ ,  $S_K$  and  $S_L$  the centers of gravity of two neighbouring elements  $K$  and  $L$  sharing the common face  $\sigma$ . The latter follows directly from the Taylor expansion.

We further recall the negative  $L^p$ -estimates [9]

$$\|v\|_{L^p(\Omega_h)} \lesssim h^{N \frac{1-p}{p}} \|v\|_{L^1(\Omega_h)} \text{ for any } 1 \leq p \leq \infty, \text{ with } \frac{N(1-p)}{p} = -N \text{ if } p = \infty, \quad (2.3)$$

and the trace inequality

$$\|v\|_{L^p(\partial K)} \lesssim h^{-\frac{1}{p}} \|v\|_{L^p(K)} \text{ for any } 1 \leq p \leq \infty, \quad (2.4)$$

for any  $v \in Q_h(\Omega_h)$ . Moreover, we have a discrete version of the Sobolev embedding theorem, see Chainais–Hillairet, Droniou [8, Lemma 6.1],

$$\|v\|_{L^6(\Omega_h)} \lesssim \|v\|_{L^2(\Omega_h)} + \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \frac{[[v]]^2}{h} dS_h \right)^{1/2} \text{ for any } v \in Q_h(\Omega_h), \quad N = 1, 2, 3. \quad (2.5)$$

Given a velocity  $\mathbf{u} \in Q_h(\Omega_h; R^N)$  and  $r \in Q_h(\Omega_h)$ , we define on each face  $\sigma \in \Sigma_{int}$  an *upwind* of  $r$  by  $\mathbf{u}$  as

$$Up[r, \mathbf{u}] = \bar{r} \bar{\mathbf{u}} \cdot \mathbf{n} - \frac{1}{2} |\bar{\mathbf{u}} \cdot \mathbf{n}| [[r]] = r^{\text{in}} [\bar{\mathbf{u}} \cdot \mathbf{n}]^+ + r^{\text{out}} [\bar{\mathbf{u}} \cdot \mathbf{n}]^-. \quad (2.6)$$

Finally, we set

$$r^{\text{up}} = \begin{cases} r^{\text{in}} & \text{if } \bar{\mathbf{u}} \cdot \mathbf{n} \geq 0 \\ r^{\text{out}} & \text{if } \bar{\mathbf{u}} \cdot \mathbf{n} < 0, \end{cases}, \quad r^{\text{down}} = \begin{cases} r^{\text{out}} & \text{if } \bar{\mathbf{u}} \cdot \mathbf{n} \geq 0 \\ r^{\text{in}} & \text{if } \bar{\mathbf{u}} \cdot \mathbf{n} < 0, \end{cases}, \quad (2.7)$$

and

$$\widetilde{[[r]]} = r^{\text{up}} - r^{\text{down}} = -[[r]] \text{sgn}(\bar{\mathbf{u}} \cdot \mathbf{n}). \quad (2.8)$$

## 2.2 Approximation scheme

In order to properly define the numerical scheme, the boundary conditions must be specified. Here, we adopt the no-flux boundary condition:

$$\mathbf{u}_h \cdot \mathbf{n} = 0, \text{ for any } \sigma \in \partial\Omega_h,$$

and  $\varrho_h, p_h$  are extrapolated, i.e.  $\partial\varrho_h/\partial\mathbf{n} = 0 = \partial p_h/\partial\mathbf{n}$ ,  $\mathbf{n}$  is an outer normal to  $\partial\Omega_h$ . We consider the numerical flux function in the form

$$F_h(r_h, \mathbf{u}_h) = Up[r_h, \mathbf{u}_h] - \mu_h [[r_h]], \quad (2.9)$$

where  $\mu_h \geq 0$  and  $Up[r_h, \mathbf{u}_h]$  is given by (2.6). The quantities  $\varrho_h \in Q_h(\Omega_h)$ ,  $\mathbf{m}_h \in Q_h(\Omega_h; R^N)$ , and  $E_h \in Q_h(\Omega_h)$  at the time level  $t$  are given by the following system of equations:

- **Continuity equation**

$$\int_{\Omega_h} D_t \varrho_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} F_h(\varrho_h, \mathbf{u}_h) [[\Phi]] dS_h = 0 \text{ for any } \Phi \in Q_h(\Omega_h), \quad (2.10)$$

where

$$\mathbf{u}_h = \frac{\mathbf{m}_h}{\varrho_h}.$$

- **Momentum equation**

$$\begin{aligned}
\int_{\Omega_h} D_t \mathbf{m}_h \cdot \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{F}_h(\mathbf{m}_h, \mathbf{u}_h) \cdot [[\Phi]] \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p}_h \mathbf{n} \cdot [[\Phi]] \, dS_h \\
= -h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\Phi]] \, dS_h \text{ for all } \Phi \in Q_h(\Omega_h, R^N),
\end{aligned} \tag{2.11}$$

where

$$p_h = (\gamma - 1) \left( E_h - \frac{1}{2} \frac{|\mathbf{m}_h|^2}{\varrho_h} \right).$$

- **Energy equation**

$$\begin{aligned}
\int_{\Omega_h} D_t E_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} F_h(E_h, \mathbf{u}_h) [[\Phi]] \, dS_h \\
- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p}_h [[\Phi \mathbf{u}_h]] \cdot \mathbf{n} \, dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p}_h \Phi [[\mathbf{u}_h]] \cdot \mathbf{n} \, dS_h \\
= -h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot \overline{\mathbf{u}}_h [[\Phi]] \, dS_h \text{ for all } \Phi \in Q_h(\Omega_h).
\end{aligned} \tag{2.12}$$

Note that our upwinding  $Up[r_h, \mathbf{u}_h]$ ,  $r_h = \varrho_h, \mathbf{m}_h, E_h$ , is based only on the sign of the normal component of velocity, instead of the sign of the eigenvalues as in the standard flux–vector splitting schemes. In addition, numerical diffusion term  $-\mu_h [[r_h]]$  is added to the numerical flux function. The parameter  $\mu_h \geq 0$  is typically of the following form

$$\mu_h = hM(h, \overline{\varrho}_h, \overline{\mathbf{m}}_h, \overline{E}_h),$$

where  $M$  is a continuous function. Unlike the convective terms, the pressure terms are appropriately averaged, cf. (2.11), (2.12). We should note that the terms on the right–hand side of (2.11), (2.12) can be interpreted as the interior penalty terms for the velocity  $\mathbf{u}_h$  that are typically used in the discontinuous Galerkin approach.

In the purely discrete version of (2.10–2.12), the operator  $D_t$  stands for

$$D_t r_h = \frac{r_h(t) - r_h(t - \Delta t)}{\Delta t},$$

where  $\Delta t > 0$  is the time step. In the semi–discrete setting considered in this paper, the functions  $[\varrho_h, \mathbf{m}_h, E_h]$  are continuous functions of the time  $t \in [0, T]$ , and  $D_t$  is interpreted as the standard differential operator,

$$D_t = \frac{d}{dt}.$$

**Remark 2.1.** By virtue of the *product rule*, the integral

$$h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot \overline{\mathbf{u}_h} [[\Phi]] dS_h = \frac{h^{\alpha-1}}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h^2]] [[\Phi]] dS_h$$

may be replaced by a more convenient expression

$$h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\Phi \mathbf{u}_h]] dS_h - h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\Phi} dS_h.$$

**Remark 2.2.** We point out that

$$\sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h} [[\Phi \mathbf{u}_h]] \cdot \mathbf{n} dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h \Phi} [[\mathbf{u}_h]] \cdot \mathbf{n} dS_h \neq \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h} \overline{\mathbf{u}_h} \cdot \mathbf{n} [[\Phi]] dS_h \quad (2.13)$$

as one might expect. Indeed, the left-hand side of (2.13) equals to

$$\sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h} \overline{\mathbf{u}_h} \cdot \mathbf{n} [[\Phi]] dS_h - \frac{1}{4} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] [[\mathbf{u}_h]] \cdot \mathbf{n} [[\Phi]] dS_h. \quad (2.14)$$

This paper is devoted to the semi-discrete version, where  $[\varrho_h, \mathbf{m}_h, E_h]$  are continuous functions of time and the approximate scheme (2.10–2.12) may be interpreted as a finite system of ODEs. It follows from the standard ODE theory that for a given initial state

$$\begin{aligned} \varrho_h(0) &= \varrho_{0,h} \in Q_h(\Omega_h), \quad \varrho_{0,h} > 0, \quad \mathbf{m}(0) = \mathbf{m}_{0,h} \in Q_h(\Omega_h; R^N), \quad E_h(0) = E_{0,h} \in Q_h(\Omega_h), \\ E_{0,h} - \frac{1}{2} \frac{|\mathbf{m}_{0,h}|^2}{\varrho_{0,h}} &> 0, \end{aligned}$$

the semi-discrete system (2.10–2.12) admits a unique solution  $[\varrho_h, \mathbf{m}_h, E_h]$  defined on a maximal time interval  $[0, T_{\max})$ , where

$$\varrho_h(t) > 0, \quad p_h(t) = (\gamma - 1) \left( E_h(t) - \frac{1}{2} \frac{|\mathbf{m}_h(t)|^2}{\varrho_h(t)} \right) > 0 \text{ for all } t \in [0, T_{\max}). \quad (2.15)$$

In particular, the absolute temperature  $\vartheta_h$  can be defined,

$$\vartheta_h(t) = \frac{p_h(t)}{\varrho_h(t)} = \frac{\gamma - 1}{\varrho_h(t)} \left( E_h(t) - \frac{1}{2} \frac{|\mathbf{m}_h(t)|^2}{\varrho_h(t)} \right).$$

As we show in Section 4, the system (2.10–2.12) admits sufficiently strong *a priori* bounds that will guarantee **(i)**  $T_{\max} = \infty$ , **(ii)** validity of (2.15) for any  $t \geq 0$ .

### 3 Entropy balance

We derive a discrete analogue of the entropy balance (1.7) associated to the semi-discrete system (2.10–2.12).

### 3.1 Renormalization

The process of renormalization requires multiplying the discrete equations by nonlinear functions of the unknowns.

#### 3.1.1 Continuity equation

Multiplying the continuity equation (1.8) by  $b'(\varrho)$  we deduce its renormalized form

$$\partial_t b(\varrho) + \operatorname{div}_x(b(\varrho)\mathbf{u}) + \left(b'(\varrho)\varrho - b(\varrho)\right)\operatorname{div}_x\mathbf{u} = h\operatorname{div}_x(\lambda\nabla_x b(\varrho)) - \lambda b''(\varrho)|\nabla_x\varrho|^2.$$

Its discrete analogue (2.10) gives rise to

$$\begin{aligned} & \int_{\Omega_h} \frac{d}{dt} b(\varrho_h)\Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[b(\varrho_h), \mathbf{u}_h][[\Phi]] dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{\mathbf{u}}_h \cdot \mathbf{n} \left[ \left[ (b(\varrho_h) - b'(\varrho_h)\varrho_h)\Phi \right] \right] dS_h \\ & = - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\varrho_h]] \left[ [b'(\varrho_h)\Phi] \right] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \left( \left[ [\widetilde{b(\varrho_h)}] \right] - b'(\varrho_h^{\text{down}})[[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}}_h \cdot \mathbf{n}| dS_h, \end{aligned} \quad (3.1)$$

for any  $\Phi \in Q_h(\Omega_h)$ , see [12, Section 4.1]. Here  $r^{\text{down}}$  and  $[[\widetilde{r_h}]]$  are given by (2.7) and (2.8), respectively.

#### 3.1.2 Transport equation

Under the assumption that  $\varrho$  satisfies (1.8), we consider a field  $b$  satisfying

$$\partial_t(\varrho b) + \operatorname{div}_x(\varrho b\mathbf{u}) = F.$$

Multiplying the equation by  $\chi'(b)$  we obtain

$$\partial_t(\varrho\chi(b)) + \operatorname{div}_x(\varrho\chi(b)\mathbf{u}) = F\chi'(b) + \operatorname{div}_x(h\mu\nabla_x\varrho)(\chi(b) - b\chi'(b)).$$

The discrete version for  $\varrho_h$  satisfying (2.10) reads:

$$\begin{aligned} & \int_{\Omega} \frac{d}{dt} (\varrho_h b_h)\chi'(b_h)\Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h b_h, \mathbf{u}_h][[\chi'(b_h)\Phi]] dS_h \\ & = \int_{\Omega} \frac{d}{dt} \varrho_h \chi(b_h)\Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \chi(b_h), \mathbf{u}_h][[\Phi]] dS_h \\ & + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\varrho_h]] \left[ [(\chi(b_h) - \chi'(b_h)b_h)\Phi] \right] dS_h \\ & + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( \left[ [\widetilde{\chi(b_h)}] \right] - \chi'(b_h^{\text{down}})[[\widetilde{b_h}]] \right) |\overline{\mathbf{u}}_h \cdot \mathbf{n}| dS_h, \end{aligned} \quad (3.2)$$

see [12, Lemma A.1, Section A.2].

## 3.2 Discrete entropy balance equation

We derive a discrete analogue of the entropy balance equation following step by step its derivation in the continuous setting.

### 3.2.1 Discrete kinetic energy equation

The discrete kinetic energy equation is obtained by taking the scalar product of (1.9) with  $\mathbf{u}_h$ , or, at the discrete level, by taking  $\Phi = \mathbf{u}_h \Phi$  in (2.11):

$$\begin{aligned} \frac{d}{dt} \int_{\Omega_h} \mathbf{m}_h \cdot \mathbf{u}_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{F}_h(\mathbf{m}_h, \mathbf{u}_h) \cdot [[\mathbf{u}_h \Phi]] \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \bar{p}_h \mathbf{n} \cdot [[\mathbf{u}_h \Phi]] \, dS_h \\ = -h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h \Phi]] \, dS_h. \end{aligned}$$

Next, we use relation (3.2) for  $b_h = \mathbf{u}_h$ ,  $\chi(|\mathbf{u}_h|) = \frac{1}{2}|\mathbf{u}_h|^2$  to compute

$$\begin{aligned} \frac{d}{dt} \int_{\Omega_h} \mathbf{m}_h \cdot \mathbf{u}_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{U} \mathbf{p}[\mathbf{m}_h, \mathbf{u}_h] \cdot [[\mathbf{u}_h \Phi]] \, dS_h \\ = \frac{d}{dt} \int_{\Omega_h} \varrho_h \mathbf{u}_h \cdot \mathbf{u}_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{U} \mathbf{p}[\varrho_h \mathbf{u}_h, \mathbf{u}_h] \cdot [[\mathbf{u}_h \Phi]] \, dS_h \\ = \frac{d}{dt} \int_{\Omega} \frac{1}{2} \varrho_h |\mathbf{u}_h|^2 \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{U} \mathbf{p} \left[ \frac{1}{2} \varrho_h |\mathbf{u}_h|^2, \mathbf{u}_h \right] [[\Phi]] \, dS_h \\ - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] \left[ \left[ \frac{1}{2} |\mathbf{u}_h|^2 \Phi \right] \right] \, dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 \, dS_h. \end{aligned}$$

Consequently, summing up the previous two observations we may infer that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} \frac{1}{2} \varrho_h |\mathbf{u}_h|^2 \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{U} \mathbf{p} \left[ \frac{1}{2} \varrho_h |\mathbf{u}_h|^2, \mathbf{u}_h \right] [[\Phi]] \, dS_h \\ = -h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h \Phi]] \, dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \bar{p}_h \mathbf{n} \cdot [[\mathbf{u}_h \Phi]] \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\mathbf{m}_h]] [[\mathbf{u}_h \Phi]] \, dS_h \\ + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] \left[ \left[ \frac{1}{2} |\mathbf{u}_h|^2 \Phi \right] \right] \, dS_h - \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 \, dS_h. \end{aligned} \tag{3.3}$$

Equation (3.3) is nothing other than the discrete kinetic energy balance associated to the approximate system (2.10–2.12).

### 3.2.2 Discrete internal energy equation

The next step is subtracting (3.3) from the total energy balance (2.12):

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h e_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( Up[\varrho_h e_h, \mathbf{u}_h] - \mu_h[[E_h]] \right) [[\Phi]] dS_h \\
&= -h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\Phi \mathbf{u}_h]] dS_h + h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \bar{\Phi} dS_h \\
&+ h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h \Phi]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h \Phi} \mathbf{n} \cdot [[\mathbf{u}_h]] dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\mathbf{m}_h]] [[\mathbf{u}_h \Phi]] dS_h \\
&- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\varrho_h]] \left[ \left[ \frac{1}{2} |\mathbf{u}_h|^2 \Phi \right] \right] dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h,
\end{aligned}$$

or, reordered,

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h e_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( Up[\varrho_h e_h, \mathbf{u}_h] - \mu_h[[\varrho_h e_h]] \right) [[\Phi]] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \bar{\Phi} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h \Phi} [[\mathbf{u}_h]] \cdot \mathbf{n} dS_h \\
&+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\varrho_h \mathbf{u}_h]] [[\mathbf{u}_h \Phi]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h[[\varrho_h]] \left[ \left[ \frac{1}{2} |\mathbf{u}_h|^2 \Phi \right] \right] dS_h \\
&- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \left[ \left[ \frac{1}{2} \varrho_h |\mathbf{u}_h|^2 \right] \right] [[\Phi]] dS_h.
\end{aligned}$$

Finally, using the product rule, we obtain

$$\begin{aligned}
& [[\varrho_h \mathbf{u}_h]] [[\mathbf{u}_h \Phi]] - \frac{1}{2} [[\varrho_h]] [ [ |\mathbf{u}_h|^2 \Phi ] ] - \frac{1}{2} [[\varrho_h |\mathbf{u}_h|^2]] [[\Phi]] \\
&= \overline{\varrho_h} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h]] \bar{\Phi} + \overline{\varrho_h} \bar{\mathbf{u}}_h \cdot [[\mathbf{u}_h]] [[\Phi]] \\
&\quad + \frac{1}{2} [[\varrho_h]] \bar{\mathbf{u}}_h \cdot [[\mathbf{u}_h \Phi]] - \frac{1}{2} [[\varrho_h]] \bar{\mathbf{u}}_h \cdot [[\mathbf{u}_h]] \bar{\Phi} - \frac{1}{2} [[\varrho_h |\mathbf{u}_h|^2]] [[\Phi]] \\
&= \overline{\varrho_h} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h]] \bar{\Phi} + \overline{\varrho_h} \bar{\mathbf{u}}_h \cdot [[\mathbf{u}_h]] [[\Phi]] + \frac{1}{2} [[\varrho_h]] |\bar{\mathbf{u}}_h|^2 [[\Phi]] - \frac{1}{2} [[\varrho_h |\mathbf{u}_h|^2]] [[\Phi]] \\
&= \overline{\varrho_h} [[\mathbf{u}_h]] \cdot [[\mathbf{u}_h]] \bar{\Phi} + \overline{\varrho_h} \bar{\mathbf{u}}_h \cdot [[\mathbf{u}_h]] [[\Phi]] - \frac{1}{2} \overline{\varrho_h} [[\mathbf{u}_h \cdot \mathbf{u}_h]] [[\Phi]] = \overline{\varrho_h} [[\mathbf{u}_h]]^2 \bar{\Phi}.
\end{aligned}$$

Consequently, we record the internal energy balance in the form

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h e_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( Up[\varrho_h e_h, \mathbf{u}_h] - \mu_h [[\varrho_h e_h]] \right) [[\Phi]] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\Phi} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\
&+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\Phi} dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h} \overline{\Phi} [[\mathbf{u}_h]] \cdot \mathbf{n} dS_h.
\end{aligned} \tag{3.4}$$

### 3.2.3 Discrete entropy balance

At this stage, we are ready to derive the discrete entropy balance together with its renormalization. Dividing equation (3.4) on  $\vartheta_h$ , we get

$$\begin{aligned}
& c_v \int_{\Omega_h} \frac{d}{dt} (\varrho_h \vartheta_h) \left( \frac{\Phi}{\vartheta_h} \right) \, dx - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \vartheta_h, \mathbf{u}_h] \left[ \left[ \frac{\Phi}{\vartheta_h} \right] \right] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\Phi}{\vartheta_h} \right)} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( \frac{\Phi}{\vartheta_h} \right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\
&+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\Phi}{\vartheta_h} \right)} dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot \mathbf{n} \overline{\varrho_h} \overline{\Phi} dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\Phi}{\vartheta_h} \right] \right] dS_h.
\end{aligned}$$

Next, by virtue of formula (3.2),

$$\begin{aligned}
& c_v \int_{\Omega_h} \frac{d}{dt} (\varrho_h \vartheta_h) \left( \frac{\Phi}{\vartheta_h} \right) \, dx - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \vartheta_h, \mathbf{u}_h] \left[ \left[ \frac{\Phi}{\vartheta_h} \right] \right] dS_h \\
&= \frac{d}{dt} \int_{\Omega} \varrho_h \log(\vartheta_h^{c_v}) \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mathbf{U} \mathbf{P}[\varrho_h \log(\vartheta_h^{c_v}), \mathbf{u}_h] [[\Phi]] dS_h \\
&+ c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu [[\varrho_h]] [(\log(\vartheta_h) - 1) \Phi] dS_h \\
&+ c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h.
\end{aligned}$$



Consequently,

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h \log(\vartheta_h^{c_v}) \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \log(\vartheta_h^{c_v}), \mathbf{u}_h][[\Phi]] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left(\frac{\Phi}{\vartheta_h}\right)} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left(\frac{\Phi}{\vartheta_h}\right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\
&+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left(\frac{\Phi}{\vartheta_h}\right)} dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot \mathbf{n} \overline{\varrho_h} \Phi dS_h \\
&- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\Phi}{\vartheta_h} \right] \right] dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\log(\vartheta_h) - 1) \Phi] dS_h \\
&- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h.
\end{aligned} \tag{3.5}$$

Finally, we consider  $b(\varrho) = \varrho \log(\varrho)$  in the renormalized equation (3.1):

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h \log(\varrho_h) \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \log(\varrho_h), \mathbf{u}_h][[\Phi]] dS_h \\
&= - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [b'(\varrho_h) \Phi] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \left( [[\widetilde{b(\varrho_h)}]] - b'(\varrho_h^{\text{down}}) [[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot \mathbf{n} \overline{\varrho_h} \Phi dS_h.
\end{aligned} \tag{3.6}$$

Subtracting (3.6) from (3.5) and introducing the entropy  $s_h = \log\left(\frac{\vartheta_h^{c_v}}{\varrho_h}\right)$  we obtain

$$\begin{aligned}
& \frac{d}{dt} \int_{\Omega_h} \varrho_h s_h \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h s_h, \mathbf{u}_h][[\Phi]] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left(\frac{\Phi}{\vartheta_h}\right)} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left(\frac{\Phi}{\vartheta_h}\right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\
&+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left(\frac{\Phi}{\vartheta_h}\right)} dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \left( [[\widetilde{b(\varrho_h)}]] - b'(\varrho_h^{\text{down}}) [[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\Phi}{\vartheta_h} \right] \right] dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\log(\vartheta_h) - 1) \Phi] dS_h
\end{aligned} \tag{3.7}$$

$$+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[b'(\varrho_h)\Phi]] dS_h, \quad \text{where } b(\varrho) = \varrho \log(\varrho).$$

This is the physical entropy balance associated to (2.10–2.12). At this stage, it is not obvious how to handle the last three integrals in (3.7), however, this will be fixed in the following section.

### 3.2.4 Entropy renormalization

Consider  $\chi$  - a non-decreasing, concave, twice continuously differentiable function on  $R$  that is bounded from above. Applying formula (3.2) in (3.7) we get

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega_h} \varrho_h \chi(s_h) \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \chi(s_h), \mathbf{u}_h] [[\Phi]] dS_h \\ &= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)} dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\chi'(s_h) \Phi)^{\text{down}} \left( [[\widetilde{b(\varrho_h)}]] - b'(\varrho_h^{\text{down}}) [[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\ &- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\chi'(s_h) \Phi)^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\ &- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\chi(s_h)}]] - \chi'(s_h^{\text{down}}) [[\widetilde{s_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\ &- c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\chi'(s_h) \Phi}{\vartheta_h} \right] \right] dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\log(\vartheta_h) - 1) \chi'(s_h) \Phi] dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[b'(\varrho_h) \chi'(s_h) \Phi]] dS_h \\ &- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\chi(s_h) - \chi'(s_h) s_h) \Phi] dS_h, \quad \text{where } b(\varrho) = \varrho \log(\varrho). \end{aligned}$$

Next, we compute

$$\begin{aligned} & - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\chi'(s_h) \Phi}{\vartheta_h} \right] \right] dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\log(\vartheta_h) - 1) \chi'(s_h) \Phi] dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[b'(\varrho_h) \chi'(s_h) \Phi]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [(\chi(s_h) - \chi'(s_h) s_h) \Phi] dS_h \end{aligned}$$

$$\begin{aligned}
&= -c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \frac{\chi'(s_h) \Phi}{\vartheta_h} \right] \right] dS_h - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\log(\vartheta_h) \chi'(s_h) \Phi]] dS_h \\
&\quad + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\log(\varrho_h) \chi'(s_h) \Phi]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[(\chi(s_h) - \chi'(s_h) s_h) \Phi]] dS_h \\
&\quad + (c_v + 1) \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\chi'(s_h) \Phi]] dS_h \\
&= -c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \vartheta_h]] \left[ \left[ \varrho_h \frac{\chi'(s_h) \Phi}{\varrho_h \vartheta_h} \right] \right] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] \left[ \left[ (\chi(s_h) - (c_v + 1) \chi'(s_h)) \Phi \right] \right] dS_h \\
&= -c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[p_h]] \left[ \left[ \varrho_h \frac{\chi'(s_h) \Phi}{p_h} \right] \right] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] \left[ \left[ (\chi(s_h) - (c_v + 1) \chi'(s_h)) \Phi \right] \right] dS_h \\
&= - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\Phi \nabla_{\varrho}(\varrho_h \chi(s_h))]] [[\varrho_h]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\Phi \nabla_p(\varrho_h \chi(s_h))]] [[p_h]] dS_h.
\end{aligned}$$

Thus we infer with the general entropy inequality

$$\begin{aligned}
&\frac{d}{dt} \int_{\Omega_h} \varrho_h \chi(s_h) \Phi \, dx - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h \chi(s_h), \mathbf{u}_h][[\Phi]] dS_h \\
&= h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)} dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h \\
&\quad + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left( \frac{\chi'(s_h) \Phi}{\vartheta_h} \right)} dS_h \\
&\quad + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\chi'(s_h) \Phi)^{\text{down}} \left( [[\widetilde{b(\varrho_h)}]] - b'(\varrho_h^{\text{down}}) [[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&\quad - c_v \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\chi'(s_h) \Phi)^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&\quad - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \Phi^{\text{down}} \varrho_h^{\text{up}} \left( [[\widetilde{\chi(s_h)}]] - \chi'(s_h^{\text{down}}) [[\widetilde{s_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h \\
&\quad - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\Phi \nabla_{\varrho}(\varrho_h \chi(s_h))]] [[\varrho_h]] dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\Phi \nabla_p(\varrho_h \chi(s_h))]] [[p_h]] dS_h, \quad b(\varrho) = \varrho \log(\varrho).
\end{aligned} \tag{3.8}$$

Note that the last two integrals in (3.8) can be rewritten using the product rule as

$$\begin{aligned}
& - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\Phi]] \left( \overline{\nabla_{\varrho}(\varrho_h \chi(s_h))} [[\varrho_h]] + \overline{\nabla_p(\varrho_h \chi(s_h))} [[p_h]] \right) dS_h \\
& - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\Phi} [[\nabla_{\varrho}(\varrho_h \chi(s_h))]] [[\varrho_h]] + [[\nabla_p(\varrho_h \chi(s_h))]] [[p_h]] dS_h.
\end{aligned} \tag{3.9}$$

The first sum in (3.9) together with the upwind term in (3.8),

$$- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( Up[\varrho_h \chi(s_h), \mathbf{u}_h] + \mu_h \left( \overline{\nabla_{\varrho}(\varrho_h \chi(s_h))} [[\varrho_h]] + \overline{\nabla_p(\varrho_h \chi(s_h))} [[p_h]] \right) \right) [[\Phi]] dS_h, \tag{3.10}$$

represent the numerical entropy flux. The rest in (3.8) and (3.9) gives the numerical entropy production, cf. [14, 17, 16]. Recall that the total entropy

$$(\varrho, p) \mapsto -\varrho \chi(s(\varrho, p)) = -\varrho \chi \left( \log \left( \frac{\varrho^{c_v}}{\varrho} \right) \right) = -\varrho \chi \left( \frac{1}{\gamma - 1} \log \left( \frac{p}{\varrho^{\gamma}} \right) \right)$$

is a convex function of the variables  $\varrho$  and  $p$ . In particular,  $-\nabla_{\varrho, p}(\varrho \chi(s(\varrho_h, p_h)))$  is monotone, and therefore the term in the second line of (3.9) is non-negative. It is worthwhile to mention that the discrete entropy inequality (3.8) is a discrete version of (1.7) with  $\kappa = c_v h \varrho \lambda$ ,  $\lambda = \frac{1}{2} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| + \mu_h$ .

## 4 Stability

Having established all necessary ingredients, we are ready to discuss the available *a priori* bounds for solutions of the semi-discrete scheme (2.10–2.12).

### 4.1 Mass and energy conservation

Taking  $\Phi \equiv 1$  in the equation of continuity (2.10) yields the total mass conservation

$$\int_{\Omega_h} \varrho_h(t, \cdot) dx = \int_{\Omega_h} \varrho_{0,h} dx = M_0 > 0, \quad t \geq 0. \tag{4.1}$$

A similar argument applied to the total energy balance yields

$$\int_{\Omega_h} E_h(t, \cdot) dx = \int_{\Omega_h} E_{0,h} dx = E_0 > 0, \quad t \geq 0. \tag{4.2}$$

## 4.2 Minimum entropy principle

An important source of *a priori* bounds is the minimum entropy principle that can be derived from the entropy balance with the choice

$$\Phi = 1, \quad \chi(s) = |s - s_0|^{-}, \quad -\infty < s_0 < \min s_h(0).$$

As

$\varrho \mapsto \varrho \log(\varrho)$  is convex,  $\vartheta \mapsto \log(\vartheta)$  concave,  $s \mapsto \chi(s)$  concave, and  $(\varrho, p) \mapsto \varrho \chi(s(\varrho, p))$  concave, all integrals on the right-hand side of (3.8) are non-negative, and we may infer that

$$\int_{\Omega_h} \varrho_h(t) |s_h(t) - s_0|^{-} dx \geq 0 \quad \text{for any } t \geq 0.$$

Consequently, we have obtained the minimum entropy principle

$$s_h(t) \geq s_0 \quad \text{for all } t \geq 0. \quad (4.3)$$

## 4.3 Positivity of the pressure, existence of the temperature

The entropy as a function of  $\varrho$  and  $p$  reads

$$s = \frac{1}{\gamma - 1} \log \left( \frac{p}{\varrho^\gamma} \right);$$

whence it follows immediately from (4.3) that

$$0 < \exp\{(\gamma - 1)s_0\} \leq \frac{p_h(t)}{\varrho_h^\gamma(t)} \quad \text{for all } t \geq 0. \quad (4.4)$$

In particular, the pressure is positive as long as the density is positive, and we may set

$$\vartheta_h(t) = \frac{p_h(t)}{\varrho_h(t)}.$$

Evoking the energy bound (4.2) we get

$$\frac{1}{2} \int_{\Omega_h} \frac{|\mathbf{m}_h(t)|^2}{\varrho_h(t)} dx + c_v \int_{\Omega_h} \varrho_h(t) \vartheta_h(t) dx \leq E_0 \quad \text{for all } t \geq 0. \quad (4.5)$$

Thus going back to (4.4) we obtain

$$\int_{\Omega_h} \varrho_h^\gamma(t) dx \lesssim \int_{\Omega_h} p_h(t) dx \lesssim E_0 \quad \text{for all } t \geq 0. \quad (4.6)$$

#### 4.4 Positivity of the density

The crucial property for the approximate scheme to be valid is positivity of the density  $\varrho_h$  at least at the discrete level, meaning for any  $h > 0$ . We will show that, for any  $T > 0$ , there exists  $\underline{\varrho} = \underline{\varrho}(h, T) > 0$ , such that  $\varrho_h(t) \geq \underline{\varrho} > 0$  for all  $t \in [0, T]$ . To see this, we first evoke the kinetic energy balance (3.3) with  $\Phi = 1$ . Seeing that

$$- \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\mathbf{m}_h]] [[\mathbf{u}_h]] dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] \left[ \left[ \frac{1}{2} |\mathbf{u}_h|^2 \right] \right] dS_h = - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 dS_h,$$

we may integrate (3.3) in time and use the energy bound (4.5) to deduce

$$\begin{aligned} & h^{\alpha-1} \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h dt + \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 dS_h dt \\ & + \frac{1}{2} \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \varrho_h^{up} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h dt \lesssim \left( 1 + \sum_{\sigma \in \Sigma_{int}} \int_0^T \int_{\sigma} \overline{p_h} \mathbf{n} \cdot [[\mathbf{u}_h]] dS_h \right) dt. \end{aligned}$$

Finally, we again use (4.5) combined with the negative  $L^p$ -estimates (2.3) and Hölder's inequality to conclude

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h dt \lesssim \omega(h), \quad (4.7)$$

where  $\omega(h)$  denotes a generic function that may blow up in the asymptotic regime  $h \rightarrow 0$ . In particular, relation (4.7) implies

$$\int_0^T \left( \sup_{\sigma \in \Sigma_{int}} [[\mathbf{u}_h]]^2 \right) dt \lesssim \omega(h), \quad (4.8)$$

with another  $\omega(h)$  generally different from its counterpart in (4.7).

Next, we revisit the renormalized equation of continuity (3.1), again with  $\Phi = 1$ , obtaining

$$\int_{\Omega_h} \frac{d}{dt} b(\varrho_h) dx + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\overline{\mathbf{u}_h}]] \cdot \mathbf{n} \overline{(b(\varrho_h) - b'(\varrho_h)\varrho_h)} dS_h \leq 0$$

for any convex  $b$ . Thus the specific choice  $b(\varrho) = |\varrho - \underline{\varrho}|^-$  gives rise to the inequality

$$\frac{d}{dt} \int_{\Omega_h} |\varrho_h - \underline{\varrho}|^- dx + \underline{\varrho} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\overline{\mathbf{u}_h}]] \cdot \mathbf{n} \overline{\mathbb{1}_{\varrho_h(t) \leq \underline{\varrho}}} dS_h \leq 0.$$

In view of (4.8), we can find a positive constant  $\underline{\varrho} = \underline{\varrho}(h, T) > 0$  small enough so that

$$\int_{\Omega_h} |\varrho_h(t) - \underline{\varrho}|^- dx < 0 \quad \text{for all } t \in [0, T].$$

In other words

$$\varrho_h(t) \geq \underline{\varrho}(h, T) > 0 \quad \text{for all } t \in [0, T]. \quad (4.9)$$

**Remark 4.1.** Of course, the estimate (4.9) is not uniform, neither with respect to  $T$  nor for  $h \rightarrow 0$ . In particular, the asymptotic limit may experience vacuum zone where the density vanishes.

## 4.5 Existence of approximate solutions

Having established positivity of the density on any compact time interval, we have closed the *a priori* bounds that guarantee global existence for the semi-discrete system at any level  $h > 0$ .

**Theorem 4.2.** *Suppose that the initial data  $\varrho_{0,h}$ ,  $\mathbf{m}_{0,h}$ ,  $E_{0,h}$  satisfy*

$$\varrho_{0,h} \geq \underline{\varrho} > 0, \quad E_{0,h} - \frac{1}{2} \frac{|\mathbf{m}_{0,h}|^2}{\varrho_{0,h}} > 0.$$

*Then the semi-discrete approximate system (2.10–2.12) admits a unique global-in-time solution  $[\varrho_h, \mathbf{m}_h, E_h]$  such that*

$$\varrho_h(t) > 0, \quad E_h(t) - \frac{1}{2} \frac{|\mathbf{m}_h(t)|^2}{\varrho_h(t)} > 0 \quad \text{for any } t \geq 0.$$

*Moreover, the renormalized entropy balance (3.8) holds.*

## 4.6 Entropy estimates

We close this section by showing the uniform bounds provided by the dissipation mechanism hidden in the entropy production rate. We start by observing that

$$\int_{\Omega_h} \varrho_h s_h(t) \, dx \lesssim 1 + \int_{\Omega_h} E_h(t) \, dx \leq 1 + E_0. \quad (4.10)$$

Indeed, in view of the minimum entropy principle established in (4.4), it is enough to observe that

$$\varrho_h \log \left( \frac{\vartheta_h^{c_v}}{\varrho_h} \right) \lesssim 1 + \varrho_h \vartheta_h \quad \text{provided } 0 < \varrho_h \lesssim \vartheta_h^{c_v}.$$

Seeing that  $\varrho_h \log(\varrho_h)$  is controlled by (4.6) we restrict ourselves to  $\varrho_h \log(\vartheta_h^{c_v})$ . Here,

$$\varrho_h \log(\vartheta_h^{c_v}) \lesssim \varrho_h \vartheta_h \lesssim E_0 \quad \text{if } \vartheta_h \geq 1,$$

while

$$|\varrho_h \log(\vartheta_h^{c_v})| \leq \vartheta_h^{c_v} |\log(\vartheta_h^{c_v})| \lesssim 1 \quad \text{for } \vartheta_h \leq 1.$$

Thus we have shown (4.10).

In accordance with (4.10), we can take  $\Phi = 1$ ,  $\chi_\varepsilon(s) = \min\{s, \frac{1}{\varepsilon}\}$  in the renormalized entropy balance (3.8). Letting  $\varepsilon \rightarrow 0$  we obtain the uniform estimate:

$$\begin{aligned}
& h^{\alpha-1} \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left(\frac{1}{\vartheta_h}\right)} dS_h dt + \frac{1}{2} \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left(\frac{1}{\vartheta_h}\right)^{\text{down}} \varrho_h^{\text{up}} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h dt \\
& + \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \bar{\varrho}_h [[\mathbf{u}_h]]^2 \overline{\left(\frac{1}{\vartheta_h}\right)} dS_h dt + \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \left( [[b(\varrho)] ] - b'(\varrho_h^{\text{down}}) [[\varrho_h]] \right) |\bar{\mathbf{u}}_h \cdot \mathbf{n}| dS_h dt \\
& - c_v \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \varrho_h^{\text{down}} \left( [[\log(\vartheta_h)] ] - \frac{1}{\vartheta_h^{\text{down}}} [[\vartheta_h]] \right) |\bar{\mathbf{u}}_h \cdot \mathbf{n}| dS_h dt \\
& - \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\nabla_{\varrho}(\varrho_h s_h)] ] [[\varrho_h]] dS_h dt - \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\nabla_p(\varrho_h s_h)] ] [[p_h]] dS_h dt \lesssim (1 + E_0),
\end{aligned} \tag{4.11}$$

where  $b(\varrho) = \varrho \log(\varrho)$ . As for the last two integrals in (4.11), we can check by direct manipulation that

$$\begin{aligned}
& - \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\nabla_{\varrho}(\varrho_h s_h)] ] [[\varrho_h]] dS_h dt - \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\nabla_p(\varrho_h s_h)] ] [[p_h]] dS_h dt \\
& = -c_v \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \bar{\varrho}_h [[\vartheta_h]] \left[ \left[ \frac{1}{\vartheta_h} \right] \right] dS_h dt + \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\log(\varrho_h)] ] dS_h dt \\
& - c_v \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \left( [[\log(\vartheta_h)] ] + \bar{\vartheta}_h \left[ \left[ \frac{1}{\vartheta_h} \right] \right] \right) [[\varrho_h]] dS_h dt.
\end{aligned}$$

Next, we show that

$$- [[\varrho_h]] \left( [[\log(\vartheta_h)] ] + \bar{\vartheta}_h \left[ \left[ \frac{1}{\vartheta_h} \right] \right] \right) \leq -\frac{1}{2} | [[\varrho_h]] | [[\vartheta_h]] \left[ \left[ \frac{1}{\vartheta_h} \right] \right]. \tag{4.12}$$

As both expression in the above inequality are invariant with respect to the change “in” and “out” and, in addition, the right-hand side is invariant with respect to the same operation in  $\varrho_h$  and  $\vartheta_h$  separately, it is enough to show (4.12) assuming  $\varrho_h^{\text{in}} \geq \varrho_h^{\text{out}}$ . In other words,

$$- [[\varrho_h]] = | [[\varrho_h]] | \geq 0.$$

Consequently, the proof of (4.12) reduces to the inequality

$$[[\log(\vartheta_h)] ] + \bar{\vartheta}_h \left[ \left[ \frac{1}{\vartheta_h} \right] \right] \leq -\frac{1}{2} [[\vartheta_h]] \left[ \left[ \frac{1}{\vartheta_h} \right] \right].$$



Denoting  $Z = \frac{\vartheta_h^{\text{out}}}{\vartheta_h^{\text{in}}}$ , we have to show

$$\log(Z) - \frac{1}{2} \left( Z - \frac{1}{Z} \right) \leq \frac{1}{2} \left( Z + \frac{1}{Z} \right) - 1$$

or

$$\log(Z) \leq Z - 1,$$

which is obvious as  $\log$  is a concave function. In view of (4.12), relation (4.11) yields

$$\begin{aligned} & h^{\alpha-1} \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} [[\mathbf{u}_h]]^2 \overline{\left( \frac{1}{\vartheta_h} \right)} dS_h dt + \frac{1}{2} \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \left( \frac{1}{\vartheta_h} \right)^{\text{down}} \varrho_h^{\text{up}} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h dt \\ & + \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \mu_h \overline{\varrho_h} [[\mathbf{u}_h]]^2 \overline{\left( \frac{1}{\vartheta_h} \right)} dS_h dt + \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \left( [[\widetilde{b(\varrho_h)}]] - b'(\varrho_h^{\text{down}}) [[\widetilde{\varrho_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h dt \\ & - c_v \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \varrho_h^{\text{up}} \left( [[\widetilde{\log(\vartheta_h)}]] - \frac{1}{\vartheta_h^{\text{down}}} [[\widetilde{\vartheta_h}]] \right) |\overline{\mathbf{u}_h} \cdot \mathbf{n}| dS_h dt \\ & - c_v \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \mu_h \min\{\varrho_h^{\text{in}}, \varrho_h^{\text{out}}\} [[\vartheta_h]] \left[ \left[ \frac{1}{\vartheta_h} \right] \right] dS_h dt + \int_0^T \sum_{\sigma \in \Sigma_{\text{int}}} \int_{\sigma} \mu_h [[\varrho_h]] [[\log(\varrho_h)]] dS_h dt \\ & \lesssim (1 + E_0). \end{aligned} \tag{4.13}$$

## 5 Consistency

We show consistency of the scheme (2.10–2.12), meaning the approximate solutions satisfy the weak formulation of the problem modulo approximation errors vanishing in the asymptotic limit  $h \rightarrow 0$ .

### 5.1 Numerical flux

Firstly, we handle the numerical fluxes in (2.10), (2.11) and the numerical entropy flux (3.10) consisting of the upwind and  $\mu_h$ -dependent terms.

#### 5.1.1 Upwinds

The upwind terms in the continuity equation (2.10), momentum equation (2.11), and the renormalized entropy balance (3.8) read

$$\int_{\Omega_h} \left( \frac{\varrho_h}{\vartheta_h} \overline{\mathbf{u}_h} \cdot \mathbf{n} - \frac{1}{2} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\varrho]] \right) [[\Phi]] dx,$$

$$\int_{\Omega_h} \left( \overline{\mathbf{m}_h} (\overline{\mathbf{u}_h} \cdot \mathbf{n}) - \frac{1}{2} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\mathbf{m}_h]] \right) \cdot [[\Phi]] \, dx,$$

and  $\int_{\Omega_h} \left( \overline{\varrho_h \chi(s_h)} \overline{\mathbf{u}_h} \cdot \mathbf{n} - \frac{1}{2} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\varrho_h \chi(s_h)]] \right) [[\Phi]] \, dx$ , respectively.

For  $\Phi \in C^1(\overline{\Omega_h})$  we get

$$\begin{aligned} \int_{\Omega_h} \varrho_h b_h \mathbf{u}_h \cdot \nabla_x \Phi \, dx &= \sum_{K \in \mathcal{T}_h} \int_K \varrho_h b_h \mathbf{u}_h \cdot \nabla_x \Phi \, dx = \sum_{K \in \mathcal{T}_h} \int_{\partial K} \varrho_h b_h \mathbf{u}_h \cdot \mathbf{n} \Phi \, dS_h \\ &= \sum_{K \in \mathcal{T}_h} \int_{\partial K} \varrho_h b_h \mathbf{u}_h \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h + \sum_{K \in \mathcal{T}_h} \int_{\partial K} \varrho_h b_h \mathbf{u}_h \cdot \mathbf{n} \overline{\Pi_h[\Phi]} \, dS_h \\ &= - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \overline{\Pi_h[\Phi]} \, dS_h \\ &= - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{\varrho_h b_h \mathbf{u}_h} \cdot \mathbf{n} [[\Pi_h[\Phi]]] \, dS_h \\ &= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{\varrho_h b_h} \overline{\mathbf{u}_h} \cdot \mathbf{n} [[\Pi_h[\Phi]]] \, dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\overline{\varrho_h b_h \mathbf{u}_h} - \overline{\varrho_h b_h} \overline{\mathbf{u}_h}) \cdot \mathbf{n} [[\Pi_h[\Phi]]] \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h \\ &= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} Up[\varrho_h b_h] [[\Pi[\Phi]]] \, dS_h + \frac{1}{2} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\varrho_h b_h]] [[\Pi[\Phi]]] \, dS_h \\ &+ \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} (\overline{\varrho_h b_h \mathbf{u}_h} - \overline{\varrho_h b_h} \overline{\mathbf{u}_h}) \cdot \mathbf{n} [[\Pi_h[\Phi]]] \, dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h. \end{aligned}$$

Seeing that

$$\overline{uv} - \overline{u} \overline{v} = \frac{1}{4} [[u]] [[v]]$$

we have to control the following error terms:

$$\begin{aligned} E_1 &= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\varrho_h b_h]] [[\Pi[\Phi]]] \, dS_h, \\ E_2 &= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} \left( \Phi - \overline{\Pi_h[\Phi]} \right) \, dS_h, \\ E_3 &= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\varrho_h b_h]] [[\mathbf{u}_h]] \cdot \mathbf{n} [[\Pi[\Phi]]] \, dS_h, \end{aligned}$$

where  $b_h$  is either 1 or  $\chi(s_h)$  or  $u_h^j$ ,  $j = 1, \dots, N$ . In view of (2.1) and the identity

$$[[\varrho_h b_h \mathbf{u}_h]] \cdot \mathbf{n} = [[\varrho_h b_h]] \overline{\mathbf{u}_h} \cdot \mathbf{n} + \overline{\varrho_h b_h} [[\mathbf{u}_h]] \cdot \mathbf{n},$$

it is enough to show that

$$\begin{aligned}
E_{1,h} &= h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| | [[\varrho_h b_h]] | dS_h \rightarrow 0, \\
E_{2,h} &= h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\varrho_h b_h}| | [[\mathbf{u}_h]] \cdot \mathbf{n} | dS_h \rightarrow 0, \\
E_{3,h} &= h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} | [[\varrho_h b_h]] | | [[\mathbf{u}_h]] \cdot \mathbf{n} | dS_h \rightarrow 0
\end{aligned} \tag{5.1}$$

as  $h \rightarrow 0$  for any fixed  $\Phi \in C^1(\overline{\Omega}_h)$ . Moreover, by virtue of the minimum entropy principle (4.4), the entropy  $s_h$  is bounded below uniformly for  $h \rightarrow 0$ . As the cut-off function  $\chi$  is supposed to be bounded from above, we may assume

$$|\chi(s_h)| \lesssim 1 \text{ for } h \rightarrow 0.$$

The following analysis leans heavily on the bound

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h dt \lesssim h^{1-\alpha} \tag{5.2}$$

that follows directly from the entropy estimates (4.13) provided

$$0 < \vartheta_h \lesssim 1 \text{ uniformly for } h \rightarrow 0. \tag{5.3}$$

Accordingly, we *suppose* that the approximate solutions satisfy (5.3). Then, as  $\gamma > 1$ , the entropy minimum principle (4.4) yields a similar bound on the density,

$$0 < \varrho_h \lesssim 1 \text{ uniformly for } h \rightarrow 0. \tag{5.4}$$

With (5.3), (5.4) at hand, the convergence of the errors  $E_{2,h}$ ,  $E_{3,h}$  for  $b_h = 1$  and  $b_h = \chi(s_h)$  reduces to showing

$$h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} | [[\mathbf{u}_h]] | dS_h \rightarrow 0.$$

To see this, we use Hölder's inequality,

$$\begin{aligned}
h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} | [[\mathbf{u}_h]] | dS_h &\leq h \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} 1 dS_h \right)^{1/2} \\
&\lesssim \sqrt{h} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \lesssim h^{1-\frac{\alpha}{2}} F_h^1, \quad \|F_h^1\|_{L^2(0,T)} \lesssim 1,
\end{aligned}$$

where the last inequality follows from the hypothesis (5.2).

In order to control the integral in  $E_{1,h}$  we need bounds on the velocity  $\mathbf{u}_h$ . They can be deduced from the total energy balance (4.5) if we make another extra *hypothesis*, namely

$$0 < \underline{\varrho} \leq \varrho_h \text{ uniformly for all } h \rightarrow 0. \quad (5.5)$$

In view of (4.4) this implies a similar lower bound on the approximate temperature,

$$0 < \underline{\vartheta} \leq \vartheta_h \text{ uniformly for all } h \rightarrow 0. \quad (5.6)$$

Under these circumstances, we easily deduce from (4.5), (4.13) the following bounds:

$$\sup_{t \in [0, T]} \|\mathbf{u}_h(t)\|_{L^2(\Omega_h)} \lesssim 1, \quad (5.7)$$

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\mathbf{u}_h]]^2 dS_h dt \lesssim 1, \quad (5.8)$$

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\varrho_h]]^2 dS_h dt \lesssim 1, \quad (5.9)$$

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| [[\vartheta_h]]^2 dS_h dt \lesssim 1. \quad (5.10)$$

In particular, we obtain the estimates

$$h^{\alpha-1} \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h dt \lesssim 1, \quad (5.11a)$$

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \lambda_h [[\varrho_h]]^2 dS_h dt \lesssim 1, \quad \int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \lambda_h [[\vartheta_h]]^2 dS_h dt \lesssim 1, \quad \lambda_h \approx |\bar{\mathbf{u}}_h \cdot \mathbf{n}| + \mu_h, \quad (5.11b)$$

which are slightly better than the standard weak BV estimates, cf. [14, 16, 17].

Now, the error term  $E_{1,h}$  for  $b_h$  either equal to 1 or  $\chi(s_h)$  can be handled as

$$\begin{aligned} h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| | [[\varrho_h b_h]] | dS_h &\lesssim h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| ( | [[\varrho_h]] | + | [[\vartheta_h]] | ) dS_h \\ &\lesssim h \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h| dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\bar{\mathbf{u}}_h \cdot \mathbf{n}| ( [[\varrho_h]]^2 + [[\vartheta_h]]^2 ) dS_h \right)^{1/2} \\ &\lesssim \sqrt{h} \|\mathbf{u}_h\|_{L^1(\Omega_h)}^{1/2} F_h^2 \lesssim \sqrt{h} F_h^2, \quad \|F_h^2\|_{L^2(0, T)} \lesssim 1. \end{aligned}$$

Thus it remains to estimate  $E_{1,h}$ ,  $E_{2,h}$ ,  $E_{3,h}$  for  $b_h = u_h^j$ . For  $E_{2,h}$ , we get

$$\begin{aligned} h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\varrho_h u_h^j}| | [[\mathbf{u}_h \cdot \mathbf{n}]] | dS_h &\lesssim h \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h}|^2 dS_h \right)^{1/2} \\ &\lesssim h^{\frac{3}{2} - \frac{\alpha}{2}} F_h^1 h^{-\frac{1}{2}} \|\mathbf{u}_h\|_{L^2(\Omega_h)} \leq h^{1 - \frac{\alpha}{2}} F_h^1, \quad \|F_h^1\|_{L^2(0,T)} \lesssim 1, \end{aligned} \quad (5.12)$$

where we have used the trace inequality, (5.2) and (5.7). As for  $E_{3,h}$  it rewrites as

$$\begin{aligned} h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} | [[\varrho_h u_h^j]] | | [[\mathbf{u}_h]] \cdot \mathbf{n} | dS_h &\lesssim h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h + h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h}| | [[\mathbf{u}_h]] | dS_h \\ &\lesssim h^{2-\alpha} F_h^3 + h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h}| | [[\mathbf{u}_h]] | dS_h, \quad \|F_h^3\|_{L^1(0,T)} \lesssim 1, \end{aligned}$$

while the last integral can be handled exactly as in (5.12). Finally, we are left with  $E_{1,h}$ , specifically,

$$\begin{aligned} h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| | [[\varrho_h u_h^j]] | dS_h \\ \lesssim h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| |\overline{\mathbf{u}_h}| | [[\varrho_h]] | dS_h + h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| | [[\mathbf{u}_h]] | dS_h, \end{aligned}$$

where the last integral can be estimated exactly as in (5.12). Next, by Hölder's inequality, the trace inequality, (5.9), we get

$$\begin{aligned} h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| |\overline{\mathbf{u}_h}| | [[\varrho_h]] | dS_h &\lesssim h \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h}|^3 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| [[\varrho_h]]^2 dS_h \right)^{1/2} \\ &\lesssim \sqrt{h} \|\mathbf{u}_h\|_{L^3(\Omega_h)}^{3/2} F_h^4, \quad \|F_h^4\|_{L^2(0,T)} \lesssim 1. \end{aligned}$$

Now, in view of the interpolation inequality

$$\|\mathbf{u}_h\|_{L^3(\Omega_h)} \lesssim \|\mathbf{u}_h\|_{L^2(\Omega_h)}^{1/2} \|\mathbf{u}_h\|_{L^6(\Omega_h)}^{1/2},$$

combined with (5.7), we obtain

$$h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| |\overline{\mathbf{u}_h}| | [[\varrho_h]] | dS_h \lesssim \sqrt{h} \|\mathbf{u}_h\|_{L^6(\Omega_h)}^{3/4} F_h^4.$$

Finally, we apply the discrete Sobolev embedding (2.5) and (5.2) to conclude

$$h \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| |\overline{\mathbf{u}_h}| | [[\varrho_h]] | dS_h \lesssim \sqrt{h} F_h^4 \left( 1 + \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \frac{[[\mathbf{u}_h]]^2}{h} dS_h \right)^{1/2} \right)^{3/4} \lesssim h^{\frac{4-3\alpha}{8}} F_h^4 F_h^5,$$

with

$$\|F_h^5\|_{L^{8/3}(0,T)} \lesssim 1.$$

Thus the error in the upwind terms satisfies (5.1) as soon as

$$0 < \alpha < \frac{4}{3},$$

and the extra hypotheses (5.3), (5.5) hold.

### 5.1.2 $\mu_h$ -dependent terms

The numerical fluxes of the continuity and momentum equations (2.10), (2.11), and the numerical entropy flux (3.10) contain  $\mu_h$ -dependent terms, namely

$$\begin{aligned} & \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\Pi[\Phi]]] dS_h, \\ & \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \mathbf{u}_h]] \cdot [[\Pi[\Phi]]] dS_h, \\ & \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \left( \overline{\nabla_{\varrho}(\varrho_h \chi(s_h))} [[\varrho_h]] + \overline{\nabla_p(\varrho_h \chi(s_h))} [[p_h]] \right) [[\Pi[\Phi]]] dS_h. \end{aligned}$$

In what follows we show they vanish in the limit  $h \rightarrow 0$ . In view of our hypotheses (5.3), (5.5), the product rule yields

$$[[p_h]] \approx [[\varrho_h]] + [[\vartheta_h]], \quad (5.13)$$

and the estimates (5.11b) imply

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]]^2 dS_h \lesssim 1, \quad (5.14)$$

$$\int_0^T \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\vartheta_h]]^2 dS_h \lesssim 1. \quad (5.15)$$

Assuming the parameter  $\mu_h$  is bounded, Hölder's inequality with (2.1) and (5.14) directly yield

$$\begin{aligned} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]] [[\Pi[\Phi]]] dS_h & \lesssim \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h h^2 dS_h \right)^{1/2} \\ & \lesssim \sqrt{h} F_h^6, \quad \|F_h^6\|_{L^2(0,T)} \lesssim 1. \end{aligned}$$

Analogously, using the product rule, the trace inequality, and bounds (5.2), (5.4), (5.14), we get

$$\begin{aligned}
& \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h \mathbf{u}_h]] \cdot [[\Pi[\Phi]]] dS_h \\
& \lesssim h \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h |\bar{\mathbf{u}}_h|^2 dS_h \right)^{1/2} + \sqrt{h} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \\
& \lesssim \sqrt{h} \|\mathbf{u}_h\|_{L^2(\Omega_h)} F_h^6 + h^{1-\frac{\alpha}{2}} F_h^1 \lesssim \sqrt{h} F_h^6 + h^{1-\frac{\alpha}{2}} F_h^1, \quad \|F_h^1\|_{L^2(0,T)}, \|F_h^6\|_{L^2(0,T)} \lesssim 1.
\end{aligned}$$

Finally, (5.13), (5.14) and (5.15) imply

$$\begin{aligned}
& \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h \left( \overline{\nabla_{\varrho}(\varrho_h \chi(s_h))} [[\varrho_h]] + \overline{\nabla_p(\varrho_h \chi(s_h))} [[p_h]] \right) [[\Pi[\Phi]]] dS_h \\
& \lesssim \sqrt{h} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\varrho_h]]^2 dS_h \right)^{1/2} + \sqrt{h} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \mu_h [[\vartheta_h]]^2 dS_h \right)^{1/2} \\
& \lesssim \sqrt{h} (F_h^6 + F_h^7), \quad \|F_h^6\|_{L^2(0,T)}, \|F_h^7\|_{L^2(0,T)} \lesssim 1.
\end{aligned}$$

## 5.2 The artificial viscosity and the pressure terms

There are two remaining terms to be handled in the momentum equation, namely,

$$h^{\alpha-1} \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\Pi[\Phi]]] dS_h,$$

and

$$\sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \bar{p}_h \mathbf{n} \cdot [[\Pi[\Phi]]] dS_h.$$

First, in accordance with (2.1),

$$\begin{aligned}
& h^{\alpha-1} \left| \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]] \cdot [[\Pi[\Phi]]] dS_h \right| \lesssim h^{\alpha-1} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} h^2 dS_h \right)^{1/2} \\
& \lesssim h^{\alpha-\frac{1}{2}} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[\mathbf{u}_h]]^2 dS_h \right)^{1/2} \lesssim h^{\alpha/2} F_h^7, \quad \|F_h^7\|_{L^2(0,T)} \lesssim 1.
\end{aligned}$$

Second,

$$\begin{aligned}
\int_{\Omega_h} p_h \operatorname{div}_x \Phi \, dx &= \sum_{K \in \mathcal{T}_h} \int_K p_h \operatorname{div}_x \Phi \, dx = \sum_{K \in \partial K_h} \int_{\partial K} p_h \Phi \cdot \mathbf{n} dS_h \\
&= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] (\Phi - \overline{\Pi[\Phi]}) \cdot \mathbf{n} dS_h + \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] \overline{\Pi[\Phi]} \cdot \mathbf{n} dS_h \\
&= \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] (\Phi - \overline{\Pi[\Phi]}) \cdot \mathbf{n} dS_h - \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} \overline{p_h} [[\Pi[\Phi]]] \cdot \mathbf{n} dS_h.
\end{aligned}$$

Here, similarly to the preceding section, the error term can be estimated as

$$\begin{aligned}
\left| \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] (\Phi - \overline{\Pi[\Phi]}) \cdot \mathbf{n} dS_h \right| &\lesssim \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]]^2 dS_h \right)^{1/2} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} h^2 dS_h \right)^{1/2} \\
&\lesssim \sqrt{h} \left( \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]]^2 dS_h \right)^{1/2}.
\end{aligned}$$

Recall that  $[[p_h]] \approx [[\varrho_h]] + [[\vartheta_h]]$ ; whence for the error to tend to zero it is enough to *assume*

$$\mu_h \gtrsim h^\beta > 0, \quad 0 \leq \beta < 1.$$

In the case of uniform rectangular/cubic elements we allow  $\mu_h = 0$ . Indeed, due to (2.2) and (2.4) we have, for any  $\Phi \in C^2(\overline{\Omega}_h; R^N)$ ,

$$\begin{aligned}
\left| \sum_{\sigma \in \Sigma_{int}} \int_{\sigma} [[p_h]] (\Phi - \overline{\Pi[\Phi]}) \cdot \mathbf{n} dS_h \right| &\lesssim \sum_{\sigma \in \Sigma_{int}} \left| [[p_h]] \right| \int_{\sigma} |\Phi - \overline{\Pi[\Phi]}| dS_h \\
&\lesssim \sum_{\sigma \in \Sigma_{int}} \left( \int_{\sigma} |[[p_h]]| dS_h \right) h^2 \lesssim h \sum_{K \in \mathcal{T}_h} \int_K |p_h| dx,
\end{aligned}$$

which tends to zero as  $p_h \in L^\infty((0, T); L^1(\Omega_h))$ .

### 5.3 Consistency formulation

Summing up the results of Subsections 5.1 and 5.2, we obtain a consistency formulation of the approximation scheme (2.10–2.12).

**Theorem 5.1.** *Let the initial data  $\varrho_{0,h}$ ,  $\mathbf{m}_{0,h}$ ,  $E_{0,h}$  satisfy the hypotheses of Theorem 4.2. Let  $[\varrho_h, \mathbf{m}_h, E_h]$  be the unique solutions of the approximate problem (2.10–2.12) on the time interval  $[0, T]$ .*



Then

$$\left[ \int_{\Omega_h} \varrho_h \varphi \, dx \right]_{t=0}^{t=\tau} = \int_0^\tau \int_{\Omega_h} [\varrho_h \partial_t \varphi + \mathbf{m}_h \cdot \nabla_x \varphi] \, dx \, dt + \int_0^\tau e_{1,h}(t, \varphi) \, dt \quad (5.16)$$

for any  $\varphi \in C^1([0, T] \times \overline{\Omega}_h)$ ;

$$\left[ \int_{\Omega_h} \mathbf{m}_h \varphi \, dx \right]_{t=0}^{t=\tau} = \int_0^\tau \int_{\Omega_h} \left[ \mathbf{m}_h \cdot \partial_t \varphi + \frac{\mathbf{m}_h \otimes \mathbf{m}_h}{\varrho_h} : \nabla_x \varphi + p_h \operatorname{div}_x \varphi \right] \, dx \, dt + \int_0^\tau e_{2,h}(t, \varphi) \, dt \quad (5.17)$$

for any  $\varphi \in C^1([0, T] \times \overline{\Omega}_h; R^N)$ ,  $\varphi \cdot \mathbf{n}|_{\Omega_h} = 0$ ;

$$\int_{\Omega_h} E_h(t) \, dx = \int_{\Omega_h} E_{0,h} \, dx; \quad (5.18)$$

$$\left[ \int_{\Omega_h} \varrho_h \chi(s_h) \varphi \, dx \right]_{t=0}^{t=\tau} \geq \int_0^\tau \int_{\Omega_h} [\varrho_h \chi(s_h) \partial_t \varphi + \chi(s_h) \mathbf{m}_h \cdot \nabla_x \varphi] \, dx \, dt + \int_0^\tau e_{3,h}(t, \varphi) \, dt \quad (5.19)$$

for any  $\varphi \in C^1([0, T] \times \overline{\Omega}_h)$ ,  $\varphi \geq 0$ , and any  $\chi$ ,

$\chi : R \rightarrow R$  a non-decreasing concave function,  $\chi(s) \leq \bar{\chi}$  for all  $s \in R$ .

If, in addition,

$$h^\beta \lesssim \mu_h \lesssim 1, \quad 0 \leq \beta < 1, \quad 0 < \alpha < \frac{4}{3}, \quad (5.20)$$

and

$$0 < \bar{\varrho} \leq \varrho_h(t), \quad \vartheta_h(t) \leq \bar{\vartheta} \text{ for all } t \in [0, T] \text{ uniformly for } h \rightarrow 0, \quad (5.21)$$

then

$$\|e_{j,h}(\cdot, \varphi)\|_{L^1(0,T)} \lesssim h^\delta \|\varphi\|_{C^1} \text{ for some } \delta > 0.$$

In the case of uniform rectangular/cubic elements the result of Theorem 5.1 remains valid for  $0 \leq \mu_h \lesssim 1$ , and  $\varphi \in C^1([0, T]; C^2(\overline{\Omega}_h; R^N))$ ,  $\varphi \cdot \mathbf{n}|_{\Omega_h} = 0$ .

**Remark 5.2.** Omitting the  $h^{\alpha-1}$ -dependent terms in (2.11–2.12) corresponds to the Lax–Friedrichs scheme with the numerical fluxes

$$\overline{r_h \mathbf{u}_h} \cdot \mathbf{n} - \lambda_h[[r_h]] = \overline{r_h \mathbf{u}_h} \cdot \mathbf{n} - \lambda_h[[r_h]] + \frac{1}{4}[[r_h]][[\mathbf{u}_h]] \cdot \mathbf{n}$$

with  $\lambda_h = \frac{1}{2} \max(\lambda_h^{\text{in}}, \lambda_h^{\text{out}})$ ,  $\lambda_h = \frac{1}{2} |\overline{\mathbf{u}_h} \cdot \mathbf{n}| + c_h$ , where  $c_h = \sqrt{\gamma \bar{\vartheta}_h}$  stands for the speed of sound. In the standard Lax–Friedrichs scheme the average  $\overline{r_h \mathbf{u}_h}$  instead of  $\overline{r_h \mathbf{u}_h}$  is used. Moreover, in the energy equation  $\overline{p_h \mathbf{u}_h}$  is used instead of (2.14) for the pressure term in the energy flux, cf. Remark 2.2. Nevertheless, the present proof under the hypotheses (5.20), (5.21) might be adapted to the standard Lax–Friedrichs scheme.

## 6 Convergence

In view of the uniform bounds (4.2), (4.5), and (4.6), the family  $\{\varrho_h, \mathbf{m}_h, E_h\}_{h>0}$  of approximate solutions is uniformly bounded in  $L^\infty(0, T; L^1(\Omega_h))$ . Moreover,  $\{\varrho_h\}_{h>0}$  is bounded in  $L^\infty(0, T; L^\gamma(\Omega_h))$  and  $\{\mathbf{m}_h\}_{h>0}$  is bounded in  $L^\infty(0, T; L^{\frac{2\gamma}{\gamma+1}}(\Omega_h; R^N))$  uniformly for  $h \rightarrow 0$ .

### 6.1 Young measure generated by the approximate solutions

In accordance with the fundamental theorem on Young measures, see Ball [1] or Pedregal [22], the family  $\{\varrho_h, \mathbf{m}_h, E_h\}_{h>0}$ , up to a suitable subsequence, generates a Young measure  $\{\mathcal{V}_{t,x}\}_{(t,x) \in (0,T) \times \Omega_h}$ . Recall that the Young measure is an object with the following properties:

- the mapping

$$\mathcal{V}_{t,x} : (t, x) \in (0, T) \times \Omega_h \mapsto \mathcal{P}(\mathcal{F})$$

is weakly-(\*) measurable, where  $\mathcal{P}$  is the space of probability measures defined on the phase space

$$\mathcal{F} = \left\{ \varrho, \mathbf{m}, E \mid \varrho \geq 0, \mathbf{m} \in R^N, E \geq 0 \right\};$$

- 

$$G(\varrho_h, \mathbf{m}_h, E_h) \rightarrow \overline{G(\varrho, \mathbf{m}, E)} \text{ weakly-(*) in } L^\infty((0, T) \times \Omega_h)$$

for any  $G \in C_c(\mathcal{F})$ , and

$$\overline{G(\varrho, \mathbf{m}, E)}(t, x) = \int_{\mathcal{F}} G(\varrho, \mathbf{m}, E) d\mathcal{V}_{t,x} \equiv \langle \mathcal{V}_{t,x}; G(\varrho, \mathbf{m}, E) \rangle \text{ for a.a. } (t, x) \in (0, T) \times \Omega_h.$$

We shall use the following result proved in [11, Lemma 2.1].

**Lemma 6.1.** *Let*

$$|G(\varrho, \mathbf{m}, E)| \leq F(\varrho, \mathbf{m}, E) \text{ for all } (\varrho, \mathbf{m}, E) \in \mathcal{F}.$$

*Then*

$$\left| \overline{G(\varrho, \mathbf{m}, E)} - \langle \mathcal{V}_{t,x}; G(\varrho, \mathbf{m}, E) \rangle \right| \leq \overline{F(\varrho, \mathbf{m}, E)} - \langle \mathcal{V}_{t,x}; F(\varrho, \mathbf{m}, E) \rangle \equiv \mu_F \text{ in } \mathcal{M}([0, T] \times \Omega).$$

### 6.2 Kinetic energy concentration defect

Under the extra hypotheses (5.21), the support of the measure  $\mathcal{V}_{t,x}$  is contained in the set

$$\text{supp}[\mathcal{V}_{t,x}] \subset \left\{ [\varrho, \mathbf{m}, E] \mid 0 < \underline{\varrho} \leq \varrho \leq \bar{\varrho}, 0 < \underline{\vartheta} \leq \vartheta \leq \bar{\vartheta} \right\} \text{ for a.a. } (t, x) \in (0, T) \times \Omega_h.$$

In particular, all non-linearities appearing in the consistency formulation (5.16 – 5.19) are weakly precompact in the Lebesgue space  $L^1((0, T) \times \Omega_h)$ , with the only exception of the convective term

$$\left\{ \frac{\mathbf{m}_h \otimes \mathbf{m}_h}{\varrho_h} \right\}_{h>0} \text{ bounded in } L^1((0, T) \times \Omega_h, R^{N \times N}).$$

For the latter we can only assert that

$$\frac{\mathbf{m}_h \otimes \mathbf{m}_h}{\varrho_h} \rightarrow \frac{\overline{\mathbf{m} \otimes \mathbf{m}}}{\varrho} \text{ weakly-} (*) \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}_h; R^{N \times N}).$$

We denote

$$\mathbb{C}_d = \frac{\overline{\mathbf{m} \otimes \mathbf{m}}}{\varrho} - \left\langle \mathcal{V}_{t,x}; \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right\rangle \in \mathcal{M}([0, T] \times \overline{\Omega}_h; R^{N \times N})$$

the associated *concentration defect measure*. As

$$\left| \frac{\mathbf{m}_h \otimes \mathbf{m}_h}{\varrho_h} \right| \lesssim \frac{|\mathbf{m}_h|^2}{\varrho_h} \leq E_h,$$

we may use Lemma 6.1 to conclude that

$$\int_0^\tau \int_{\overline{\Omega}_h} 1 \, d|\mathbb{C}_d| \lesssim \int_{\Omega_h} E_0 \, dx - \int_{\Omega_h} \langle \mathcal{V}_{\tau,x}; E \rangle \, dx \text{ for a.a. } \tau \in [0, T]. \quad (6.1)$$

The quantity on the right-hand side of (6.1) is called *energy dissipation defect* and inequality (6.1) plays a crucial role in the concept of *dissipative measure-valued (DMV) solutions* to the complete Euler system introduced in [6].

### 6.3 Limit problem

We say that a family of probability measures  $\{\mathcal{V}_{t,x}\}_{(t,x) \in (0,T) \times \Omega_h}$  is a (DMV) solution to the complete Euler system (1.1–1.3) if:

•

$$\left[ \int_{\Omega_h} \langle \mathcal{V}_{t,x}; \varrho \rangle \varphi \, dx \right]_{t=0}^{t=\tau} = \int_0^\tau \int_{\Omega_h} [\langle \mathcal{V}_{t,x}; \varrho \rangle \partial_t \varphi + \langle \mathcal{V}_{t,x}; \mathbf{m} \rangle \cdot \nabla_x \varphi] \, dx \, dt$$

for any  $\varphi \in C^1([0, T] \times \overline{\Omega}_h)$ ;

•

$$\begin{aligned} & \left[ \int_{\Omega_h} \langle \mathcal{V}_{t,x}; \mathbf{m} \rangle \cdot \varphi \, dx \right]_{t=0}^{t=\tau} \\ &= \int_0^\tau \int_{\Omega_h} \left[ \langle \mathcal{V}_{t,x}; \mathbf{m} \rangle \cdot \partial_t \varphi + \left\langle \mathcal{V}_{t,x}; \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right\rangle : \nabla_x \varphi + \langle \mathcal{V}_{t,x}; p \rangle \operatorname{div}_x \varphi \right] \, dx \, dt \\ &+ \int_0^\tau \int_{\overline{\Omega}_h} \nabla_x \varphi : d\mathbb{C}_d \end{aligned}$$

for any  $\varphi \in C^1([0, T] \times \overline{\Omega}_h; R^N)$ ,  $\varphi \cdot \mathbf{n}|_{\Omega_h} = 0$ ;

•

$$\int_{\Omega_h} \langle \mathcal{V}_{\tau,x}; E \rangle \, dx \leq \int_{\Omega_h} E_0 \, dx$$

for a.a.  $\tau \in [0, T]$ ;

•

$$\left[ \int_{\Omega_h} \langle \mathcal{V}_{t,x}; \varrho \chi(s) \rangle \varphi \, dx \right]_{t=0}^{t=\tau} \geq \int_0^\tau \int_{\Omega_h} [\langle \mathcal{V}_{t,x}; \varrho \chi(s) \rangle \partial_t \varphi + \langle \mathcal{V}_{t,x}; \chi(s) \mathbf{m} \rangle \cdot \nabla_x \varphi] \, dx \, dt$$

for any  $\varphi \in C^1([0, T] \times \bar{\Omega}_h)$ ,  $\varphi \geq 0$ , and any  $\chi$ ,

$\chi : R \rightarrow R$  a non-decreasing concave function,  $\chi(s) \leq \bar{\chi}$  for all  $s \in R$ ;

•

$$\int_0^\tau \int_{\bar{\Omega}_h} 1 \, d|\mathbb{C}_d| \lesssim \int_{\Omega_h} E_0 \, dx - \int_{\Omega_h} \langle \mathcal{V}_{\tau,x}; E \rangle \, dx$$

for a.a.  $\tau \in [0, T]$ .

Summing up the preceding discussion, we can state the following result.

**Theorem 6.2.** *Let the initial data  $\varrho_{0,h}$ ,  $\mathbf{m}_{0,h}$ ,  $E_{0,h}$  satisfy*

$$\varrho_{0,h} \geq \underline{\varrho} > 0, \quad E_{0,h} - \frac{1}{2} \frac{|\mathbf{m}_{0,h}|^2}{\varrho_{0,h}} > 0.$$

*Let  $[\varrho_h, \mathbf{m}_h, E_h]$  be the solution of the scheme (2.10–2.12) such that*

$$h^\beta \lesssim \mu_h \lesssim 1, \quad 0 \leq \beta < 1, \quad 0 < \alpha < \frac{4}{3},$$

*and*

$$0 < \bar{\varrho} \leq \varrho_h(t), \quad \vartheta_h(t) \leq \bar{\vartheta} \text{ for all } t \in [0, T] \text{ uniformly for } h \rightarrow 0.$$

*Then the family of approximate solutions  $\{\varrho_h, \mathbf{m}_h, E_h\}_{h>0}$  generates a Young measure  $\{\mathcal{V}_{t,x}\}_{(t,x) \in (0,T) \times \Omega_h}$  that is a (DMV) solution of the complete Euler system (1.1–1.3).*

Finally, evoking the weak (DMV)–strong uniqueness result proved in [6, Theorem 3.3] we conclude with the following corollary.

**Corollary 6.3.** *In addition to the hypotheses of Theorem 6.2, suppose that the complete Euler system (1.1–1.3) admits a Lipschitz–continuous solution  $[\varrho, \mathbf{m}, E]$  defined on  $[0, T]$ .*

*Then*

$$\varrho_h \rightarrow \varrho, \quad \mathbf{m}_h \rightarrow \mathbf{m}, \quad E_h \rightarrow E \text{ (strongly) in } L^1((0, T) \times \Omega_h).$$

## Conclusion

In the present paper we have studied the convergence of a new finite volume method for multi-dimensional Euler equations of gas dynamics. As the Euler system admits highly oscillatory solutions, in particular they are ill-posed in the class of weak entropy solutions for  $L^\infty$ -initial data [13], it is more natural to investigate the convergence in the class of *dissipative measure-valued (DMV) solutions*. The (DMV) solutions represent the most general class of solutions that still satisfy the weak-strong uniqueness property. Thus, if the strong solution exists the (DMV) solution coincides with the strong one on its lifespan, cf. [6].

Our study is inspired by the work of Guermond and Popov [19] who proposed a viscous regularization of the compressible Euler equations satisfying the minimum entropy principle and positivity preserving properties. They also showed the connection to the two-velocities Brenner's model [3, 4, 5], which is a base of our new finite volume method (2.10–2.12). The method is (i) positivity preserving, i.e. discrete density, pressure and temperature are positive on any finite time interval, (ii) entropy stable and (iii) satisfies the minimum entropy principle. Moreover, the discrete entropy inequality allows us to control certain weak BV-norms, cf. (5.11). These results together with a priori estimates (4.1–4.9) yield the consistency of the new finite volume method under mild hypothesis. Indeed, instead of conventional convergence results based on rather unrealistic hypothesis on uniform boundedness of all physical quantities, we only require that the discrete temperature is bounded and vacuum does not appear, cf. (5.21). In Theorem 6.1 we have shown that the numerical solutions of the finite volume method (2.10–2.12) generate the (DMV) solution of the Euler equations. Consequently, using the recent result on the (DMV)-strong uniqueness, we have proven the convergence to the strong solution on its lifespan.

It seems that the hypothesis on  $\varrho_h$  can be relaxed, though removing the boundedness of  $\vartheta_h$  remains open. This can be an interesting question for future study. Moreover, in order to preserve the Galilean invariance of the Brenner model (1.8–1.10) it is possible to consider the symmetric gradient in the  $h^\alpha$ -diffusion terms and the same convergence result can be shown. As far as we know the present convergence result is the first result in the literature, where the convergence of a finite volume method has been proven for multi-dimensional Euler equations assuming only that the gas remains in its non-degenerate region.

## References

- [1] J.M. Ball. A version of the fundamental theorem for Young measures. *In Lect. Notes in Physics 344, Springer-Verlag*, pages 207–215, 1989.
- [2] A. Bardow and H. C. Öttinger. Consequences of the Brenner modification to the Navier–Stokes equations for dynamic light scattering. *Phys. A* **373**: 88–96, 2007.
- [3] H. Brenner. Kinematics of volume transport. *Phys. A*, **349**: 11–59, 2005.

- [4] H. Brenner. Navier-Stokes revisited. *Phys. A*, **349**(1-2): 60–132, 2005.
- [5] H. Brenner. Fluid mechanics revisited. *Phys. A*, **349**:190–224, 2006.
- [6] J. Březina and E. Feireisl. Measure-valued solutions to the complete Euler system. 2017. arxiv preprint No. 1702.04878, to appear in *J. Math. Soc. Japan*.
- [7] J. Březina and E. Feireisl. Measure-valued solutions to the complete Euler system revisited. 2017. arxiv preprint No. 1710.10751, to appear in *Z. Angew. Math. Mech*.
- [8] C. Chainais–Hillairet and J. Droniou. Finite volume schemes for non-coercive elliptic problems with Neumann boundary conditions. *IMA J. Numer. Anal.* **31**(1): 61–85, 2011.
- [9] P. G. Ciarlet. The Finite Element Method for Elliptic Problems. *Classics in Applied Mathematics, Society for Industrial and Applied Mathematics*, 2002.
- [10] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. *Handbook of numerical analysis* **7**: 713–1018, 2000.
- [11] E. Feireisl, P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann. Dissipative measure-valued solutions to the compressible Navier–Stokes system. *Calc. Var. Partial Diff.* **55**(6): 55–141, 2016.
- [12] E. Feireisl, T. Karper, and A. Novotný. A convergent numerical method for the Navier–Stokes–Fourier system. *IMA J. Numer. Anal.* **36**(4): 1477–1535, 2016.
- [13] E. Feireisl, C. Klingenberg, O. Kreml, and S. Markfelder. On oscillatory solutions to the complete Euler system. 2017. arxiv preprint No. 1710.10918.
- [14] E. Feireisl, M. Lukáčová-Medvid’ová, and H. Mizerová. Convergence of finite volume schemes for the Euler equations via dissipative measure-valued solutions. submitted, 2018.
- [15] U. K. Fjordholm, R. Käppeli, S. Mishra, and E. Tadmor. Construction of approximate entropy measure valued solutions for hyperbolic systems of conservation laws. *Found. Comp. Math.*, pages 1–65, 2015.
- [16] U. S. Fjordholm, S. Mishra, and E. Tadmor. Arbitrarily high-order accurate entropy stable essentially non-oscillatory schemes for systems of conservation laws. *SIAM J. Numer. Anal.* **50**(2): 544–573, 2012.
- [17] U. S. Fjordholm, S. Mishra, and E. Tadmor. On the computation of measure-valued solutions. *Acta Numer.* **25**: 567–679, 2016.
- [18] C. J. Greenshields and J. M. Reese. The structure of shock waves as a test of Brenner’s modifications to the Navier–Stokes equations. *J. Fluid Mech.* **580**: 407–429, 2007.

- [19] J. L. Guermond and B. Popov. Viscous regularization of the Euler equations and entropy principles. *SIAM J. Appl. Math.* **74**(2): 284–305, 2014.
- [20] Z. Guo and K. Xu. Numerical Validation of Brenner’s Hydrodynamic Model by Force Driven Poiseuille Flow. *Adv. Appl. Math. Mech.* **1**(3): 391–401, 2009.
- [21] H. C. Öttinger, H. Struchtrup, and M. Liu. Inconsistency of a dissipative contribution to the mass flux in hydrodynamics. *Phys. Rev. E* **80**(056303): 1–8, 2009.
- [22] P. Pedregal. *Parametrized measures and variational principles*. Birkhäuser, Basel, 1997.
- [23] R. Rannacher and S. Turek. Simple nonconforming quadrilateral Stokes element. *Numer. Meth. Par. D. E.* **8**(2): 97–111, 1992.
- [24] E. Tadmor. Entropy stability theory for difference approximations of nonlinear conservation laws and related time dependent problems. *Acta Numer.* **12**: 451–512, 2003.
- [25] E. Tadmor. The numerical viscosity of entropy stable schemes for systems of conservation laws. *Math. Comp.* **49**(179): 91–103, 1987.
- [26] E. Tadmor. Minimum entropy principle in the gas dynamic equations. *Appl. Num. Math.* **2**: 211–219, 1986.