



INSTITUTE OF MATHEMATICS

THE CZECH ACADEMY OF SCIENCES

**Electromagnetic fields with vanishing
scalar invariants**

Marcello Ortaggio

Vojtěch Pravda

Preprint No. 13-2016

PRAHA 2016

Electromagnetic fields with vanishing scalar invariants

Marcello Ortaggio* and Vojtěch Pravda†
Institute of Mathematics of the Czech Academy of Sciences
Žitná 25, 115 67 Prague 1, Czech Republic

November 13, 2015

Abstract

We determine the class of p -forms \mathbf{F} which possess vanishing scalar invariants (VSI) at arbitrary order in a n -dimensional spacetime. Namely, we prove that \mathbf{F} is VSI *if and only if* it is of type N, its multiple null direction ℓ is “degenerate Kundt”, and $\mathcal{L}_\ell \mathbf{F} = 0$. The result is theory-independent. Next, we discuss the special case of Maxwell fields, both at the level of test fields and of the full Einstein-Maxwell equations. These describe electromagnetic non-expanding waves propagating in various Kundt spacetimes. We further point out that a subset of these solutions possesses a *universal* property, i.e., they also solve (virtually) any generalized (non-linear and with higher derivatives) electrodynamics, possibly also coupled to Einstein’s gravity.

PACS 04.50.+h, 04.20.Jb, 04.40.Nr

Contents

1	Introduction	2
1.1	Background	2
1.2	Main result	4
2	Explicit form of VSI electromagnetic fields and adapted coordinates	5
2.1	Electromagnetic field and spacetime metric	5
2.2	Maxwell’s equations	7
2.2.1	The special case $p = 1$	7
2.2.2	The special cases $n = 4, 3$	7
2.3	Maxwell-Chern-Simons’ equations	7
2.3.1	Generic case $k \geq 2$	8
2.3.2	Special case $k = 1$	8
2.4	Universal solutions of generalized electrodynamics (test fields)	8
3	Einstein-Maxwell solutions	9
3.1	General equations	9
3.1.1	VSI spacetimes with VSI Maxwell fields	11
3.1.2	pp -waves with VSI Maxwell fields	11
3.2	Universal Einstein-Maxwell solutions	12

*ortaggio@math.cas.cz

†pravda@math.cas.cz

A	Some of the Newman-Penrose equations for Kundt spacetimes	13
A.1	General Kundt spacetimes	13
A.2	Kundt spacetimes of aligned Riemann type II	14
B	Proof of Theorem 1.5	15
B.1	Proof of “2. \Rightarrow 1.”	15
B.1.1	Preliminaries	15
B.1.2	Proof	16
B.2	Proof of “1. \Rightarrow 2.”	16
B.2.1	Preliminaries	16
B.2.2	Proof	18
C	VSI₁ and VSI₂ p-forms	19

1 Introduction

1.1 Background

Synge [1] called electromagnetic *null* fields those characterized by the vanishing of the two Lorentz invariants, i.e.,

$$F_{ab}F^{ab} = 0, \quad F_{ab}{}^*F^{ab} = 0. \quad (1)$$

From a physical viewpoint, null Maxwell fields characterize electromagnetic plane waves [1] as well as the asymptotic behaviour of radiative systems (cf. [2] and references therein). Fields satisfying (1) single out a unique null direction ℓ such that the corresponding energy-momentum tensor can be written as $T_{ab} = A\ell_a\ell_b$ [3] (cf. also, e.g., [1, 4]). They also possess the unique property that their field strength at any spacetime point can be made as small (or large) as desired in a suitably boosted reference frame [1, 5]. Moreover, null solutions to the sourcefree Maxwell equations can be associated with shearfree congruences of null geodesics (and viceversa) via the Mariot-Robinson theorem [2, 4] and are therefore geometrically privileged.

There are further reasons that motivate the interest in fields with the property (1). On the gravity side, an analog of null electromagnetic fields is given by metrics of Riemann type III and N, for which all the zeroth-order scalar invariants constructed from the Riemann tensor vanish identically (see [4] in 4D and [6] in higher dimensions).¹ These have remarkable properties. For example, all type N Einstein spacetimes are automatically vacuum solutions of quadratic [8] (in particular, Gauss-Bonnet [9]) and Lovelock gravity [10]. Furthermore, it has been known for some time [11–13] that certain type N pp -wave solutions in general relativity (in vacuum or with dilaton and form fields) are classical solutions to string theory to all orders in σ -model perturbation theory – in fact, they are also solutions to *any* gravity theory in which the “corrections” to the field equations can be expressed in terms of scalars and tensors constructed from the field strengths and their covariant derivatives (see [11, 13, 14] for details), and are in this sense *universal* [15].² Apart from being of Riemann type N and universal, the metrics considered in [11–13] have the special property that all

¹Hereafter, by “zeroth-order” invariants of a certain tensor we refer to the *algebraic* ones, i.e., those not involving covariant derivatives of the given tensor. Additionally, we will restrict ourselves to *polynomial* scalar invariants (cf. Definition 5.1 of [7]).

²To be precise, in the terminology of [15] the universal property refers only to certain Einstein metrics, whereas here we clearly use the term “universal” in a broader sense (which applies not only to the metric – not necessarily Einstein – but to the full solution, also including possible matter fields).

the scalar invariants constructed from the Riemann tensor *and* its covariant derivatives vanish and thus belong to the VSI class of spacetimes [6, 16] (cf. Definition 1.2 below).³

In view of these results for gravity, one may wonder whether certain null (or VSI) Maxwell fields possess a similar “universal” property and thus solve also generalized theories of electrodynamics. In fact, it was already known to Schrödinger [5, 19] that all null Maxwell fields solve the equations for the electromagnetic field in any non-linear electrodynamics (NLE; cf., e.g., [20]) – this was later extended to the full Einstein-Maxwell equations including the electromagnetic backreaction on the spacetime geometry [21–23]. However, the case of theories more general than NLE (including also derivatives of the field strength in the Lagrangian, cf., e.g., [24, 25]) seems not to have been investigated systematically from this viewpoint. As a first step in this direction, it is the purpose of the present paper to determine the class of electromagnetic fields for which all the scalar invariants constructed from the field strength and its derivatives vanish identically (VSI), which is obviously a subset of null fields. We will also point out a few examples possessing the universal property, while a more detailed study of the latter will be presented elsewhere.

Various extensions of Einstein-Maxwell gravity exist in which the number of spacetime dimensions n may be greater than four, and the electromagnetic field is represented by a rank- p form (cf., e.g., [26, 27]). These theories have attracted interest in recent years, motivated, in particular, by supergravity and string theory. We will therefore consider null fields with arbitrary $n \geq 3$ and p (with $1 \leq p \leq n - 1$ to avoid trivial cases). The relevant generalization of the concept of null fields is straightforward: all the zeroth-order scalar polynomial invariant constructed out of a p -form \mathbf{F} vanish (thus generalizing (1)) iff [28] \mathbf{F} is of type N in the null alignment classification of [29] (cf. Corollary 1.4 below; for the case $p = 2$ this was proven earlier in [6]). For this reason, in this paper we shall use the terminology “null” and “type N” interchangeably, when referring to p -forms.

It will be also useful to observe that the type N condition of [29] for p -forms can be easily rephrased in a manifestly frame-independent way, which we thus adopt as a definition here (see also [30] when $p = 2$):

Definition 1.1 (p -forms of type N). *At a spacetime point, a p -form \mathbf{F} is of type N if it satisfies*

$$\ell^a F_{ab_1 \dots b_{p-1}} = 0, \quad \ell_{[a} F_{b_1 \dots b_p]} = 0, \quad (2)$$

where ℓ is a null vector (this follows from (2) and need not be assumed). The second condition can be equivalently replaced by $\ell^a {}^* F_{ab_1 \dots b_{n-p-1}} = 0$ (cf. [1, 31] for $n = 4$, $p = 2$).

The most general algebraic form of a null p -form \mathbf{F} is thus known (eq. (6) below). We will determine what are the necessary and sufficient conditions for a p -form \mathbf{F} living in a certain spacetime to be VSI, in the sense of the following definition:

Definition 1.2 (VSI tensors). *A tensor in a spacetime with metric g_{ab} is VSI_I if the scalar polynomial invariants constructed from the tensor itself and its covariant derivatives up to order I ($I = 0, 1, 2, 3, \dots$) vanish. It is VSI if all its scalar polynomial invariants of arbitrary order vanish. As in [6, 32], if the Riemann tensor of g_{ab} is VSI (or VSI_I), the spacetime itself is said to be VSI (or VSI_I).*

For the purposes of the present paper, it will be convenient to recall Corollary 3.2 of [28], which can be expressed as

Theorem 1.3 (Algebraic VSI theorem [28]). *A tensor is VSI_0 iff it is of type III (or more special).*

³In the vacuum case, recent analysis [17, 18] has extended the result of [11–13] in various directions. In particular, it is now clear that the VSI property is neither a sufficient nor a necessary condition for “universality” (however, universal spacetimes must be CSI [17], i.e., with constant scalar invariants). Moreover, certain non- pp -wave spacetimes of Weyl type III, N [17] and II (or D) [18] can also be universal.

Again, the tensor type refers to the algebraic classification of [29] (see also the review [7]). Now, recalling that a non-zero p -form \mathbf{F} can only be of type G, II (D) or N (cf. [6, 33, 34]),⁴ it immediately follows from Theorem 1.3 that

Corollary 1.4 (VSI₀ p -forms). *A p -form \mathbf{F} is VSI₀ iff it is of type N.*

1.2 Main result

The main result of this paper (proven in appendix B) is the following

Theorem 1.5 (VSI p -forms). *The following two conditions are equivalent:*

1. *a non-zero p -form field \mathbf{F} is VSI in a spacetime with metric g_{ab}*
2. *(a) \mathbf{F} possesses a multiple null direction ℓ , i.e., it is of type N*
(b) $\mathcal{L}_\ell \mathbf{F} = 0$
(c) g_{ab} is a degenerate Kundt metric, and ℓ is the corresponding Kundt null direction.

We can already make a few observations about the implications of Theorem 1.5.

Remark 1.6 (Degenerate Kundt metrics). *The definition of degenerate Kundt metrics [37, 38] is reproduced in Appendix A.2 (Definition A.1). By Proposition A.2 (with Propositions 7.1 and 7.3 of [7]), it follows that, e.g., all VSI spacetimes, all pp -waves, and all Kundt Einstein (in particular, Minkowski and (A)dS) or aligned pure radiation spacetimes necessarily belong to this class. As a consequence, when considering VSI p -forms coupled to gravity, the “degenerate” part of condition 2c above becomes automatically trivially true (since a null p -form gives rise to an aligned pure radiation term in the energy-momentum tensor, cf. section 3). It is also worth remarking that, in four dimensions, the degenerate Kundt spacetimes are the only metrics not determined by their curvature scalar invariants [37], and they are thus of particular relevance for the equivalence problem [37, 38].*

Remark 1.7 (VSI vector field). *Condition 2c of Theorem 1.5 implies that any affinely parametrized principal null vector ℓ of a VSI \mathbf{F} is itself VSI. Indeed, in the special case $p = 1$, thanks to condition 2c, condition 2a is trivially satisfied and condition 2b simply means that ℓ is affinely parametrized; i.e., for $p = 1$, Theorem 1.5 reduces to: a vector field ℓ is VSI in a spacetime with metric g_{ab} iff ℓ is Kundt and affinely parameterized, and g_{ab} is a degenerate Kundt metric w.r.t ℓ .*

Remark 1.8 (Theory independence of the result). *In Theorem 1.5, the p -form \mathbf{F} is not assumed to satisfy any particular field equations and the result is thus rather general. On the other hand, if \mathbf{F} is taken to be closed (i.e., $d\mathbf{F} = 0$) then condition 2b automatically follows from the type N condition 2a, and need not be assumed. Otherwise, condition 2b is needed to ensure (together with 2a and 2c) that $\nabla \mathbf{F}$ is of type III (or more special). Note that $\mathcal{L}_\ell \mathbf{F} = \nabla_\ell \mathbf{F}$ if \mathbf{F} is of type N and ℓ is Kundt.*

Remark 1.9 (VSI₃ \Rightarrow VSI for a p -form). *From the proof of Theorem 1.5 (Appendix B) it follows, in fact, that if \mathbf{F} is VSI₃ then it is necessarily VSI. Condition 1 could thus be accordingly relaxed in the proof “1. \Rightarrow 2.”. By contrast, recall that in the case of the Riemann tensor one has VSI₂ \Rightarrow VSI [6, 16, 32]. For completeness, necessary and sufficient conditions for \mathbf{F} to be VSI₁ or VSI₂ are given in Appendix C.*

Remark 1.10 (ϵ -property). *From Lemma B.4 (Appendix B), it is easy to see that a tensor is VSI if and only if, given an arbitrary non-negative integer N , there exist a reference frame in which its*

⁴However, it is well-known that for $p = 2$ and $n = 4$ the only possible types are D and N [4, 31]. Note also that the type G does not occur for $p = 2$ and n even [30, 35, 36]. For $p = 1$ the type is G, D or N when the vector \mathbf{F} is timelike, spacelike or null, respectively (cf., e.g., [7]; in particular, $p = 1$ is the only case of interest when $n = 3$).

components and those of its covariant derivatives up to order N can be made as small as desired (the proof of this statement is essentially the same as the one given in [32] for the Riemann tensor). This applies, in particular, to a p -form \mathbf{F} and is an extension of the early observations of [1, 5] for the case $N = 0$.

In the rest of the paper we discuss further implications of Theorem 1.5 from a physical viewpoint. In section 2 we give the explicit form of VSI p -forms, the associated (degenerate Kundt) background metric and the corresponding Maxwell equations in adapted coordinates. We also observe that null (and thus VSI) Maxwell fields are “immune” to adding a Chern-Simons term to the Maxwell equations (except when it is linear, i.e., for $n = 2p - 1$). More generally, we point out that certain VSI Maxwell fields solve *any* generalized electrodynamics. In section 3 we consider VSI p -forms in the full Einstein-Maxwell theory (which solve also certain supergravities due to the vanishing of the Chern-Simons term). We discuss consequences of the Einstein equations, also mentioning a few examples, and we comment on various subclasses of exact solutions (such as VSI spacetimes and pp -waves) with arbitrary n and p . We observe that certain VSI Maxwell fields solve any generalized electrodynamics also when the coupling to gravity is kept into account. Appendix A contains a summary of properties of Kundt spacetimes which are useful in this paper. Appendix B gives the proof of Theorem 1.5. For completeness, in Appendix C we present the necessary and sufficient conditions for a p -form to be VSI_1 or VSI_2 , which can be seen as an “intermediate” result between Corollary 1.4 and Theorem 1.5.

Notation

In an n -dimensional spacetime we set up a frame of n real vectors $\mathbf{m}_{(a)}$ which consists of two null vectors $\boldsymbol{\ell} \equiv \mathbf{m}_{(0)}$, $\mathbf{n} \equiv \mathbf{m}_{(1)}$ and $n - 2$ orthonormal spacelike vectors $\mathbf{m}_{(i)}$, with $a, b \dots = 0, \dots, n - 1$ while $i, j \dots = 2, \dots, n - 1$ [39] (see also the review [7] and references therein). For indices i, j, \dots there is no need to distinguish between subscripts and superscripts. The Ricci rotation coefficients L_{ab} , N_{ab} and $\overset{i}{M}_{ab}$ are defined by [39]

$$L_{ab} = \ell_{a;b}, \quad N_{ab} = n_{a;b}, \quad \overset{i}{M}_{ab} = m_{a;b}^{(i)}, \quad (3)$$

and satisfy the identities [39]

$$L_{0a} = N_{1a} = N_{0a} + L_{1a} = \overset{i}{M}_{0a} + L_{ia} = \overset{i}{M}_{1a} + N_{ia} = \overset{i}{M}_{ja} + \overset{j}{M}_{ia} = 0. \quad (4)$$

Covariant derivatives along the frame vectors are denoted as

$$D \equiv \ell^a \nabla_a, \quad \Delta \equiv n^a \nabla_a, \quad \delta_i \equiv m^{(i)a} \nabla_a. \quad (5)$$

2 Explicit form of VSI electromagnetic fields and adapted coordinates

In this section we study VSI Maxwell *test* fields, i.e., without taking into account their backreaction on the spacetime geometry (which will be discussed in section 3).

2.1 Electromagnetic field and spacetime metric

It is useful to express explicitly the conditions 2 of Theorem 1.5 in a null frame adapted to $\boldsymbol{\ell}$ (but otherwise arbitrary), as defined above. Condition 2a reads (cf., e.g., [33])

$$F_{ab_1 \dots b_{p-1}} = p! F_{1i_1 \dots i_{p-1}} \ell_{[a} m_{b_1}^{(i_1)} \dots m_{b_{p-1}}^{(i_{p-1})}]. \quad (6)$$

The ‘‘Kundt part’’ of condition 2c means (A1), i.e.,

$$L_{i0} = 0, \quad L_{ij} = 0. \quad (7)$$

Conditions 2b and the remaining part of condition 2c are more conveniently represented in a null frame *parallelly transported along* ℓ (i.e., such that (A2) holds), where they take the form (cf. also appendix A, and recall that a Kundt metric for which the Riemann tensor and its *first* covariant derivative are of aligned type II is necessarily degenerate Kundt [37, 38])

$$DF_{1i_1\dots i_{p-1}} = 0, \quad R_{010i} = 0, \quad DR_{0101} = 0, \quad \text{with (A2)}. \quad (8)$$

In adapted coordinates, degenerate Kundt metrics are described by [37, 38]

$$ds^2 = 2du [dr + H(u, r, x)du + W_\alpha(u, r, x)dx^\alpha] + g_{\alpha\beta}(u, x)dx^\alpha dx^\beta, \quad (9)$$

where $\ell = \partial_r$ is the Kundt vector, $\alpha, \beta = 2 \dots n-1$, x denotes collectively the set of coordinates x^α , and $W_{\alpha,rr} = 0 = H_{,rrr}$ (thanks to which the second and third of (8) are identically satisfied), i.e.,

$$W_\alpha(u, r, x) = rW_\alpha^{(1)}(u, x) + W_\alpha^{(0)}(u, x), \quad (10)$$

$$H(u, r, x) = r^2H^{(2)}(u, x) + rH^{(1)}(u, x) + H^{(0)}(u, x). \quad (11)$$

In these coordinates one has (cf. the coordinate-independent expression (A3))

$$\ell_{a;b}dx^a dx^b = (2rH^{(2)} + H^{(1)})du^2 + \frac{1}{2}W_\alpha^{(1)}(du dx^\alpha + dx^\alpha du). \quad (12)$$

The corresponding VSI p -form (6) reads

$$\mathbf{F} = \frac{1}{(p-1)!} f_{\alpha_1 \dots \alpha_{p-1}}(u, x) du \wedge dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{p-1}}, \quad (13)$$

where $f_{\alpha_1 \dots \alpha_{p-1}} \equiv F_{u\alpha_1 \dots \alpha_{p-1}}$ is r -independent due to the first of (8). In these coordinates we have (with the definition (B9))

$$\mathcal{F}^2 = f_{\alpha_1 \dots \alpha_{p-1}} f^{\alpha_1 \dots \alpha_{p-1}}, \quad (14)$$

where, from now on, it is understood that the indices of $f^{\alpha_1 \dots \alpha_{p-1}}$ are raised using the transverse metric $g^{\alpha\beta}$. \mathcal{F}^2 parametrizes the field strength of \mathbf{F} and is invariant under Lorentz transformations preserving ℓ , as well as under transformations of the spatial coordinates $x \mapsto x'(x)$.

At this stage, $g_{\alpha\beta}$, $W_\alpha^{(1)}$, $W_\alpha^{(0)}$, $H^{(2)}$, $H^{(1)}$, $H^{(0)}$ and $F_{u\alpha_1 \dots \alpha_{p-1}}$ are all arbitrary functions of u and x (recall that the invariant condition $W_\alpha^{(1)} = 0 \Leftrightarrow L_{i1} = 0$ defines a special subfamily of Kundt metrics, cf. Appendix A.1). In general, the associated Weyl and Ricci tensor are both of aligned type II (as follows from Definition A.1 in Appendix A.2). The b.w. 0 components of all curvature tensors (and thus also their curvature invariants, of all orders) of the metric (9) are independent of the functions $W_\alpha^{(0)}$, $H^{(1)}$ and $H^{(0)}$ [40] (as summarized in Proposition 7.2 of [7]). Restrictions coming from the Einstein equations are described below in section 3. We emphasize here that although the p -form (13) is VSI, the spacetime (9) (with (10), (11)) in general is not (not even VSI₀) [6, 16, 41], i.e., it may admit some non-zero invariants constructed from the Riemann tensor and its derivatives. However, all mixed invariants (i.e., those involving Riemann *and* \mathbf{F} together, along with their derivatives) are necessarily zero (since \mathbf{F} possesses only negative b.w.s and Riemann only non-positive ones, and similarly for their derivatives.)

The metric (9) includes, in particular, spacetimes of constant curvature (Minkowski and (A)dS). Therefore, the p -form (13) can be used to describe VSI test fields in such backgrounds, as a special case.

2.2 Maxwell's equations

The construction of VSI p -forms has been, so far, purely geometric. Using (9) and (13), the source-free Maxwell equations $d\mathbf{F} = 0$ and $d^*\mathbf{F} = 0$ reduce, respectively, to

$$f_{[\alpha_2 \dots \alpha_{p-1}, \alpha_1]} = 0, \quad (\sqrt{\tilde{g}} f^{\beta \alpha_1 \dots \alpha_{p-2}})_{,\beta} = 0, \quad (15)$$

where $\tilde{g} \equiv \det g_{\alpha\beta} = -\det g_{ab} \equiv -g$. Effectively, these are Maxwell's equations for the $(p-1)$ -form \mathbf{f} in the $(n-2)$ -dimensional Riemannian geometry associated with $g_{\alpha\beta}$. In other words, the most general VSI \mathbf{F} that solves Maxwell's equations is given by (13) in the spacetime (9) with (10), (11), where \mathbf{f} is *harmonic* w.r.t. $g_{\alpha\beta}$.⁵ Recall, however, that \mathbf{f} can also depend on u .

2.2.1 The special case $p = 1$

The case $p = 1$ (or $p = n - 1$ by duality), for which $\mathbf{F} = \boldsymbol{\ell}$ is a vector field, is special. Indeed by Theorem 1.5 (see also Remark 1.7), if $\boldsymbol{\ell}$ is VSI then it must be degenerate Kundt and affinely parametrized. Further, by a boost, $\boldsymbol{\ell}$ can always be rescaled (still remaining VSI) such that $L_{1i} = L_{i1}$ (cf., e.g., [6, 17]). It is then easy to see that Maxwell's equations are satisfied (i.e., $\ell^a{}_{;a} = 0 = \ell_{[a;b]}$, cf. (A3)). Therefore, *for $p = 1$ to any VSI \mathbf{F} it can always be associated a solution to the sourcefree Maxwell equations* (in this statement, “VSI” can also be relaxed to “VSI₁”, since only the Kundt property of $\boldsymbol{\ell}$ has been employed, cf. also Proposition C.1)

2.2.2 The special cases $n = 4, 3$

Case $n = 4$ We have seen (Theorem 1.5) that, for any n and p , if a p -form \mathbf{F} is VSI then the multiply aligned null direction $\boldsymbol{\ell}$ is necessarily (degenerate) Kundt and thus, in particular, geodesic and shearfree. In the case $n = 4, p = 2$, the Robinson theorem [2, 4, 43] then implies that the family of null bivectors associated with $\boldsymbol{\ell}$ includes a solution to the sourcefree Maxwell equations. Namely, from a VSI \mathbf{F} one can always obtain a solution \mathbf{F}' to the Maxwell equations by means of a Lorentz transformation (spin and boost) preserving the null direction defined by $\boldsymbol{\ell}$ (of course, it may happen that \mathbf{F} is already a solution and no transformation is needed). The 2-form \mathbf{F}' will thus be also multiply aligned with $\boldsymbol{\ell}$ and therefore it will be still VSI (condition 2b of Theorem 1.5 will hold due to $d\mathbf{F}' = 0$, cf. Remark 1.8). Along with the observation in section 2.2.1 for the case $p = 1$ (or $p = 3$), one concludes that for $n = 4$, given a VSI field, one can always associate with it a VSI solution to the sourcefree Maxwell equations (as above, it suffices to assume that \mathbf{F} is VSI₁). This does not seem to be true when $n > 4$, in general.

Case $n = 3$ In three dimensions the situation is even simpler, since the only case of interest is $p = 1$ (dual to $p = 2$). The observation in section 2.2.1 thus immediately implies that also *for $n = 3$ to any VSI (or VSI₁) \mathbf{F} it can always be associated a solution to the sourcefree Maxwell equations.*

2.3 Maxwell-Chern-Simons' equations

Maxwell's equations (15) for a p -form field admit a generalization which includes a Chern-Simons term, i.e., $d\mathbf{F} = 0$ and $d^*\mathbf{F} + \alpha\mathbf{F} \wedge \dots \wedge \mathbf{F} = 0$, where $\alpha \neq 0$ is an arbitrary constant. The second term in the latter equation contains k factors \mathbf{F} , and the corresponding number of spacetime dimensions is given by $n = p(k+1) - 1$. Such modifications of Maxwell's equations appear, for example, in (the

⁵Note that eqs. (15) apply to type N Maxwell fields \mathbf{F} in *any* Kundt spacetime, so also to type N \mathbf{F} that are not VSI (i.e., we have not used (10) and (11) to obtain (15)). In the case $p = 2$, this agrees with the results of [42], once specialized to Maxwell fields of type N.

bosonic sector of) minimal supergravity in five and eleven dimensions (with $n = 5$, $p = 2$, $k = 2$ and $n = 11$, $p = 4$, $k = 2$, respectively – cf., e.g., [44] and references therein).

2.3.1 Generic case $k \geq 2$

Now, let us note that any p -form of type N (6) satisfies

$$\mathbf{F} \wedge \mathbf{F} = 0, \quad (16)$$

so that for a \mathbf{F} of type N, the Chern-Simons term vanishes identically, provided $k \geq 2$. Therefore, a type N solution \mathbf{F} of Maxwell’s theory (15) is automatically also a solution of Maxwell-Chern-Simons’ theory. This applies, in particular, to VSI solutions. This has been also noticed, e.g., in [45] in the case of certain VSI pp -waves in $n = 11$ supergravity.

2.3.2 Special case $k = 1$

The special case $k = 1$ results in a linear theory

$$d\mathbf{F} = 0, \quad d^*\mathbf{F} + \alpha\mathbf{F} = 0 \quad (n = 2p - 1). \quad (17)$$

It is clear that here the Maxwell equations are modified non-trivially also for type N fields. The second of (15) now has to be replaced by

$$(\sqrt{\tilde{g}} f^{\beta\alpha_1 \dots \alpha_{p-2}})_{,\beta} - \alpha \sqrt{\tilde{g}} \star f^{\alpha_1 \dots \alpha_{p-2}} = 0, \quad (18)$$

where \star is the Hodge dual in the transverse geometry $g_{\alpha\beta}$ (not to be confused with the n -dimensional Hodge dual $*$ in the full spacetime g_{ab}). Recall that a linear Chern-Simon term appears, e.g., in topological massive electrodynamics in three dimensions ($n = 3$, $p = 2$, $k = 1$) [46,47] (and references therein).

2.4 Universal solutions of generalized electrodynamics (test fields)

Theories of electrodynamics described by a Lagrangian depending also on the derivatives of the field strength have been proposed long ago in [24,25]. More recently, interest in higher-derivative theories has been also motivated by string theory, cf., e.g., [48–50] and references therein. In this context, the electromagnetic field is typically represented by a closed 2-form \mathbf{F} whose field equations contain “correction terms” constructed in terms of \mathbf{F} and its covariant derivatives.

The class of VSI Maxwell fields defined in this paper can be employed to identify a subset of VSI solutions that are “universal”, i.e., solving simultaneously any electrodynamics whose field equations can be expressed as $d\mathbf{F} = 0$, $*d*\tilde{\mathbf{F}} = 0$, where $\tilde{\mathbf{F}}$ can be any p -form constructed from \mathbf{F} and its covariant derivatives.⁶ It can be shown, for example, that any VSI Maxwell \mathbf{F} is universal if the background is a Kundt spacetime of Weyl and traceless-Ricci type III (aligned) with $DR = 0 = \delta_i R$. In particular, Ricci flat and Einstein Kundt spacetimes of Weyl type III/N/O can occur (an explicit example is given in section 3.2), the latter including Minkowski and (A)dS. Details and more general examples (also of Weyl type II) will be presented elsewhere.

It should be emphasized that the above definition of universality includes terms with arbitrary higher-order derivative corrections. If one restricts oneself to theories in which $\tilde{\mathbf{F}}$ is constructed

⁶We assume that $\tilde{\mathbf{F}}$ is constructed polynomially from these quantities (however, the scalar coefficients appearing in such polynomials need not be polynomials of the scalar invariants of \mathbf{F} and its covariant derivatives – cf., for instance, Born-Infeld’s theory). The same type of construction will be assumed for the energy-momentum tensor of generalized theories in section 3.2 (but see also footnote 11).

algebraically from \mathbf{F} , i.e., without taking derivatives of \mathbf{F} (like NLE, for which $n = 2p = 4$ and the modified Maxwell equations are of the form $d(f_1^* \mathbf{F} + f_2 \mathbf{F}) = 0$), then these admit as solutions *all* null solutions of Maxwell's equations (not necessarily VSI), without any restriction on the background geometry. This has been known for a long time in NLE [5, 19] (see also, e.g., [51]).

3 Einstein-Maxwell solutions

3.1 General equations

What discussed so far applies to VSI test fields, since we have not considered the consequences of the backreaction on the spacetime geometry. In the full Einstein-Maxwell theory this is described by the energy-momentum tensor associated with \mathbf{F}

$$T_{ab} = \frac{\kappa_0}{8\pi} \left(F_{ac_1 \dots c_{p-1}} F_b{}^{c_1 \dots c_{p-1}} - \frac{1}{2p} g_{ab} F^2 \right), \quad (19)$$

where $F^2 = F_{\alpha_1 \dots \alpha_p} F^{\alpha_1 \dots \alpha_p}$. With (6), T_{ab} in (19) takes the form of aligned pure radiation, and Einstein's equations with a cosmological constant $R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = 8\pi T_{ab}$ reduce to

$$R_{ab} = \frac{2\Lambda}{n-2} g_{ab} + \kappa_0 \mathcal{F}^2 \ell_a \ell_b. \quad (20)$$

The Ricci tensor of (9) thus must satisfy $R_{01} = \frac{2}{n-2} \Lambda$, $R_{ij} = \frac{2}{n-2} \Lambda \delta_{ij}$, $R_{1i} = 0$ and $R_{11} = \kappa_0 \mathcal{F}^2$ (with $R = 2n\Lambda/(n-2)$, while $R_{00} = 0 = R_{0i}$ identically since the Riemann type is II by construction). Using these, the b.w. 0 components of the Einstein equations imply that in (9), (11) (as follows readily from [42])

$$\mathcal{R}_{\alpha\beta} = \frac{2\Lambda}{n-2} g_{\alpha\beta} + \frac{1}{2} W_\alpha^{(1)} W_\beta^{(1)} - W_{(\alpha||\beta)}^{(1)}, \quad (21)$$

$$2H^{(2)} = \frac{\mathcal{R}}{2} - \frac{n-4}{n-2} \Lambda + \frac{1}{4} W^{(1)\alpha} W_\alpha^{(1)}, \quad (22)$$

where $\mathcal{R}_{\alpha\beta}$, \mathcal{R} and $||$ denote, respectively, the Ricci tensor, the Ricci scalar and the covariant derivative associated with $g_{\alpha\beta}$, and $W^{(1)\alpha} \equiv g^{\alpha\beta} W_\beta^{(1)}$. The first of these is an “effective Einstein equation” for the transverse metric, while the second one determines the function $H^{(2)}$. Note that the functions $W_\alpha^{(0)}$, $H^{(1)}$ and $H^{(0)}$ do not appear here.

With (21), the contracted Bianchi identity in the transverse geometry, i.e., $2\mathcal{R}_{\alpha\beta}{}^{||\beta} = \mathcal{R}_{,\alpha}$, tells us (after simple manipulations with the Ricci identity) that $W_\alpha^{(1)}$ is constrained by

$$W_{\alpha||\beta}^{(1)\beta} = \frac{1}{2} W^{(1)\beta} \left(3W_{\alpha||\beta}^{(1)} - W_{\beta||\alpha}^{(1)} \right) + W_\alpha^{(1)} \left(W^{(1)\beta}{}_{||\beta} - \frac{1}{2} W^{(1)\beta} W_\beta^{(1)} - \frac{2\Lambda}{n-2} \right). \quad (23)$$

Finally, the Einstein equations of negative b.w., which can be used to determine the functions

$H^{(1)}$ and $H^{(0)}$, reduce to [42]⁷

$$2H_{,\alpha}^{(1)} = -g_{\alpha\beta,u}{}^{||\beta} + 2W_{[\alpha||\beta]}^{(0)\beta} - 2W^{(0)\beta}W_{\alpha||\beta}^{(1)} + (W^{(0)\beta}W_{\beta}^{(1)})_{,\alpha} + W_{\alpha,u}^{(1)} + 2(\ln\sqrt{\tilde{g}})_{,\alpha} \\ + W_{\alpha}^{(1)} \left[W^{(0)\beta}W_{\beta}^{(1)} - W^{(0)\beta}{}_{||\beta} + (\ln\sqrt{\tilde{g}})_{,u} \right] + \frac{4\Lambda}{n-2}W_{\alpha}^{(0)}, \quad (24)$$

$$\Delta H^{(0)} + W^{(1)\alpha}H_{,\alpha}^{(0)} + W^{(1)\alpha}{}_{||\alpha}H^{(0)} = W^{(0)\beta}W_{\beta}^{(0)} \left(\frac{1}{2}W^{(1)\alpha}{}_{||\alpha} - \frac{2\Lambda}{n-2} \right) \\ + H^{(1)} \left[W^{(0)\alpha}{}_{||\alpha} - (\ln\sqrt{\tilde{g}})_{,u} \right] - \frac{1}{2}(W^{(0)\alpha}W_{\alpha}^{(1)})^2 + W^{(0)[\alpha||\beta]}W_{[\alpha||\beta]}^{(0)} + W_{\alpha,u}^{(0)}{}_{||\alpha} \\ - W^{(0)\beta} \left(2W^{(1)\alpha}W_{[\alpha||\beta]}^{(0)} + W_{\beta,u}^{(1)} - 2H_{,\beta}^{(1)} \right) - (\ln\sqrt{\tilde{g}})_{,uu} + \frac{1}{4}g^{\alpha\beta}{}_{,u}g_{\alpha\beta,u} - \kappa_0\mathcal{F}^2, \quad (25)$$

where Δ is the Laplace operator in the geometry of the transverse metric $g_{\alpha\beta}$ (not to be confused with the symbol \triangle defined in (5)). In addition to (21)–(25), the equations for the electromagnetic field (15) must also be satisfied (note that the only metric functions entering there are the $g_{\alpha\beta}$). When $\mathcal{F}^2 = 0$, eqs. (21)–(25) represent the vacuum Einstein equations for the most general Kundt spacetime.

Let us observe that the functions $\mathcal{F}^2(u, x)$ and $H^{(0)}(u, x)$ enter only (25), which is linear in $H^{(0)}(u, x)$. It follows, e.g., that given any Kundt Einstein spacetime (necessarily of the form (9) with (10), (11)), one can add to it an electromagnetic (and gravitational) wave by just appropriately choosing a new function $H^{(0)}(u, x)$, and leaving the other metric functions unchanged (amounting, in fact, to a generalized Kerr-Schild transformation; cf. Theorem 31.1 and section 31.6 of [4] in four dimensions). The resulting solution will describe a wave-like VSI p -form field propagating in the chosen Kundt Einstein spacetime. If the Einstein “seed” is VSI (or CSI), so will be the corresponding Einstein-Maxwell spacetime (see [40] and the comments in section 2.1).

The simplest such examples one can construct are electromagnetic and gravitational “plane-fronted” waves (with $W_{\alpha}^{(0)} = 0$) propagating in a constant curvature background, giving rise to Kundt waves of Weyl type N. These are well-known in four dimensions for any value of Λ [53, 54] (and include, e.g., the Siklos waves when $\Lambda < 0$; see also [4, 55, 56]), and have been considered also in arbitrary dimensions [57] (for $p = 2$, but a generalization to any p is straightforward). Similarly, one can construct, e.g., electrovac waves of Weyl type II in (anti-)Nariai product spaces for $n = 4$ [58] and higher [52]. All these examples are CSI spacetimes, and for $\Lambda = 0$ they become VSI (pp - or Kundt waves) spacetimes (cf. sections 3.1.1 and 3.1.2). More general (e.g., with $W_{\alpha}^{(0)} \neq 0$) degenerate Kundt metrics with null Maxwell fields are also known (see [4, 56, 59] and references therein for $n = 4$ and, e.g., [52, 60] in higher dimensions – more references in section 3.1.1 below in the case of VSI spacetimes).

In general, for all Einstein-Maxwell solutions with a VSI p -form \mathbf{F} , since all the mixed invariants are zero (cf. section 2.1) and since the Ricci tensor is constructed out of \mathbf{F} (eq. (20)), the only possible non-zero scalar invariants are those constructed from the Weyl tensor (and its derivatives), and the Ricci scalar R . We also remark that, thanks to the observations of section 2.3, all such

⁷Eqs. (24) and (25) are equations (76) and (73) of [42] (once the non-null part of the electromagnetic field is set to zero there, and up to minor reshuffling and different notation). Note that the remaining Einstein equations (71), (72) and (75) given in [42] are identically satisfied as a consequence of (21)–(25), (15) of the present paper, and can thus be dropped (cf. [4] when $n = 4$; for $n > 4$ this has also been discussed in [52] in the special case $W_{\alpha}^{(1)} = 0$). To verify this statement one needs repeated use of the Ricci identity and standard identities such as $g_{ab,c} = g_{ad}\Gamma_{bc}^d + g_{bd}\Gamma_{ac}^d$ and $\Gamma_{ab}^b = (\ln\sqrt{-g})_{,a} = \frac{1}{2}g^{bc}g_{bc,a}$. In the case of (72, [42]) one also needs the identity $g_{\alpha\beta,u}{}^{||\alpha\beta} - g^{\alpha\beta}g_{\alpha\beta,u}{}_{||\gamma}{}^{\gamma} = g^{\alpha\beta}(\tilde{\Gamma}_{\alpha\beta,u}^{\gamma} - \tilde{\Gamma}_{\gamma\beta,u}^{\alpha}) = g^{\alpha\beta}\mathcal{R}_{\alpha\beta,u}$ (where $\tilde{\Gamma}_{\alpha\beta,u}^{\gamma}$ are the Christoffel symbols of $g_{\alpha\beta}$ – in the last equality the (contracted) standard definition of the Riemann tensor has been employed).

Einstein-Maxwell solutions having $n = 5, p = 2$ or $n = 11, p = 4$ are also solutions (with the same \mathbf{F}) of the bosonic sector of 5D minimal supergravity (ungauged or gauged) and 11D supergravity, respectively (further comments and references in section 3.1.1).

3.1.1 VSI spacetimes with VSI Maxwell fields

The special case when the spacetime metric is VSI is of particular interest. All VSI metrics in HD (in particular, with Ricci type N) are given in [41] (see also [61]). In a VSI spacetime, one can always choose coordinates such that the metric is given by (9) with [41]

$$g_{\alpha\beta} = \delta_{\alpha\beta}, \quad W_\alpha^{(1)} = -\delta_{\alpha,2} \frac{2\epsilon}{x_2}, \quad H^{(2)} = \frac{\epsilon}{2(x^2)^2} \quad (\epsilon = 0, 1). \quad (26)$$

The VSI assumption implies $\Lambda = 0$, so that the Einstein equations (20) give $R_{ab} = \kappa_0 \mathcal{F}^2 \ell_a \ell_b$, i.e., the Ricci tensor is of aligned type N (and (21) and (22) are satisfied identically). This constrains the functions $W_\alpha^{(0)}$, $H^{(1)}$ and $H^{(0)}$, as detailed in [41]. The Weyl tensor is of type III aligned with $\ell = \partial_r$ (it becomes of type N for special choices of $W_\alpha^{(0)}$, including $W_\alpha^{(0)} = 0$,⁸ and of type O under further conditions on $H^{(0)}$ [41]). The Maxwell field is still given by (13). Some VSI spacetimes (more general than VSI pp -waves) coupled to null p -forms have been discussed in [61] in the context of type IIB supergravity (but note that a few of those are of Ricci type III due to the presence of an additional non-trivial dilaton⁹).

As a special subcase, for $\epsilon = 0 = H^{(1)}$ one obtains VSI pp -waves with a null p -form, for which the Weyl type can only be III(a) or more special (since the Ricci type is N [41]). In 4D, these are well-known and necessarily of Weyl type N (which coincides with the type III(a) for $n = 4$ [7]) or O (cf. section 24.5 of [4] and references therein). In higher dimensions, their role in the context of supergravity and string theory has been known for some time, see, e.g., [11–13, 65–68]. More recently, some of these (with n arbitrary, $p = 2, 3$ and $W_\alpha = W_\alpha^{(0)}(u, x) \neq 0$) have been interpreted as “charged gyratons” [69, 70].¹⁰ From the viewpoint of supersymmetry, it is worth recalling also that a VSI spacetime admitting a timelike or null Killing vector field must necessarily be a VSI pp -wave (see the appendix of [61]). Supersymmetric VSI pp -waves coupled to null p -forms are indeed well-known in various supergravities (see, e.g., [11, 65, 66, 68, 71, 72] and references therein).

3.1.2 pp -waves with VSI Maxwell fields

As mentioned above, spacetimes admitting a null Killing vector field are of special interest for supersymmetry. A null Killing vector field aligned with the Ricci tensor is necessarily Kundt (cf.,

⁸We note that, when the Weyl type is N, the metric takes the Kerr-Schild form – the argument given in section 4.2.2 of [62] in the vacuum case holds also for Ricci type N. When the Weyl type is III, they are of the more general “extended Kerr-Schild” form [63].

⁹We did not consider a dilaton φ in our discussion, but this can be easily included. The dilaton φ itself cannot be VSI unless zero, being a scalar field. However, if we want $\varphi_{,a}$ to be VSI, then it must obviously be a null vector field. If we also require the mixed invariant $F^a{}_{c_1 \dots c_{p-1}} F_b{}^{c_1 \dots c_{p-1}} \varphi_{,a} \varphi_{,b}$ to vanish, we immediately obtain $\varphi_{,a} \propto \ell_a = (du)_a$, which implies $\varphi = \varphi(u)$ (as assumed in [11, 13, 61, 64]). This is also a sufficient condition for $\varphi_{,a}$ to be VSI (since $\varphi_{,a} = \varphi' \ell_a$ and ℓ is degenerate Kundt, cf. Remark 1.7) and for all mixed invariants (i.e., containing φ , the Riemann/Maxwell tensors and their derivatives) to vanish as well. However, $\varphi_{;ab} = \varphi'' \ell_a \ell_b + \varphi' \ell_{a;b}$ is a symmetric rank-2 tensor (of aligned type III or more special) which contains also non-zero components of b.w. -1 iff $\varphi' W_\alpha^{(1)} \neq 0$ (cf. (12)) and appears on the r.h.s. of Einstein’s equations [61], thus modifying the Ricci type.

¹⁰It was indeed observed in [69] that the ansatz used there leads to VSI spacetimes in which also all the electromagnetic scalar invariants vanish. This appears as a special subcase of the result given in Theorem 1.5.

e.g., Proposition 8.21 of [7]). Now, in the metric (9), the Kundt vector $\ell = \partial_r$ is Killing iff

$$W_\alpha^{(1)} = 0, \quad H^{(2)} = 0 = H^{(1)}, \quad (27)$$

which is equivalent to requiring (9) to be a pp -wave [73], i.e., $\ell_{a;b} = 0$ (cf. (12)).

For pp -waves the Einstein equations (20) imply $\Lambda = 0$ (cf., e.g., Proposition 7.3 of [7]), so that $R_{ab} = \kappa_0 \mathcal{F}^2 \ell_a \ell_b$ is of type N. By (21), it follows that the transverse metric $g_{\alpha\beta}(u, x)$ must be Ricci-flat (if it is flat, as happens necessarily for $n = 4, 5$, one has VSI pp -waves, cf. section 3.1.1), and (22) is then identically satisfied. Constraints on the functions $W_\alpha^{(0)}$ and $H^{(0)}$ follow from the remaining Einstein equations (24), (25), i.e.,

$$\begin{aligned} 2W_{[\alpha|\beta]}^{(0)\beta} &= g_{\alpha\beta,u} \parallel^\beta - 2(\ln \sqrt{\tilde{g}})_{,u\alpha}, \\ \Delta H^{(0)} &= W^{(0)[\alpha|\beta]} W_{[\alpha|\beta]}^{(0)} + W_{\alpha,u}^{(0)\parallel\alpha} - (\ln \sqrt{\tilde{g}})_{,uu} + \frac{1}{4} g^{\alpha\beta}_{,u} g_{\alpha\beta,u} - \kappa_0 \mathcal{F}^2. \end{aligned} \quad (28)$$

Similarly as for Ricci flat pp -waves (cf. Table 2 of [74] and Proposition 7.3 of [7]), here the Weyl type is II'(abd) (which reduces to III(a) for $n = 5$ and to N for $n = 4$) or more special. The Maxwell field is given by (13).

Some CSI (non-VSI) pp -waves coupled to null forms in Ricci-flat direct products arise as special cases of the solutions of [75] (where $n = 11$ and $p = 4$).

3.2 Universal Einstein-Maxwell solutions

In section 2.4 we commented on the role of a subset of the VSI Maxwell fields as universal solutions of all generalized electrodynamics on certain backgrounds (test fields). More generally, some of those can also be used to construct exact solutions of full general relativity, i.e., keeping into account the backreaction of the electromagnetic field on the spacetime geometry. This is described by Einstein's equations in which, however, the T_{ab} associated with the electromagnetic field is determined in the generalized electrodynamics (in terms of \mathbf{F} and its covariant derivatives, thus being generically different from (19) – cf., e.g., [25]). The class of universal Einstein-(generalized-)Maxwell solutions deserves a more detailed study, which we will present elsewhere. Here we only point out some examples. Namely, it can be shown that all VSI spacetimes with $L_{i1} = 0 = L_{1i}$ (i.e., the recurrent ones) coupled to an aligned VSI p -form field that solve the standard Einstein-Maxwell equations (and are thus of Ricci type N) are also exact solutions of gravity coupled to generalized electrodynamics,¹¹ provided $p > 1$ and $\delta_i F_{1j_1 \dots j_{p-1}} = 0$ (in an “adapted” parallelly transported frame, i.e., such that $M_{jk}^i = 0$). Within this family, metrics of Weyl type N are necessarily pp -waves, for which such a universal property was pointed out in [11, 13, 14], at least for certain values of p ([11] considered only plane waves, but included also Yang-Mills field). But metrics of Weyl type III are also permitted, including pp -waves ($L_{11} = 0$) and also genuinely recurrent ($L_{11} \neq 0$) spacetimes (for $n = 4$, $p = 3$

¹¹To be precise, we should exclude from the discussion possible peculiar theories admitting an energy-momentum tensor T_{ab} that vanishes for certain non-zero electromagnetic fields (or at least for the “universal” ones). One possible way to ensure this is, for example, to consider only theories for which T_{ab} is of the form $T_{ab} \propto T_{ab}^{EM} + \text{“corrections”}$, where T_{ab}^{EM} is the energy-momentum tensor of the standard Einstein-Maxwell theory, and the “corrections” are terms that go to zero faster than T_{ab}^{EM} in the limit of weak fields. Additionally, it is also understood that a constant rescaling of a universal p -form \mathbf{F} (or, alternatively, of the corresponding metric, and so also the Ricci tensor) may be necessary when going from one theory to another.

this was discussed in [64]). One explicit example of the latter solutions in 4D is given by¹²

$$ds^2 = 2du \left[dr + \frac{1}{2} (xr - xe^x - 2\kappa_0 e^x c^2(u)) du \right] + e^x (dx^2 + e^{2u} dy^2), \quad (29)$$

$$\mathbf{F} = e^{x/2} c(u) du \wedge \left(-\cos \frac{ye^u}{2} dx + e^u \sin \frac{ye^u}{2} dy \right). \quad (30)$$

Similarly as in section 2.4, the above discussion applies to generalized electrodynamics with arbitrary higher-order derivative corrections. As a special case, the fact that Einstein-Maxwell solutions with aligned null electromagnetic fields (not necessarily VSI) are also solution of NLE coupled to gravity was previously demonstrated in [21–23].

Acknowledgments

This work has been supported by research plan RVO: 67985840 and research grant GAČR 13-10042S.

A Some of the Newman-Penrose equations for Kundt spacetimes

A.1 General Kundt spacetimes

By assumption ℓ is geodesic and Kundt (i.e., expansionfree, shearfree and twistfree), so that (recall the definitions (3)) [6, 39, 76]

$$L_{i0} = 0, \quad L_{ij} = 0. \quad (A1)$$

Without loss of generality we can use *an affine parametrization and a frame parallelly transported along ℓ* , such that, in addition to (A1), we also have [6, 39, 76]

$$L_{10} = 0, \quad \overset{i}{M}_{j0} = 0, \quad N_{i0} = 0. \quad (A2)$$

Thanks to these, the covariant derivatives of the frame vectors take the form [6, 39]

$$\ell_{a;b} = L_{11} \ell_a \ell_b + L_{1i} \ell_a m_b^{(i)} + L_{i1} m_a^{(i)} \ell_b, \quad (A3)$$

$$n_{a;b} = -L_{11} n_a \ell_b - L_{1i} n_a m_b^{(i)} + N_{i1} m_a^{(i)} \ell_b + N_{ij} m_a^{(i)} m_b^{(j)}, \quad (A4)$$

$$m_{a;b}^{(i)} = -N_{i1} \ell_a \ell_b - L_{i1} n_a \ell_b - N_{ij} \ell_a m_b^{(j)} + \overset{i}{M}_{j1} m_a^{(j)} \ell_b + \overset{i}{M}_{kl} m_a^{(k)} m_b^{(l)}. \quad (A5)$$

From the Ricci identities (11g) and (11k) of [76] with (A1), (A2) it follows immediately

$$R_{0i0j} = 0, \quad R_{0ijk} = 0, \quad (A6)$$

¹²This solution has been obtained by adding a null \mathbf{F} to a type III vacuum spacetime found by Petrov (eq. (31.40) in [4]), cf. section 3.1. A parallelly transported frame satisfying $\overset{i}{M}_{jk} = 0$ is given by $\ell = \partial_r$, $\mathbf{n} = \partial_u - \frac{1}{2} (xr - xe^x - 2\kappa_0 e^x c^2(u)) \partial_r$, $\mathbf{m}_2 = e^{-x/2} \left(\cos \frac{ye^u}{2} \partial_x - e^{-u} \sin \frac{ye^u}{2} \partial_y \right)$, $\mathbf{m}_3 = e^{-x/2} \left(\sin \frac{ye^u}{2} \partial_x + e^{-u} \cos \frac{ye^u}{2} \partial_y \right)$. Note that by setting $\kappa_0 = 0$ in (29), this solution represents a universal *test* Maxwell field in a Petrov type III vacuum spacetime, relevant to the discussion in section 2.4.

which implies that all Kundt spacetimes are of aligned Riemann type I, with the further restriction $R_{0ijk} = 0$ (cf. [38, 42]). Furthermore, the Ricci identities (11b), (11e), (11n), (11a), (11j), (11m) and (11f) of [76] read

$$DL_{1i} = -R_{010i}, \quad DL_{i1} = -R_{010i}, \quad (\text{A7})$$

$$D\overset{i}{M}_{jk} = 0, \quad (\text{A8})$$

$$DL_{11} = -L_{1i}L_{i1} - R_{0101}, \quad (\text{A9})$$

$$DN_{ij} = -R_{0j1i}, \quad (\text{A10})$$

$$D\overset{i}{M}_{j1} = -\overset{i}{M}_{jk}L_{k1} - R_{01ij}, \quad (\text{A11})$$

$$DN_{i1} = -N_{ij}L_{j1} + R_{101i}, \quad (\text{A12})$$

while the commutators [6] needed in this paper simplify to

$$\Delta D - D\Delta = L_{11}D + L_{i1}\delta_i, \quad (\text{A13})$$

$$\delta_i D - D\delta_i = L_{1i}D. \quad (\text{A14})$$

In adapted coordinates, the general Kundt line-element can be written as (9), where $\ell = \partial_r$ and all the metric functions depend arbitrarily on their arguments. In those coordinates $L_{1i} = L_{i1}$ (since $\ell_a = (du)_a$), cf. (12). From (12), (A3) it follows that ℓ is *recurrent* iff $W_\alpha^{(1)} = 0 \Leftrightarrow L_{1i} = L_{i1} = 0$, which is equivalent to $[\delta_i, D] = 0$ – this condition can be used to invariantly characterize subfamilies of Kundt spacetimes.

A.2 Kundt spacetimes of aligned Riemann type II

The above results hold for *any* Kundt spacetime. If one now restricts to the Kundt spacetimes of *aligned Riemann type II* (i.e., we assume $R_{010i} = 0$ in addition to (A6)),¹³ using (A1), (A2) the Bianchi identities (B3), (B5), (B12), (B1), (B6) and (B4) of [39] reduce to

$$DR_{01ij} = 0, \quad (\text{A15})$$

$$DR_{0i1j} = 0, \quad (\text{A16})$$

$$DR_{ijkl} = 0, \quad (\text{A17})$$

$$DR_{101i} - \delta_i R_{0101} = -R_{0101}L_{i1} - R_{01is}L_{s1} - R_{0i1s}L_{s1}, \quad (\text{A18})$$

$$DR_{1kij} + \delta_k R_{01ij} = R_{01ij}L_{k1} - 2R_{0k1[i}L_{j]1} + R_{ksij}L_{s1} - 2R_{01[i]s}\overset{s}{M}_{j]k}, \quad (\text{A19})$$

$$DR_{1i1j} - \Delta R_{0j1i} - \delta_j R_{101i} = R_{0101}N_{ij} - R_{01is}N_{sj} + R_{0s1i}N_{sj} + R_{0j1s}\overset{s}{M}_{i1} + R_{0s1i}\overset{s}{M}_{j1} \\ + 2R_{101i}L_{[1j]} + R_{1ijs}L_{s1} + R_{101s}\overset{s}{M}_{ij}. \quad (\text{A20})$$

Note that here (A7) reduces to $DL_{1i} = 0 = DL_{i1}$, so that differentiation of (A9) gives

$$D^2L_{11} = -DR_{0101}. \quad (\text{A21})$$

An important subset of Kundt spacetimes of Riemann type II is given by the *degenerate* Kundt metrics [37, 38], defined by

Definition A.1 (Degenerate Kundt metrics [37, 38]). *A Kundt spacetime is “degenerate” if the Kundt null direction ℓ is also a multiple null direction of the Riemann tensor and of its covariant derivatives of arbitrary order (which are thus all of aligned type II, or more special).*

¹³This is equivalent to saying that ℓ is a multiply aligned null directions of both the Weyl and the Ricci tensors.

It is worth emphasizing that, in fact, the degenerate condition is automatically met at all orders once it is satisfied by the Riemann tensor and its *first* derivative (Theorem 4.2 and section 7 of [38]). For degenerate Kundt spacetimes we have the following

Proposition A.2 (Conditions for degenerate Kundt metrics). *A Kundt spacetime is degenerate iff it is of aligned Riemann type II and $\ell^a R_{,a} = 0$ (using an affine parameter and a parallelly transported frame, the latter condition is equivalent to any of the following: $DR = 0$, $DR_{01} = 0$ or (for $n > 3$) $DC_{0101} = 0^{14}$). A Kundt spacetime for which the tracefree part of the Ricci tensor is of aligned type III is necessarily degenerate.*

Proof. It is a result of from [37,38] that for Kundt spacetimes the degenerate condition is equivalent to the Riemann type II with $DR_{0101} = 0$. The first part of the proposition thus simply follows from $DR_{0101} = 0$ and the contracted Bianchi identities. The second part follows using Proposition 2 of [76] (implying the Weyl type II – cf. also Proposition 7.1 of [7]) and, again, the contracted Bianchi identities (whose component of b.w. +1 gives $DR = 0$). \square

For Kundt spacetimes of Riemann type II, the degenerate condition is also equivalent to $D^2 L_{11} = 0$ (cf. (A21)). Note that the assumptions on the Ricci tensor in the second part of Proposition A.2 are of physical interest since they correspond to the case when the energy-momentum tensor is triply aligned with ℓ . See Remark 1.6 for further comments (and Proposition 6.1 of [38] for an alternative invariant characterization of degenerate Kundt metrics – discussed for $n = 4$ in [38], but holding for any n).

B Proof of Theorem 1.5

B.1 Proof of “2. \Rightarrow 1.”

B.1.1 Preliminaries

Before starting with the proof, let us make a few helpful observations on the strategy we shall adopt. First, by assumption 2a, \mathbf{F} is VSI₀. If we are able to show that all its covariant derivatives are of aligned type III (or more special), then we are done with the proof.

Now, by assumption 2c, ℓ is Kundt. Using an affine parameter and a frame parallelly transported along ℓ , this implies that the covariant derivatives of the frame vectors do not produce terms of higher b.w., cf. (A3)–(A5) (i.e., $\ell_{a;b}$ has only components of b.w. -1 or less, etc.). Together with assumption 2b (which here can be written as the first of (8)), this immediately shows that $\nabla \mathbf{F}$ is of aligned type III, as required (cf. also Proposition C.1 in Appendix C). The problem is now to show that the same is true for covariant derivatives of \mathbf{F} of *arbitrary* order.

To this end, the *balanced-scalar approach* of [6,16] will be useful. This approach can be applied to various tensors (or spinors). In the context of VSI tensors, the main idea is to show that (under proper assumptions) the covariant derivative of a tensor of type III is necessarily of aligned type III and thus, *by induction*, the tensor under consideration is VSI. Let us thus recall the relevant definition of [6,16]:

Definition B.1 (Balanced scalars and tensors [6,16]). *In a frame parallelly transported along an affinely parameterized geodesic null vector field ℓ , a scalar η of b.w. b under a constant boost is a “balanced scalar” if $D^{-b}\eta = 0$ for $b < 0$ and $\eta = 0$ for $b \geq 0$. A tensor whose components are all balanced scalars is a “balanced tensor”.*

¹⁴Recall that for $n = 3$ the Weyl tensor vanishes identically and therefore the condition $DC_{0101} = 0$ is trivial.

Note, in particular, that balanced tensors are of type III (or more special), multiply aligned with ℓ . Restating the inductive method mentioned above in more technical terms, this will thus consist in showing that the covariant derivative of a balanced tensor is again a balanced tensor (Lemma B.3 below, which will then apply to \mathbf{F}).¹⁵

B.1.2 Proof

We are thus ready to prove the direction “2. \Rightarrow 1.” of Theorem 1.5.

By assumption 2c, ℓ is degenerate Kundt (and thus geodesic). Eq. (A1) is satisfied and, employing an affine parameter and a parallelly transported frame, also (A2) and (A6) hold, along with $R_{010i} = 0$ and $DR_{0101} = 0$. Using the Ricci and Bianchi identities and the commutators summarized in appendix A, one easily arrives at

$$DL_{1i} = 0, \quad DL_{i1} = 0, \quad D\overset{i}{M}_{jk} = 0, \quad (\text{B1})$$

$$D^2N_{ij} = 0, \quad D^2\overset{i}{M}_{j1} = 0, \quad D^2L_{11} = 0, \quad D^3N_{i1} = 0. \quad (\text{B2})$$

Together with the commutators (A13) and (A14), this suffices to readily extend Lemma 4 of [6] (see [6, 16] for more technical details), i.e.,

Lemma B.2 (Balanced scalars in degenerate Kundt spacetimes). *In a degenerate Kundt spacetime, employing an affine parameter and a parallelly transported frame, if η is a balanced scalar of b.w. b , then all the following scalars (ordered by b.w.) are also balanced: $D\eta$; $L_{1i}\eta$, $L_{i1}\eta$, $\overset{i}{M}_{jk}\eta$, $\delta_i\eta$; $L_{11}\eta$, $N_{ij}\eta$, $\overset{i}{M}_{j1}\eta$, $\Delta\eta$, $N_{i1}\eta$.*

Now, considering (A3)–(A5) and Lemma B.2, one can easily extend also Lemma 6 of [6] (cf. also [16] for a spinorial version of in four dimensions), i.e.,

Lemma B.3 (Derivatives of balanced tensors in degenerate Kundt spacetimes). *In a degenerate Kundt spacetime, the covariant derivative of a balanced tensor is again a balanced tensor.*

Next, by assumptions 2a and 2b, we also have that (6) and the first of (8) hold. But these two equations precisely mean that \mathbf{F} is a balanced tensor. By Lemma B.3, the covariant derivatives of arbitrary order of \mathbf{F} are thus balanced tensors. In particular, they all possess only components of negative boost weight, which implies that \mathbf{F} is VSI, as we wanted to prove.

B.2 Proof of “1. \Rightarrow 2.”

B.2.1 Preliminaries

Let us start by proving two useful lemmas (of some interest in their own) about general tensors with vanishing invariants.¹⁶ The first of these (Lemma B.4) follows directly from Theorem 2.1 and Corollary 3.2 of [28] (since \mathbf{T} and its covariant derivatives up to order I are clearly not characterized by their invariants). We nevertheless provide an independent simple proof.

¹⁵In the balanced-scalar approach, it is convenient to assign a b.w. to all the Newman-Penrose quantities, and this is why we consider only *constant* boosts in the Definition B.1 (so that, e.g., L_{11} has b.w. -1 and, if η has b.w. b , then $D\eta$ and $\Delta\eta$ have, respectively, b.w. $(b+1)$ and $(b-1)$, etc. – these quantities would *not* admit a b.w. under a general boost [33]). It is important to observe that there is no loss of generality here as far as our proof is concerned – cf. also [6, 17, 18].

¹⁶To avoid confusion, let us emphasize that in the present section B.2.1, and only here, \mathbf{T} can be any tensor and does not necessarily coincide with the energy-momentum tensor (19).

Lemma B.4 (Alignment of VSI tensors). *If a tensor field \mathbf{T} is VSI_I , \mathbf{T} and its covariant derivatives up to order I are of aligned type III (or more special).*

Proof. The case $I = 0$ is contained in the algebraic VSI theorem (Theorem 1.3), so we need to discuss only the case $I \geq 1$. Additionally, the lemma is trivially true in the case $\nabla\mathbf{T} = 0$, so that we can assume hereafter $\nabla\mathbf{T} \neq 0$. \mathbf{T} being VSI_I implies that \mathbf{T} and its covariant derivatives up to order I are all VSI_0 (Definition 1.2). That all these tensors are of type III thus follows immediately from [28]. It remains to be proven that they are all aligned (i.e., for all of them the same null vector ℓ defines a multiple null direction such that all the non-negative b.w. components vanish).

Since \mathbf{T} is of type III, it admits a *unique* multiply aligned null direction such that all the non-negative b.w. components vanish. Furthermore, such tensor cannot admit a distinct null direction with respect to which all positive b.w. components vanish. Let us work in a null frame such that ℓ is parallel to this unique null direction. In this frame, \mathbf{T} possesses only components of negative b.w.. Now, let us consider two possible cases separately. *i*) First, if \mathbf{T} has only components of b.w. -2 or less, then the components of $\nabla\mathbf{T}$ have b.w. 0 or less (since the covariant derivative of a tensor can, at most, raise the b.w. of $+2$), i.e., also $\nabla\mathbf{T}$ is multiply aligned to ℓ . But $\nabla\mathbf{T}$ must be of type III (as noticed above), therefore the only possibility is that the components of $\nabla\mathbf{T}$ have, in fact, b.w. -1 (or less) in the frame we are using, so $\nabla\mathbf{T}$ is of *aligned* type III. *ii*) On the other hand, if \mathbf{T} has some components of b.w. -1 , we cannot use the same argument, since some components of $\nabla\mathbf{T}$ will have b.w. $+1$, in general. Let us thus assume this is indeed the case (if not, i.e., if components of $\nabla\mathbf{T}$ have only b.w. 0 or less, then we can proceed as in case *i*) and let us consider, instead, the tensor product $\mathbf{T}_\times \equiv \mathbf{T} \times \nabla\mathbf{T}$. Obviously, the components of \mathbf{T}_\times cannot have b.w. greater than 0 (cf., e.g., Proposition A.11 of [34]). But since \mathbf{T} is VSI_1 , then \mathbf{T}_\times must be of type III, which thus implies that the components of \mathbf{T}_\times can only have b.w. -1 or smaller in a frame adapted to ℓ . However, it is not difficult to see that this cannot be true if $\nabla\mathbf{T}$ possesses some components of b.w. $+1$ (as we assumed), thus leading to a contradiction. In other words, if \mathbf{T} is VSI_1 then $\nabla\mathbf{T}$ can only have components of b.w. 0 or less; but then, in fact, these can be only of b.w. -1 or less (since $\nabla\mathbf{T}$ must be of type III, similarly as in point *i*)), so that, again, $\nabla\mathbf{T}$ is of *aligned* type III.

Combining *i*) and *ii*) we have proven the lemma for the case $I = 1$. Clearly the same argument extends to any higher I , i.e., the proof is complete. □

In turn, this can be used to prove the following result for rank-2 tensors.

Lemma B.5 (T_{ab} VSI_2 implies Kundt). *If a 2-tensor field T_{ab} is VSI_2 then : (a) T_{ab} , $T_{ab;c}$ and $T_{ab;cd}$ are of aligned type III (or more special); (b) the corresponding multiple null direction ℓ is necessarily Kundt.*

Proof. Point (a) follows immediately from Lemma B.4. It will be used in the following.

Now, thanks to (a), in an adapted null frame we can write

$$T_{ab} = T_{1i}\ell_a m_b^{(i)} + T_{i1}m_a^{(i)}\ell_b + T_{11}\ell_a\ell_b. \quad (\text{B3})$$

It is convenient to first prove (b) in the special case $T_{1i} = 0 = T_{i1}$, i.e., $T_{ab} = T_{11}\ell_a\ell_b$ (of course with $T_{11} \neq 0$). Let us define a compact notation for the covariant derivatives

$$T_{abc_1\dots c_I}^{(I)} \equiv T_{ab;c_1\dots c_I} \quad (I = 1, 2, 3, \dots). \quad (\text{B4})$$

For our purposes, it will now suffice to require that certain components of b.w. 0 of $\mathbf{T}^{(1)}$ and $\mathbf{T}^{(2)}$ vanish (in view of (a)). First, requiring $T_{i10}^{(1)} = 0$ one obtains (using also the first of (3))

$$L_{i0} = 0, \quad (\text{B5})$$

i.e., ℓ must be geodesic (note that the VSI_1 property of \mathbf{T} suffices to prove this). Next, the condition $T_{01ij}^{(2)} = 0$ is equivalent to $L_{ki}L_{kj} = 0$, which, by tracing, leads to

$$L_{ij} = 0, \quad (\text{B6})$$

i.e. (together with (B5)), ℓ must be a Kundt null direction, which thus proves (b) in the case $T_{ab} = T_{11}\ell_a\ell_b$.

Finally, for a generic T_{ab} (eq. (B3)) we can apply the same argument to the tensor $\tilde{T}_{ab} \equiv T_{ac}T_b{}^c = (T_{1i}T_{i1})\ell_a\ell_b$ if $T_{1i} \neq 0$ (or to $\tilde{T}_{ab} \equiv T_{ca}T_b{}^c = (T_{i1}T_{i1})\ell_a\ell_b$ if $T_{i1} \neq 0$), so the proof of (b) is now complete (where we used the fact that if T_{ab} is VSI_2 then \tilde{T}_{ab} must obviously also be VSI_2). \square

B.2.2 Proof

Let us now prove the direction “1. \Rightarrow 2.” of Theorem 1.5. We assume that \mathbf{F} is VSI. It is, in particular, VSI_0 and thus, by Corollary 1.4, \mathbf{F} must be of type N, i.e., condition 2a is proven. In an adapted frame this means that (6) holds. It remains to prove that conditions 2b and 2c are also satisfied, i.e., we need to show (recall (7) and (8))

- (i) $L_{i0} = 0, L_{ij} = 0$
- (ii) $R_{010i} = 0, DR_{0101} = 0$ (in a parallelly transported frame)
- (iii) $DF_{1i_1\dots i_{p-1}} = 0$ (in a parallelly transported frame).

It is convenient to define the following 2-tensor \mathbf{T} ¹⁷

$$T_{ab} = \frac{\kappa_0}{8\pi} F_{ac_1\dots c_{p-1}} F_b{}^{c_1\dots c_{p-1}}. \quad (\text{B7})$$

Since \mathbf{F} is VSI, then \mathbf{T} must also be VSI, thus condition (i) follows immediately from Lemma B.5 applied to \mathbf{T} , so that ℓ is Kundt. (Alternatively, condition (i) can also be proven using Proposition C.1 in Appendix C – note also that so far we used only the assumption that \mathbf{F} is VSI_2 .)

For the next steps, it is useful to observe that, by (6), \mathbf{T} is also of type N w.r.t. ℓ , i.e.,

$$T_{ab} = \frac{\kappa_0}{8\pi} \mathcal{F}^2 \ell_a \ell_b, \quad (\text{B8})$$

where

$$\mathcal{F}^2 \equiv F_{1i_1\dots i_{p-1}} F_{1i_1\dots i_{p-1}} \geq 0, \quad (\text{B9})$$

and has b.w. -2 . Since the b.w. of \mathbf{F} and $\nabla^{(I)}\mathbf{F}$ is always -1 or less (Lemma B.4), also the b.w. of $\mathbf{T}^{(I)}$ (defined in (B4)) must always be -2 (or less; cf. also Proposition A.11 of [34]). This condition will be used below.

From now on, we employ an affine parameter along the Kundt null vector ℓ and a frame parallelly transported along it, so that (A2) holds. As observed in appendix A, this implies $R_{0i0j} = 0 = R_{0ij0} = 0$ (eq. (A6)).

Next, requiring $\nabla\mathbf{F}$ to be of type III (namely, $(\nabla\mathbf{F})_{01i_1\dots i_{p-1}} = 0$) is now equivalent to condition (iii), which is thus also proven (again, this alternatively follows from Proposition C.1).

Condition (iii) in turns implies $D(\mathcal{F}^2) = 0$, so that by a boost we can set $\mathcal{F}^2 = 1$ in (B8) (while preserving the affine parametrization of ℓ and the parallel transport of the frame), i.e., from now on $T_{ab} = \frac{\kappa_0}{8\pi} \ell_a \ell_b$.

¹⁷For a VSI_0 field \mathbf{F} , \mathbf{T} equals the associated energy-momentum tensor (19), since $F^2 = 0$.

Now, imposing $T_{11i0}^{(2)} = 0$ and $T_{i110}^{(2)} = 0$ (these components have b.w. -1 and thus must vanish, as observed above) gives, respectively,

$$DL_{1i} = 0, \quad DL_{i1} = 0. \quad (\text{B10})$$

By (A7) this in turn implies

$$R_{010i} = 0, \quad (\text{B11})$$

so that the Riemann type is II or more special (recall that we already obtained $R_{0i0j} = 0 = R_{0ijk}$ above).

Finally, requiring $T_{11100}^{(3)} = 0$ (b.w. -1) leads to

$$D^2L_{11} = 0, \quad (\text{B12})$$

and thus, by (A9) (with (B10)),

$$DR_{0101} = 0, \quad (\text{B13})$$

which completes the proof of condition (ii), and the proof is now complete.

Note that, in fact, we have used only up to the third derivatives of \mathbf{F} in the argument above (cf. Remark 1.9).

C VSI_1 and VSI_2 p -forms

A bridge between Corollary 1.4 and Theorem 1.5 is provided by the following result (see also Remark 1.9).

Proposition C.1 (VSI_1 and VSI_2 p -forms). *A p -form \mathbf{F} is VSI_1 iff it is of type N, $\mathcal{L}_\ell \mathbf{F} = 0$, ℓ is Kundt. It is VSI_2 iff it is of type N, $\mathcal{L}_\ell \mathbf{F} = 0$, ℓ is Kundt and (at least) doubly aligned with the Riemann tensor.*

In particular, this means that for solutions of the Einstein-Maxwell theory, we have that $\text{VSI}_1 \Rightarrow \text{VSI}$, since condition 2c of Theorem 1.5 is automatically satisfied (cf. Remark 1.6). Similarly, $\text{VSI}_1 \Rightarrow \text{VSI}$ also for a \mathbf{F} in the classes of spacetimes mentioned in Remark 1.6 which are necessarily degenerate Kundt.

Proof. Let us first prove the “only if” part. We assume that \mathbf{F} is VSI_1 . Then \mathcal{T} must have the same property. In particular, in an adapted frame, we have (B8). Further, requiring $T_{i10}^{(1)} = 0$ (recall definition (B4)) we obtain $L_{i0} = 0$ (i.e., ℓ is geodesic). From now on we can thus employ an affine parameter and a frame parallelly transported along ℓ . The condition $(\nabla \mathbf{F})_{01i_1 \dots i_{p-1}} = 0$ gives $DF_{1i_1 \dots i_{p-1}} = 0$. From $(\nabla \mathbf{F})_{i_1 \dots i_{p+1}} = 0$ and $(\nabla \mathbf{F})_{j10i_1 \dots i_{p-2}} = 0$ we obtain, respectively,

$$F_{1[i_1 \dots i_{p-1} L_{i_p}]j} = 0, \quad (\text{C1})$$

$$F_{1j i_1 \dots i_{p-2} L_{jk}} = 0. \quad (\text{C2})$$

Contracting (C1) with $L_{i_p k}$ and using (C2) leads to $F_{1i_1 \dots i_{p-1} L_{jk} L_{jl}} = 0$. Further contraction with δ_{kl} gives $L_{jk} L_{jk} = 0$ and thus $L_{jk} = 0$, i.e., ℓ is Kundt. This is all we needed to prove as for the VSI_1 statement.

If, additionally, \mathbf{F} is VSI_2 , then from $T_{11i0}^{(2)} = 0$ and $T_{i110}^{(2)} = 0$ we obtain $DL_{1i} = 0 = DL_{i1}$. By (A7) this implies $R_{010i} = 0$, so that (recall also (A6)) ℓ is doubly aligned with the Riemann tensor, as we wanted to prove.

The “if” part of the proposition can be proven similarly by reversing the above steps (essentially showing that under the conditions given in Proposition C.1 the first covariant derivatives (and for VSI_2 also the second covariant derivatives) of \mathbf{F} are b.w. negative). \square

References

- [1] J. L. Synge, *Relativity: the Special Theory*. North-Holland, Amsterdam, 1955.
- [2] R. Penrose and W. Rindler, *Spinors and Space-Time*, vol. 2. Cambridge University Press, Cambridge, 1986.
- [3] H. S. Ruse, *On the geometry of the electromagnetic field in general relativity*, *Proc. London Math. Soc. (2)* **41** (1936) 302–322.
- [4] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*. Cambridge University Press, Cambridge, second ed., 2003.
- [5] E. Schrödinger, *Contributions to Born's new theory of the electromagnetic field*, *Proc. Roy. Soc. London Ser. A* **150** (1935) 465–477.
- [6] A. Coley, R. Milson, V. Pravda, and A. Pravdová, *Vanishing scalar invariant spacetimes in higher dimensions*, *Class. Quantum Grav.* **21** (2004) 5519–5542.
- [7] M. Ortogio, V. Pravda, and A. Pravdová, *Algebraic classification of higher dimensional spacetimes based on null alignment*, *Class. Quantum Grav.* **30** (2013) 013001.
- [8] T. Málek and V. Pravda, *Type III and N solutions to quadratic gravity*, *Phys. Rev. D* **84** (2011) 024047.
- [9] A. Pravdová and V. Pravda, *The Newman-Penrose formalism in higher dimensions: vacuum spacetimes with a non-twisting geodesic multiple Weyl aligned null direction*, *Class. Quantum Grav.* **25** (2008) 235008.
- [10] H. S. Reall, N. Tanahashi, and B. Way, *Causality and hyperbolicity of Lovelock theories*, *Class. Quantum Grav.* **31** (2014) 205005.
- [11] R. Güven, *Plane waves in effective theories of superstrings*, *Phys. Lett. B* **191** (1987) 275–281.
- [12] D. Amati and C. Klimčík, *Nonperturbative computation of the Weyl anomaly for a class of nontrivial backgrounds*, *Phys. Lett. B* **219** (1989) 443–447.
- [13] G. T. Horowitz and A. R. Steif, *Spacetime singularities in string theory*, *Phys. Rev. Lett.* **64** (1990) 260–263.
- [14] G. T. Horowitz, *Singularities in string theory*, in *Strings '90* (R. Arnowitt, R. Bryan, M. J. Duff, D. Nanopoulos, C. N. Pope, and E. Sezgin, eds.), pp. 163–175. World Scientific, Singapore, 1990.
- [15] A. A. Coley, G. W. Gibbons, S. Hervik, and C. N. Pope, *Metrics with vanishing quantum corrections*, *Class. Quantum Grav.* **25** (2008) 145017.
- [16] V. Pravda, A. Pravdová, A. Coley, and R. Milson, *All spacetimes with vanishing curvature invariants*, *Class. Quantum Grav.* **19** (2002) 6213–6236.
- [17] S. Hervik, V. Pravda, and A. Pravdová, *Type III and N universal spacetimes*, *Class. Quantum Grav.* **31** (2014) 215005.
- [18] S. Hervik, T. Málek, V. Pravda, and A. Pravdová, *Type II universal spacetimes*, 1503.08448.
- [19] E. Schrödinger, *A new exact solution in non-linear optics (two-wave-system)*, *Proc. Roy. Irish Acad.* **A49** (1943) 59–66.
- [20] J. Plebański, *Lectures on non-linear electrodynamics*. Nordita, Copenhagen, 1970.
- [21] S. Kichenassamy, *Sur le champ électromagnétique singulier en théorie de Born-Infeld*, *C. R. Hebd. Seanc. Acad. Sci.* **248** (1959) 3690–3692.

- [22] H. Kremer and S. Kichenassamy, *Sur le champ électromagnétique singulier dans une théorie du type Born-Infeld*, *C. R. Hebd. Seanc. Acad. Sci.* **250** (1960) 1192–1194.
- [23] A. Peres, *Nonlinear electrodynamics in general relativity*, *Phys. Rev.* **122** (1961) 273–274.
- [24] F. Bopp, *Eine lineare Theorie des Elektrons*, *Ann. Physik* **430** (1940) 345–384.
- [25] B. Podolsky, *A generalized electrodynamics. Part I – Non-quantum*, *Phys. Rev.* **62** (1942) 68–71.
- [26] C. Teitelboim, *Gauge invariance for extended objects*, *Phys. Lett. B* **167** (1986) 63–68.
- [27] M. Henneaux and C. Teitelboim, *p-form electrodynamics*, *Found. Phys.* **16** (1986) 593–617.
- [28] S. Hervik, *A spacetime not characterized by its invariants is of aligned type II*, *Class. Quantum Grav.* **28** (2011) 215009.
- [29] R. Milson, A. Coley, V. Pravda, and A. Pravdová, *Alignment and algebraically special tensors in Lorentzian geometry*, *Int. J. Geom. Meth. Mod. Phys.* **2** (2005) 41–61.
- [30] L. M. Sokolowski, F. Occhionero, M. Litterio, and L. Amendola, *Classical electromagnetic radiation in multidimensional space-times*, *Ann. Physics* **225** (1993) 1–47.
- [31] G. Hall, *Symmetries and Curvature Structure in General Relativity*. World Scientific, Singapore, 2004.
- [32] N. Pelavas, A. Coley, R. Milson, V. Pravda, and A. Pravdová, *VSI_i spacetimes and the ϵ -property*, *J. Math. Phys.* **46** (2005) 063501.
- [33] M. Durkee, V. Pravda, A. Pravdová, and H. S. Reall, *Generalization of the Geroch-Held-Penrose formalism to higher dimensions*, *Class. Quantum Grav.* **27** (2010) 215010.
- [34] S. Hervik, M. Ortaggio, and L. Wylleman, *Minimal tensors and purely electric or magnetic spacetimes of arbitrary dimension*, *Class. Quantum Grav.* **30** (2013) 165014.
- [35] G. Bergqvist and J. Senovilla, *Null cone preserving maps, causal tensors and algebraic Rainich theory*, *Class. Quantum Grav.* **18** (2001) 5299–5326.
- [36] R. Milson, *Alignment and the classification of Lorentz-signature tensors*, [gr-qc/0411036](#).
- [37] A. Coley, S. Hervik, and N. Pelavas, *Spacetimes characterized by their scalar curvature invariants*, *Class. Quantum Grav.* **26** (2009) 025013.
- [38] A. Coley, S. Hervik, G. O. Papadopoulos, and N. Pelavas, *Kundt spacetimes*, *Class. Quantum Grav.* **26** (2009) 105016.
- [39] V. Pravda, A. Pravdová, A. Coley, and R. Milson, *Bianchi identities in higher dimensions*, *Class. Quantum Grav.* **21** (2004) 2873–2897. See also V. Pravda, A. Pravdová, A. Coley and R. Milson *Class. Quantum Grav.* **24** (2007) 1691 (corrigendum).
- [40] A. Coley, S. Hervik, and N. Pelavas, *Lorentzian manifolds and scalar curvature invariants*, *Class. Quantum Grav.* **27** (2010) 102001.
- [41] A. Coley, A. Fuster, S. Hervik, and N. Pelavas, *Higher dimensional VSI spacetimes*, *Class. Quantum Grav.* **23** (2006) 7431–7444.
- [42] J. Podolský and M. Žofka, *General Kundt spacetimes in higher dimensions*, *Class. Quantum Grav.* **26** (2009) 105008.
- [43] I. Robinson, *Null electromagnetic fields*, *J. Math. Phys.* **2** (1961) 290.
- [44] T. Ortín, *Gravity and strings*. Cambridge University Press, Cambridge, second ed., 2015.

- [45] J. M. Figueroa-O’Farrill and G. Papadopoulos, *Homogeneous fluxes, branes and a maximally supersymmetric solution of M-theory*, *JHEP* **0108** (2001) 036.
- [46] J. F. Schonfeld, *A mass term for three-dimensional gauge fields*, *Nucl. Phys. B* **185** (1981) 157–171.
- [47] S. Deser, R. Jackiw, and S. Templeton, *Topologically massive gauge theories*, *Ann. Phys.* **140** (1982) 372–411. See also S. Deser and R. Jackiw (1988), Erratum, *Ann. Phys.* 185:406.
- [48] O. D. Andreev and A. A. Tseytlin, *Partition function representation for the open superstring effective action: Cancellation of Möbius infinities and derivative corrections to Born-Infeld lagrangian*, *Nucl. Phys. B* **311** (1988) 205–252.
- [49] L. Thorlacius, *Born-Infeld string as a boundary conformal field theory*, *Phys. Rev. Lett.* **80** (1998) 1588–1590.
- [50] W. Chemissany, R. Kallosh, and T. Ortin, *Born-Infeld with higher derivatives*, *Phys. Rev. D* **85** (2012) 046002.
- [51] J. Bičák and J. Slavík, *Non-linear electrodynamics in the Newman-Penrose formalism*, *Acta Phys. Polon.* **B6** (1975) 489–508.
- [52] P. Krtouš, J. Podolský, A. Zelnikov, and H. Kadlecová, *Higher-dimensional Kundt waves and gyratons*, *Phys. Rev. D* **86** (2012) 044039.
- [53] A. García Díaz and J. F. Plebański, *All nontwisting N’s with cosmological constant*, *J. Math. Phys.* **22** (1981) 2655–2658.
- [54] I. Ozsváth, I. Robinson, and K. Rózga, *Plane-fronted gravitational and electromagnetic waves in spaces with cosmological constant*, *J. Math. Phys.* **26** (1985) 1755–1761.
- [55] J. Bičák and J. Podolský, *Gravitational waves in vacuum spacetimes with cosmological constant. I. Classification and geometrical properties of nontwisting type N solutions*, *J. Math. Phys.* **40** (1999) 4495–4505.
- [56] J. B. Griffiths and J. Podolský, *Exact Space-Times in Einstein’s General Relativity*. Cambridge University Press, Cambridge, 2009.
- [57] Y. N. Obukhov, *Generalized plane-fronted gravitational waves in any dimension*, *Phys. Rev. D* **69** (2004) 024013.
- [58] J. Podolský and M. Ortaggio, *Explicit Kundt type II and N solutions as gravitational waves in various type D and O universes*, *Class. Quantum Grav.* **20** (2003) 1685–1701.
- [59] J. Griffiths, P. Docherty, and J. Podolský, *Generalized Kundt waves and their physical interpretation*, *Class. Quantum Grav.* **21** (2004) 207–222.
- [60] M. M. Caldarelli, D. Klemm, and E. Zorzan, *Supersymmetric gyratons in five dimensions*, *Class. Quantum Grav.* **24** (2007) 1341–1358.
- [61] A. Coley, A. Fuster, S. Hervik, and N. Pelavas, *Vanishing scalar invariant spacetimes in supergravity*, *JHEP* **0705** (2007) 032.
- [62] M. Ortaggio, V. Pravda, and A. Pravdová, *Higher dimensional Kerr-Schild spacetimes*, *Class. Quantum Grav.* **26** (2009) 025008.
- [63] T. Málek, *Extended Kerr-Schild spacetimes: general properties and some explicit examples*, *Class. Quantum Grav.* **31** (2014) 185013.
- [64] A. A. Coley, *A class of exact classical solutions to string theory*, *Phys. Rev. Lett.* **89** (2002) 281601.

- [65] J. Kowalski-Glikman, *Vacuum states in supersymmetric Kaluza-Klein theory*, *Phys. Lett. B* **134** (1984) 194–196.
- [66] C. Hull, *Exact pp-wave solutions of 11-dimensional supergravity*, *Phys. Lett. B* **139** (1984) 39.
- [67] A. A. Tseytlin, *String vacuum backgrounds with covariantly constant null Killing vector and two-dimensional quantum gravity*, *Nucl. Phys. B* **390** (1993) 153–172.
- [68] E. Bergshoeff, R. Kallosh, and T. Ortin, *Supersymmetric string waves*, *Phys. Rev. D* **47** (1993) 5444–5452.
- [69] V. P. Frolov and A. Zelnikov, *Gravitational field of charged gyratons*, *Class. Quantum Grav.* **23** (2006) 2119–2128.
- [70] V. P. Frolov and F.-L. Lin, *String gyratons in supergravity*, *Phys. Rev. D* **73** (2006) 104028.
- [71] K. P. Tod, *All metrics admitting super-covariantly constant spinors*, *Phys. Lett. B* **121** (1983) 241–244.
- [72] J. P. Gauntlett, J. B. Gutowski, C. M. Hull, S. Pakis, and H. S. Reall, *All supersymmetric solutions of minimal supergravity in five dimensions*, *Class. Quantum Grav.* **20** (2003) 4587–4634.
- [73] H. W. Brinkmann, *Einstein spaces which are mapped conformally on each other*, *Math. Ann.* **94** (1925) 119–145.
- [74] M. Ortaggio, *Bel-Debever criteria for the classification of the Weyl tensor in higher dimensions*, *Class. Quantum Grav.* **26** (2009) 195015.
- [75] T. Dereli and M. Gürses, *The generalized Kerr-Schild transform in eleven-dimensional supergravity*, *Phys. Lett. B* **171** (1986) 209–211.
- [76] M. Ortaggio, V. Pravda, and A. Pravdová, *Ricci identities in higher dimensions*, *Class. Quantum Grav.* **24** (2007) 1657–1664.